

**ENVIRONMENTAL AND SOCIAL IMPACT ASSESSMENT STUDY
FOR
SIERRA TROPICAL LTD's AGRICULTURAL PROJECT IN LUGBU CHIEFDOM BO
DISTRICT**

EXECUTIVE SUMMARY & MAIN REPORT

Prepared by

CEMMATS Group Ltd



Freetown, Sierra Leone

on behalf of:

**SIERRA TROPICAL
LIMITED (STL)**

**September
2016**

DOCUMENT HISTORY

Version History	Date	Reviewer
-----------------	------	----------

Title	Environmental and Social Impact Assessment for the Sierra Tropical Ltd Agricultural Project: Executive Summary and Main Report
Authors	Anthony Mansaray; Arnold Okoni-Williams; Bartholomew Bockarie; Joe Lappia; Leonard B. Buckle; Ralph Bona; Rashidu Sinnah; Vanessa James
Date	September 2016
Subject	Environmental and Social Impact Assessment
Publisher	CEMMATS Group Ltd
Type	Client Report
Description	ESIA for the Sierra Tropical Ltd
Contributors	Joseph Gbassa; Josephine Turay; Mariama Jalloh
Format	Microsoft™ Word 2007
Source	Text
Rights	© CEMMATS Group Ltd
Identifier	
Language	English
Relation	
Coverage	Sierra Leone, 2016

CONSULTANT'S DISCLAIMER

CEMMATS Group Ltd (hereafter, 'CEMMATS') has prepared this Environmental and Social Impact Assessment (ESIA) Report for the sole use of the Client and for the intended purposes as stated in the Contract between the Client and CEMMATS under which this work was completed. This ESIA Report may not be relied upon by any other party without the express written agreement of CEMMATS and/or the Client.

CEMMATS has exercised due and customary care in conducting this ESIA but has not, save as specifically stated, independently verified information provided by others. No other warranty, expressed or implied is made in relation to the conduct of the ESIA or the contents of this Report. Therefore, CEMMATS assumes no liability for any loss resulting from errors, omissions or misrepresentations made by others. This Report has been prepared at the request of the Client. The use of this Report by unauthorised third parties without written authorisation from CEMMATS shall be at their own risk, and CEMMATS accepts no duty of care to any such third party.

Any recommendations, opinions or findings stated in this Report are based on circumstances and facts as they existed at the time CEMMATS performed the work. Any changes in such circumstances and facts upon which this Report is based may adversely affect any recommendations, opinions or findings contained in this Report.

No part of this Report may be copied or duplicated without the express written permission of the Client and/or CEMMATS. Where field investigations have been carried out, these have been restricted to a level of detail required to achieve the stated objectives of the work referred to in the Contract. This work has been undertaken in accordance with CEMMATS' Quality System.

Signed by:



Andrew Keili

CEMMATS Group Ltd
Beyoh House
7A Cantonment Road
Off King Harman Road
Brookfields
Freetown
Sierra Leone
Email: akeili@cemmatssl.com
Tel: +232 76602174
Website: www.cemmatssl.com

ACKNOWLEDGEMENTS

We wish to extend our appreciation to various government of Sierra Leone (GOSL) ministries, departments, agencies, institutions, statutory bodies, organisations and individuals whose assistance, either directly or indirectly, made the Environmental and Social Impact Assessment (ESIA) study possible. We are particularly grateful to the assistance availed to us by the Environment Protection Agency Sierra Leone (EPA-SL), especially for providing us with useful take off guidance to carry out the ESIA study.

We acknowledge the assistance and cooperation of Sierra Tropical Limited, who availed us relevant documents and information, to successfully carry out this study.

We are thankful to the local authorities and members of the various communities visited in the Lugbu Chiefdom for their diverse assistance during the main field visit.

Last but not least, we would like to thank all those who, in one way or another contributed to making this ESIA study successful.

TABLE OF CONTENTS

DOCUMENT HISTORY	ii
ACKNOWLEDGEMENTS	iv
TABLE OF CONTENTS	v
LIST OF FIGURES	ix
LIST OF TABLES	x
LIST OF ACRONYMS	xii
GLOSSARY	xiv
EXECUTIVE SUMMARY	16
1 INTRODUCTION	40
1.1 Project Background	40
1.2 National Perspective	40
1.3 Project Area	41
1.3.1 Brief History of Lugbu Chiefdom	42
1.4 Environmental and Social Impact Assessment Process	42
1.4.1 Stages of the ESIA Process	42
1.4.2 Purpose of the ESIA Study	43
1.4.3 Objectives of the ESIA Study	43
1.4.4 ESIA Consultants and Teams	44
1.5 Description of the Terms of Reference (TOR)	45
1.6 Scope of Work	53
1.7 Assumptions and Limitations of Study	53
1.8 ESIA Study Boundaries	53
1.9 Organisation of the ESIA Report(s)	54
1.9.1 The ESIA Report	54
1.9.2 Management Plans featured in the ESMP	55
2 PROJECT DESCRIPTION	57
2.1 Specific Project Components	62
2.1.1 Phase 1	62
2.1.2 Phases 2 and 3	78
2.2 Timescale of Operations	80
3 ANALYSES OF PROJECT ALTERNATIVES	81
3.1 The “No Project Option”	81
3.2 The Project Site	82
3.3 Project Technology & Operations Option	82
4 POLICY, LEGAL, REGULATORY and INSTITUTIONAL CONTEXT	84
4.1 Policies and Plans	84

4.1.1	National Environmental Policy, 1994.....	84
4.1.2	Agricultural Policy.....	85
4.1.3	Draft National Lands Policy, 2015	85
4.1.4	Forestry Policy, 2010.....	86
4.1.5	Conservation and Wildlife Policy, 2010.....	87
4.1.6	Disaster Management Preparedness Plan, 2006	87
4.2	Legislation.....	88
4.2.1	Environment Protection Agency Act, 2008/2010.....	88
4.2.2	The Forestry Act, 1988	89
4.2.3	Land Tenure and Ownership.....	89
4.2.4	Fisheries Act, 2007	90
4.2.5	Wildlife Conservation Amendment Act, 1990	90
4.2.6	Factories Act – 1974	90
4.2.7	Local Government Act, 2004.....	93
4.3	Regulations.....	94
4.3.1	Forestry Regulations, 1990	94
4.3.2	Fisheries Regulations	94
4.3.3	Draft Wildlife Regulation, 1997	95
4.4	Institutional Context.....	95
4.4.1	Ministry of Water Resources	95
4.4.2	Ministry of Fisheries and Marine Resources	95
4.4.3	Ministry of Agriculture, Forestry and Food Security	96
4.4.4	Ministry of Lands, Country Planning and the Environment	96
4.4.5	EPA-SL.....	96
4.5	International Conventions Policies, Codes, Protocols and Guidelines	97
4.5.1	International Conventions.....	97
4.5.2	International Lending Institutions Policies, Standards and Guidelines	101
5	BASELINE SURVEY AND CONDITION	103
5.1	Physical Environment	105
5.1.1	Climate.....	105
5.1.2	Air Quality and Dust.....	109
5.1.3	Noise	111
5.1.4	Geology.....	112
5.1.5	Landform.....	113
5.1.6	Soils.....	113
5.1.7	Hydrology	123
5.2	Biological environment.....	146
5.2.1	Ecology	146
5.2.2	Creation of Buffer Zones	156

5.2.3	High Conservation Value Assessment.....	159
6	DESCRIPTION OF THE SOCIAL ENVIRONMENT.....	161
6.1	Introduction	161
6.1.1	National Socio-Political Context	161
6.1.2	Local Governance Structure	163
6.1.3	Local Regional Socio-Economic Context.....	163
6.1.4	Socio-Economic Condition of Project Area (Lugbu Chiefdom)	168
6.2	Socio-economic Status and Living Conditions	169
6.2.1	Methodology Used for Socio-Economic Baseline Survey	169
6.3	Results of Questionnaires administered to Landholders	172
6.3.1	Status of Respondents	172
6.3.2	Gender of Landholder	173
6.3.3	Marital Status of Landholder	174
6.3.4	Number of households living in a dwelling unit.....	175
6.3.5	Educational Level of Landholders	175
6.3.6	Size of Landholders' Land.....	176
6.3.7	Availability of Sufficient Land for food production.....	177
6.3.8	Income from Land Cultivation.....	177
6.3.9	Proportion of upland to be leased to Sierra Tropical	178
6.3.10	Sufficiency of Remaining Land for Family's Food Cultivation Requirements 178	
6.3.11	Preferred Land Lease Payment Method.....	179
6.3.12	Intended use of lease rent.....	179
6.3.13	Main Occupation of Landholder	180
6.3.14	Secondary Occupation of Landholder	181
6.3.15	Landholders' Awareness and Perception of STL Project.....	182
6.3.16	Perceived Impacts of Project	183
6.3.17	Food Security	184
6.3.18	Health.....	186
6.3.19	Education	187
6.3.20	Water and Sanitation.....	190
7	IDENTIFICATION OF POTENTIAL IMPACTS	193
7.1	Introduction	193
7.2	Environmental and Social Impact Assessment	193
7.2.1	Methodology	193
8	CONCLUSION	225
8.1	Summary	225
8.1.1	Components of the ESIA	225
8.1.2	Key Assessment Findings	226

8.2	Conclusion.....	226
9	REFERENCES	228
10	APPENDICES	230

LIST OF FIGURES

Figure 1.9-1: Proposed Location of STL Project Features Spanning 1 st , 2 nd and 3 rd Phases of the Project	58
Figure 2.1-1: Field showing Natural contours of a selected field.....	62
Figure 2.1-2: Water flow and natural drainage pattern.....	63
Figure 2.1-3: The field layout is adapted to the relief. The center road follows the watershed and the orientation of the field blocks is approximately parallel to the contour lines.	64
Figure 2.1-4: Pineapple Farm Plots (Previously owned by ALA).....	67
Figure 5.1-1: Map of Water Quality Sampling Sites at or in the vicinity of the Project site.	131
Figure 5.1-2: Location in the Sewa River at Heima where water is fetched for drinking and domestic use	137
Figure 5.1-3: Sampling at Heima Village	137
Figure 5.1-4: Hand-pump well in Kormende.....	138
Figure 6.3-1: Status of Respondent.....	173
Figure 6.3-2: Gender of Landholders.....	173
Figure 6.3-3: Age Distribution of Landholders	174
Figure 6.3-4: Marital Status of Landholder	174
Figure 6.3-5: Number of Wives per Male Landholder	175
Figure 6.3-6: Number of households per dwelling unit.....	175
Figure 6.3-7: Educational Level of Landholders	176
Figure 6.3-8: Size of Land Sufficient for Food Production Requirements.....	177
Figure 6.3-9: Income Generated from Land Cultivation	177
Figure 6.3-10: Proportion of upland land to be leased to Sierra Tropical	178
Figure 6.3-11: Sufficiency of Remaining Land for Food Cultivation Requirements	179
Figure 6.3-12: Intended use of lease rent to landholder.....	180
Figure 6.3-13: Occupation of Landholders	180
Figure 6.3-14: Secondary Occupations.....	181
Figure 6.3-15: Income from Secondary Occupation.....	182
Figure 6.3-16: Perception about Project	183
Figure 6.3-17: Perceived Positive Outcomes of Project	183
Figure 6.3-18: Experience of Food Shortage	185
Figure 6.3-19: Illnesses experienced by Landholders' Families	187
Figure 6.3-20: Sources of Health Care	187
Figure 6.3-21: Children of Primary School age (6 – 13) in Landholders' Households.....	188
Figure 6.3-22: Children in Landholders' Households of Primary School Age, Enrolled in School	188
Figure 6.3-23: Ease of Meeting Financial Requirements of School	189
Figure 6.3-24: Landholders with Children of Secondary School Age in their Households ..	189
Figure 6.3-25: Children of Secondary School Age attending School.....	190
Figure 6.3-26: Households' Primary Water Sources	191
Figure 6.3-27: Types of Toilet Facilities or Alternatives	192
Figure 6.3-28: Waste Disposal Methods.....	192

LIST OF TABLES

Table 1.3-1: Lugbu Chiefdom.....	42
Table 1.4-1 ESIA Consultants and Teams.....	44
Table 1.5-1 Terms of Reference of the ESIA Study.....	46
Table 2.1-1: Land Preparation Flow Diagram.....	65
Table 2.1-2: Fertilizers.....	73
Table 2.1-3: Pesticides.....	74
Table 2.1-4: Plant Growth Regulators.....	75
Table 2.1-5: Post Harvest Chemicals.....	75
Table 4.5-1 : List of International Conventions and Agreements to which Sierra Leone is Party	97
Table 5-4.5-1: Physical Site measurement points.....	104
Table 5.1-1: Summary of Some Climatic Data for Bo	106
Table 5.1-2: Annual Water Budgets (mm) for the Transitional Agro-Climatic Region and for BO	107
Table 5.1-3: Wind speed measurements for settlements within and around the project area	108
Table 5.1-4: Dust level within and around the project area	110
Table 5.1-5: Noise level data for settlements within and around project area.....	111
Table 5.1-6: Typical representatives of Soil/Landform Physical Properties in Sierra Tropical Limited Project Area.....	116
Table 5.1-10: Sensitive Receptors	125
Table 5.1-11: Calculation of Monthly Evapotranspiration Using Thornthwaite Method for Sierra Leone	126
Table 5.1-12: Calculation of the Monthly Evt using the Blaney & Criddle Method for Sierra Leone.....	127
Table 5.1-13: Monthly Water Balance for Sierra Leone using average data and ETP by Thornthwaite method	128
Table 5.1-14: Numerical Standards	130
Table 5.1-16: Sample Chemical Data	136
Table 5.2-1: Site number, names, GPS data and comments of areas visited during the survey.	147
Table 5.2-2: Google Earth (February 2016) image of part of the main areas and photos of the vegetation and landscape features of the Sierra Tropical Limited proposed concession zone in the Lugbu Chiefdom, in the Bo District.....	150
Table 5.2-3: Tree species of IUCN conservation status observed forest patches and gallery forests within the project's concession zones	151
Table 5.2-4: Amphibians and reptile species encountered, their IUCN status and habitat distribution	153
Table 5.2-5: Sites identified for the creation of buffer zones within concession zones. The sites are marked by yellow polygons.	157
Table 5.2-6: Sites identified for buffer zone establishment within the Sierra Tropical Limited Concession zone.....	157

Table 6.1-1: Information on National Social Indicators	162
Table 6.1-2: Population of Lugbu Chiefdom	169
Table 6.1-3: Social Facilities in Lugbu	169
Table 6.3-2: Preferred Lease Rent Payment Method	179
Table 6.3-3: Income from Primary Occupation	180
Table 6.3-4: Landholder's Awareness of Project	182
Table 6.3-5: Perceived Negative Outcomes of Project	183
Table 6.3-6: Proposed Remedies for Perceived Negative	184
Table 6.3-8: Coping Strategies during Food Shortages	186
Table 6.3-9: Reasons why children of primary school age are not in School	188
Table 7.2-2: Environmental and Social significance scale	194
Table 7.2-3: Impact Assessment Scale	194
Table 7.2-4: Degree of Difficulty to Mitigate	194
Table 7.2-5: Categories of Impact	195
Table 7.2-6: Land Preparation and Construction Phase - Environmental and Mitigation Measures	197
Table 7.2-7: Land Preparation and Construction Phase - Social Impacts and Mitigation Measures	202
Table 7.2-8: Operational Phase – Environmental Impacts and Mitigation Measures	209
Table 7.2-9: Operations Phase - Social Impacts and Mitigation Measures	216
Table 7.2-10: Decommissioning Phase – Environmental and social impacts	221

LIST OF ACRONYMS

⁰ C	Degrees Celsius
%	Percentage
"	Inch
Al	Aluminium
AMSL	above mean sea level
CBD	Convention on Biodiversity
CBO	community-based organisation
CDAP	Community Development Action Plan
CEMMATS	Construction Engineering Maintenance, Manufacturing and Technical Services
CI	Corrugated Iron
CITES	Convention on International Trade in Endangered Species on wild flora and fauna
Cl	Chloride
cm	centimetre
Cm ²	Square centimetre
dB	decibels
DO	Dissolved Oxygen
EC	Electrical Conductivity
EPA-SL	Environment Protection Agency – Sierra Leone
ESIA	Environmental and Social Impact Assessment
ESMP	Environmental and Social Management Plan
ERP	Emergency Response Plan
FAO	Food and Agricultural Organization
GDP	Gross Domestic Product
GIS	Geographic Information Systems
GoSL	Government of Sierra Leone
GPS	Global Positioning System
HC	Hydrocarbons
HDI	Human Development Index
IFC	International Finance Corporation
IMR	Infant Mortality Rate
IVS	Inland Valley Swamp
JSS	Junior Secondary School
K+	Potassium ions
kg	kilogram
km	kilometre
Km ²	Square kilometre
Le	Leones
m	metre
MDA	Ministries, Departments and Agencies
MFIs	micro-finance institutions
mg	milligram

mg/L	Milligram per litre
mg/m ³	Milligram per cubic metre
mm	millimetre
m/s	Metre per second
N	North
NE	North-east
NGO	Non-Governmental Organization
PAC	Project Affected Communities
PAPs	Project Affected Persons
PCDP	Public Consultation and Disclosure Plan
PM	Particulate Matter
PRSP	Poverty reduction Strategy Paper
RH	Relative Humidity
RPF	Resettlement Policy Framework
Si	Silicon
t	tons
TOR	Terms of Reference
WMP	Waste Management Plan

GLOSSARY

Aquatic Ecosystem	An aquatic area where living and non-living elements of the environment interact. This includes the physical, chemical, and biological processes and characteristics of rivers, lakes, and wetlands and the plants and animals associated with them
Avifauna	Birds
Board of EPA-SL	This is a board of directors that form the governing body of EPA-SL; it is headed by the Executive Chairperson and consists of mainly representatives of a number of line ministries.
Client	Individual or organization which uses the services or advice of a professional person or organization
Community	A group of interacting families or households, living in some proximity (i.e., in space, time, or relationship) that shares common values and has social cohesion and is recognized as 'separate' by other similar groups.
Community Development Action Plan (CDAP):	A CDAP is an action plan to address key community issues that are based on the expressed development needs and aspirations of the local residents of a community.
Conservation	<p>The planning, management, and implementation of an activity with the objective of protecting the essential physical, chemical, and biological characteristics of the environment against degradation;</p> <p>The process of managing biological resources (e.g., timber, fish) to ensure replacement by re-growth or reproduction of the part harvested before another harvest occurs. A balance between economic growth and environmental and natural resource protection.</p>
Ecosystem	A community of interdependent organisms together with the environment they inhabit and with which they interact
Environmental and social management plan:	A plan of action for the management of impacts on the environment and human settlements as a result of the project, as well as maintaining compliance with relevant legislature
EPA-SL "checklist"	A list of procedures developed and provided by EPA-SL to be systematically followed by a client for the conduct of ESIA and the issuance of an EIA license

Human Development Index:	The Human Development Index (HDI) is a composite statistic used by the UN to rank countries by level of "human development"
pH	A measure of the intensity of the acid or base chemistry of the water. A pH of 7 is neutral, while below 7 is acidic and above 7 is basic. pH in surface water is regulated by the geology and geochemistry of an area and is affected by biological activity. The distribution of aquatic organisms and the toxicity of some common pollutants are strongly affected by pH
Potable Water	Water that is suitable for human consumption
Project Affected Persons	Any person who, as a result of the implementation of a project, loses the right to own, use, or otherwise benefit from a built structure, land (residential, agricultural, or pasture), annual or perennial crops and trees, or any other fixed or moveable asset, either in full or in part, permanently or temporarily. Also, includes any person whose use of, and access to, natural resources is lost or restricted with an expected adverse effect on their livelihoods
Project Proponent	An individual or organization having responsibility for acquiring all necessary consents, and if successful, implementing a project
RAMSAR	Convention on wetlands of international importance
Runoff	Water that moves across (or through) soils on the land during snowmelt or rainstorms
Socio-economic data/study	Data or study to determine and describe social and economic factors characteristic of an area, such as a District, or defined population entity such as a community.
Social Indicators	A direct and valid statistical measure often used to monitor levels and changes over time of specified social factors such as average family size
Stakeholders	Individuals, groups, organizations, and institutions interested in and/or potentially affected by a project and/or having the ability to influence a project

EXECUTIVE SUMMARY

Introduction

Sierra Tropical Ltd (STL) proposes to embark on an agro-processing project involving large scale planting and processing of tropical fruits in Sierra Leone using approximately 15,000 hectares. It is envisioned to be fully integrated from cultivating, growing and harvesting pineapple, mango, papaya and other tropical fruit and processing them into manufactured products. The project is proposed to be implemented in Sierra Leone as it has the necessary favorable conditions, such as ideal weather, plentiful land, duty concessions, and a business friendly government and administration particularly with regards agriculture.

The project will initially grow up to 4,335 hectares of leased land in Lugbu Chiefdom, Bo District, and will be implemented in phases, starting with the development of a pineapple nursery, spreading out in phases to eventually cover the maximum area desired and possible.

If the first phase of the project is successful and the quality/productivity of the fruits meet our requirements, a manufacturing facility will be constructed in the later stages of the second phase, to process the raw materials into manufactured products in various packages of cans, drums, plastics and boxes.

Environmental and Social Impact Assessment Process

Prior to commencement of any project that may affect the environment and communities, it is mandated by legislation that an Environmental and Social Impact Assessment (ESIA) study be done, and, upon approval by EPA-SL, a licence is secured.

The Sierra Leone Environment Protection Agency Act, (SLEPAA) 2008 and the EIA Supplementary Acts, 2010 describe the requirements and process for securing an EIA licence, which is laid out in a “checklist” prepared by EPA-SL. In short, the client first applies to the local regulatory body, Environment Protection Agency, Sierra Leone (EPA-SL) for an EIA licence. EPA-SL requires that a screening form be filled and submitted with the application letter, after which a decision is made on the category of the project; this is followed by a scoping report. EPA-SL will then decide on the terms of reference (TOR) to be drafted by the project proponent or an independent consultant hired by the proponent.

On the approval of the agency, the consultant carries out an assessment of the environmental and social impacts of their planned operations on ecosystems and communities in the project area. A report is prepared at the end of the study and submitted to EPA-SL for review. If approved, the proponent will then be requested to conduct public disclosure meetings with relevant stakeholders on the findings and recommendations of the study, and incorporate comments, suggestions and requests made during those meetings into a public consultation and disclosure report. Finally, all reports pertaining to the ESIA study are then forwarded to the Board of EPA-SL for a decision to be made on the issue of the licence.

Project Description

Through Sierra Tropical Ltd., Dole Asia Holdings will establish a new agro-processing project in Sierra Leone. It has elected to set up its project in Sierra Leone because there are the necessary favourable conditions, such as ideal weather, plentiful land, duty concessions, business friendly government and administration especially to agriculture.

The project is envisioned to be fully integrated from cultivating and marketing fresh pineapple papaya and tropical fruits as well as processing those fruits into manufactured products in various packages of cans, drums, plastics and boxes, thereby adding further value to the product. The manufacturing aspect of the project will involve the construction and installation of a cannery and a canning plant. The finished products will be exported to Europe, United States, Middle East and Africa, as product of Sierra Leone, primarily under the Dole label.

The acquisition of the entire 15,000 hectares required for the project will be done gradually, over a series of phases. Phase 1 will last approximately 2 years, and will utilise a minimum of 800 hectares. The second phase will be implemented in the third year and will last for 3 years, with a proposed expansion to a minimum of 4,000 hectares. For years 6 to 10, more than 2,000 hectares roughly may be acquired, and with the proven success of Phase 2, expansion will continue to increase steadily. Using this projection, and assuming success at each stage, the project is estimated to continue for over well over 50 years.

Phase 1—Technical Feasibility Study

Phase 1 will be a technical feasibility study phase in order to confirm the viability of establishing a pineapple plantation in the areas identified for the project site. While the Company estimates completion of this phase approximately two years from commencement, the company may need to extend this period in order to fully accomplish its feasibility study.

The main focus of Phase 1 in addition to initial commercial planting of pineapple is to propagate planting materials through development of a nursery.

Phase 2 -First Expansion

If Sierra Tropical Ltd. determines to continue with the project following Phase 1, it then intends to further expand its operations. In this phase, the company intends to lease more land and increase plantation capability. Furthermore, the Company intends to establish a cannery and also install additional equipment for IQF Frozen products and other processes for the expansion of this project.

Phase 3 and beyond —Long Term Goal

Sierra Tropical Limited intends to expand its operations further in this phase, depending on the success of Phases 1 and 2. The Company's long term goal is to develop and establish a large scale agriculture and manufacturing operation in Sierra Leone. It aims to be the major

exporter of high quality and competitive pineapple and other tropical fruit products in the major markets in Europe, United States, Middle East and Africa. It also aims to be a major source of employment in the country, continuously developing and improving the skills of Sierra Leoneans.

Policy, Legal, Regulatory and Institutional Context

Sierra Leonean policy, legal and regulatory requirements are outlined to ensure that all project related requirements are taken into full consideration throughout the project.

The following was assessed:

- Policies and Plans:
 - National Environmental Policy (1994);
 - Agriculture Policy (2009);
 - National Land Policy (2015);
 - Forest Policy (2010);
 - National Biodiversity Strategy and Action Plan (2003);
 - Conservation and Wildlife Policy (2010);
 - Disaster Management Preparedness Plan (2006).
- Legislation
 - *Sierra Leone Environment Protection Agency Act (2008 / 2010)*;
 - *The Forestry Act (1988)*;
 - *Land Tenure and Ownership*
 - *Fisheries Act (2007)*;
 - *Wildlife Conservation Amendment Act (1990)*;
 - *Factories Act (1974)*
 - *Local Government Act (2004)*.
- Institutional Context:
 - The Ministry of Water Resources
 - The Ministry of Fisheries and Marine Resources
 - The Ministry of Agriculture, Forestry and Food Security

- Ministry of Lands, Country Planning and the Environment
- The Environment Protection Agency Sierra Leone

Baseline Survey and Condition

The baseline assessment was carried out on the physical, biological and social environments. Descriptions of the existing environment include primary and secondary data and information from relevant and available sources;

Physical Environment

Climate

Information and climatic data relating directly to the Freetown ports area was available from the Sierra Leone Meteorological Department. Field measurements, specifically wind speed, were used in conjunction with secondary data obtained, to represent climatic conditions within the project area and its surroundings. Climatic and other data for Bo area have been used to represent conditions within the project area.

Summary of Some Climatic Data for Bo

Climatic Variable		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Monthly Mean Temp (°C) (1976-1980)	Max	31.9	33.6	33.9	33.1	31.8	30.3	28.5	27.9	29.3	30.7	31.0	30.
	Min	20.1	20.8	21.6	22.2	22.4	20.0	21.8	21.2	21.8	21.5	21.7	20.8
Average Relative Humidity (%) (1976 – 2005)	9 a.m.	78.7	79.1	78.1	78.9	81.4	86.0	89.4	91.1	88.7	86.3	83.6	79.8
	3 p.m.	62.4	63.4	64.0	65.9	70.5	75.5	81.0	82.8	78.8	76.0	74.1	67.9
Monthly Means of Rainfall (mm) (2007)		4.3	5.2	46.8	93.7	176.5	446.3	649.1	637	445.6	290.4	125.2	12.1

Air Quality and Dust

The quantity of dust particles (PM10) in the air was recorded within and around twenty one (21) settlements in the study area.

From the analysis of the data collected the dust levels at the time of the study were so low that the dust probe barely recorded levels of particulate matter. This was due to the period during which the measurements were taken (rainy season), with rainfall occurring during the time of data collection.

The WHO threshold for PM10 is 50µg/m³; the maximum values recorded range between 0.001mg/m³ (1µg/m³) and 0.011 mg/m³ (11 µg/m³) while the average values is 0.00 mg/m³ (0 µg/m³) in all the settlements. All values are within the WHO recommended standards indicating that the settlements have an environmentally friendly atmosphere during the rainy season which permits healthy living conditions.

Noise

Noise levels were measured and recorded within and around twenty one (21) settlements in the project area.

The noise levels ranged from 38 decibels (dB) in Momandu to 71 decibels (dB) in Garinga. Levels were influenced by vehicular movements and ambient noise of normal community activities, normally referred to as community noise. Noise levels in some of the communities measured exceed 55db which is the average community noise level threshold. In terms of industrial environment (which the project area will soon become), noise levels are within the World Health Organisation (WHO) recommended limits, which stipulates that hearing protection gear must be used for noise exposure levels above 85 dB.

Geology

The project area geological study was done mostly through desk studies on information gathered applicable to the site and its general surroundings.

The Project area geology is within the Basement Granite and Acid Gneiss Terrain of the geology of Sierra Leone as it stretches along the Sewa River alluvial diamond field and covers approximately 7.5 to 17 km of reach to the diamondiferous Sewa River.

Landform

There are four main landforms occurring within the project area which is described below:

- **Isolated hills** - short, straight to convex, gentle to moderately sloping.
- **Interfluves** - medium length undulating, very gentle to gentle sloping.
- **Valley swamps** - level to nearly level with varying width and locally channelled
- **Floodplains** - level to almost flat with varying width and locally channelled and terraced.

Soils

An exploratory Soil Survey, Water and Climate Assessment of Sierra Leone was conducted by Dr. Carlos Gauggel, Mr. Carlos Mandujano and Mr. Romeo Patricio in 2011. The assessment resulted in the following conclusions:

1. There are about 2.8 million ha of suitable soils for pineapple production in the Lower and Inner Coastal Plains of Sierra Leone out of which 1.8 million can be irrigated using the major rivers and very close to roads and townships.
2. In general, in this study, soil quality for pineapple production, as the first priority, has been ranked A, B, B2 and C in this report. Availability for water has been considered

for this ranking. The available number of ha per zone is as follows : Area A 630000 ha, Area B 720000 ha, Area B2 360000 ha, Area C 180000 ha

3. Irrigation has to be implemented from December to May, river water and water reservoirs are the main source of water for irrigation. The number of ha indicated above have access to irrigation.
4. Primary roads and bridges are in very good shape and can handle containers. Second grade roads are very varied in condition, the condition of unpaved roads is a major issue in the rainy season.
5. Surface drainage and sub-surface drainage, in some areas, are a must for successful pineapple production due to heavy rains during the rainy season.
6. All soils will require an initial cross-way deep sub soiling to loosen the soil matrix and to break up any laterite or iron-stone layers, followed by disk harrowing, this is a must for the success of the project and it is probably required once (deep sub soiling).
7. Due to water availability, sloping landscapes, and the high risk of soil erosion, drip or low discharge irrigation is the best option.
8. The low temperatures (December-March), limited sun-light hours during some months (July-August) constitute the major constraint for pineapple production in Sierra Leone. Soil and water-wise it is very good area for pineapple cultivation with vast land and water resources available.

Hydrology

Sierra Leone falls within a tropical climate zone with peculiar wet and dry seasons splitting the year in equal halves:

- ❑ Rainy Season (May –November): Thunderstorms and squalls and Steady rains;
- ❑ Dry season (December – April): Dry weather with high humidity and Dry weather with low humidity.

These seasons could be subdivided into four (4) distinct types of weather. (i) Heavy squalls and thunderstorms, (ii) steady rains (iii) Harmattan season characterised by dry and windy weather with low humidity and (iv) Hot and humid weather.

Sierra Leone has a complex drainage pattern that includes numerous rivers and smaller creeks and streams. The larger rivers originate in the Fouta Djallon highlands of Guinea and generally flow from northeast to southwest across Sierra Leone. The country's drainage system includes nine major rivers and minor coastal creeks and tidal streams. The river watersheds range in size from 14,140 km² for the Sewa River to less than 385 km² for the smaller watersheds.

Field investigations were conducted within the study area and its immediate environs to obtain water quality data. This involved identifying water points and channels within the study area. The methodology used for the study includes the following:

- The field party travelled to the study area to identify sites for sampling and measurements and to acquaint residents and project officials with the purpose of the study;
- Questionnaires were administered to residents in the villages within the study area by CEMMATS Social team to obtain data on several issues regarding water supply and quality and sanitation issues;
- Samples were collected from the identified water sources. Some physical parameters were tested in-situ and the remaining samples were transported to a laboratory for further physical, chemical and biological tests;
- The World Health Organisation's drinking water guidelines were used as a benchmark for drinking water contamination levels; and
- Critical observations of the nature of water systems and sanitary conditions in and around the project site.

Details of the analysis and results can be found in section 5.1.7 (Hydrology) of the report.

Biological Environment

Ecology

The Lugbu Chiefdom is located in the Bo District, Southern Province of Sierra Leone, and is biogeographically part of the western extent of the Guinea-Congo forest biome and the Upper Guinea Forest.

The landscape is now dominated by vast areas of agricultural fallow land (farm bush) as a consequence of the widespread traditional cut and burn agriculture. Much of the remnant forests are kept as sacred groves (society bush and cemeteries) and occur as gallery forest along river courses. No forest reserves exist within the chiefdom, but some of the gallery forest and sacred groves are still healthy and can be protected.

Social Environment

National Level

Sierra Leone covers a total area of 71,740 km² and had an estimated population of 4.9 million in 2004 (Sierra Leone Population and Housing Census, 2004) but according to the 2015 Housing and Population Census result, the population has increased to 7,075,641. Political instability and poor economic growth led to the brutal and destructive 10 year civil war which officially ended in 2002.

National Social Indicators

Key Social Indicators	Rate	Source
GDP per capita	\$497.89 in 2015	http://www.tradingeconomics.com/sierra-leone/gdp-per-capita

Key Social Indicators	Rate	Source
Economic growth rate	13.8% in 2014	World Bank, 2015
Infant mortality rate	87/1000	World Bank, 2015
Life expectancy at birth	46 years	HDI, 2015
Maternal Mortality ratio	1410/100,000	World Bank, 2014
Population Growth rate	2.19%	(World Bank, 2014)
Adult literacy	44.46	(World Bank, 2012)
Illiteracy rate	55.54%	(World Bank, 2012)
Primary school enrolment	1,353,723	(MEST), (WAEC, 2008/2010)
Net primary enrolment rate	78%	(MEST, 2014)
Gender parity in primary school enrolment for boys and girls	1.1 : 1	(MEST, 2014)

District Level

Administratively, Sierra Leone is divided into four distinct areas: the Northern Province with its headquarters in Makeni, the Southern Province with Bo as its headquarters, the Eastern Province with Kenema as its headquarters and the Western Area comprising the Freetown Peninsular with Freetown as its headquarters.

Bo District is the largest of the four administrative Districts in the Southern Region and occupies a total area of 7,003km² and a population of 574,201 people (Sierra Leone 2015 Population And Housing Census, 2016). Agriculture is the largest economic sector in the district with more than half of the population of Bo District depending on it as their main source of livelihood. Thus economic growth and poverty reduction in the district will only be sustained with development in this sector as the District is endowed with cultivable land (Upland and Inland Valley Swamps) and adequate rain-fall. The expansion of this sector in the district will stimulate growth in other sector like the supply of input, job creation, marketing, processing etc.

Chieftdom Level

Lugbu chieftdom is one of 15 chieftdoms in the Bo District. The general occupations of the inhabitants are agriculture, diamond mining and fishing. Agriculture, engaged in by more than two thirds of the inhabitants, is dominated by crop farming; this is mostly done on subsistence basis and employs rudimentary production techniques with the main crops being rice, cassava, groundnut, potatoes, yam, cocoyam and palm oil plantations. Fishing is also done in the Sewa River for both subsistence and commercial purposes.

A socio-economic baseline survey was carried out covering twenty-two (22) towns and villages in the Lugbu Chieftdom. Details of this assessment can be found in section 6.2 (Socio-economic Status and Living Conditions) of this report.

Identification of Potential Impacts

Land Preparation and Construction Stage

Impacts at this stage are often temporary. The main concerns at the construction stage will be environmental and occupational health and safety. Risks would be reduced by strict adherence to best construction management practices. In relation to land preparation for nursery development, the main concerns are loss of biodiversity and erosion. The following table presents environmental and social impacts anticipated during this stage, which remain of medium and above impact after mitigation. The full tables of impacts are presented in Table 7.2.6 (Land Preparation and Construction Phase - Environmental and Mitigation Measures) and Table 7.2.7 (Land Preparation and Construction Phase - Social Impacts and Mitigation Measures).

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Loss of habitat	Clearing of vegetation for nursery developme nt and constructio n	Clearing of vegetation in the concession area will result in loss of vegetation cover resulting in loss of habitat for various species. This may cause interruption of natural cycles including breeding and mating. Different species of flora may be lost which will not be reinstated when the area is reforested.	Certain	Moderate	Buffer zones will be established as well as botanical gardens/ ecological corridors. Although certain species lost during clearing may not be replaced, the cleared areas will eventually be re-vegetated with trees. A new thriving and balanced ecosystem will develop	Achievable	Medium
Loss of biodivers ity	Clearing of vegetation for nursery developme nt and constructio n	Clearing of vegetation, soil treatment and preparation before planting will cause local species within the project area to migrate to other areas, leaving the project site devoid of important species (mammals, reptiles, birds and insects) which contribute to the natural balance of the ecosystem within this area.	Certain	Moderate	Buffer zones will be established as well as botanical gardens/ ecological corridors. Some animals may return to the plantation areas, but natural processes may continue to be disrupted with operational activities including tree tending and treatment, harvest, etc.	Difficult	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Soil erosion	Site preparation for nursery development and construction	Exposure of loose soil resulting from vegetation clearing, soil preparation activities and pre-construction activities.	Certain	High	<p>Implementation of erosion prevention measures in particular contour farming and cover cropping to minimise excessive runoff and erosion. The installation of sediment traps in drainage lines and storm water collection channels/settling ponds.</p> <p>Vegetation will be left around rivers and surface water bodies to minimise sediment entry. Fields will also be surrounded by vegetation to create buffers.</p> <p>The risk of soil erosion will reduce once planting commences; the same applies once construction commences and the soil is no longer exposed.</p>	Achievable	Medium
Hydrology	Site preparation for nursery development and construction	Clearing of vegetation will result in an alteration of the natural hydrological cycle as evapotranspiration in the area will be reduced. Similarly without vegetation to break the volume of rainfall	Certain	High	This will be a short lived impact which will be reversed once the fruit trees begin to grow.	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
		reaching the soils, there will be an increase in surface runoff which has direct impacts on soil erosion and sedimentation in water bodies.			Ground cover crops and mulching will be used to reduce the impact of the rainfall on the soil.		
Water quality	Site preparation for nursery development and construction	Surface water runoff from the site may contribute to the total suspended solids and turbidity of the receiving water bodies which may affect the aquatic life. Runoff may also carry off construction related debris, spills or waste not properly handled.	Likely	High	Implementation of erosion prevention measures including ground cover (grass and other ground cover plants), the installation of sediment traps and storm water collection channels/settling ponds. Chemical spills in the field will not occur as all chemicals will be mixed at mixing stations which will be designed to contain spills during mixture and loading. Waste will be stored in secure waste receptacles where they will not be exposed to runoff.	Achievable	Medium
Noise pollution	Daily operations	Noise levels within the project area will increase particularly as a result of construction and other mechanised activities.	Certain	Moderate	Modern equipment will be used which are fitted with noise suppression devices to prescribed standards.	Difficult	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
					Noise generating agricultural activities such as land preparation (every 3 years), maintenance (fortnightly) and harvesting (every 12 – 14 months) will occur periodically and not on a daily basis.		
Loss of land	Project developme nt	Loss of land may be an issue even after land lease agreements have been reached.	Likely	High	<p>Intensive community consultations and additional sensitization meetings will be conducted prior to the commencement of this stage of the project to ensure that landholders' views and opinions are heard and taken into consideration. Locals will be made aware of the implications and potential impact of a large project starting in their communities, including the pitfall of giving away too much land.</p> <p>Land will be leased in phases as the project progresses in order to prevent leased land remaining unused over an extended period.</p>	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Community Health and Safety	Daily Activities	Community residents will be exposed to increased vehicular traffic and other operational activities which being unfamiliar puts them at risk of safety hazards.	Likely	High	Regular community consultations and community health and safety sensitization will be conducted periodically throughout the life of the project.	Achievable	Moderate

Operational Stage

Once best practices have been observed during the initial stages of the project - planning/design set up and construction stages - much of the threat to the safety and integrity of the environment and society will be reduced to levels defined by legislation and best practices.

The following table presents environmental and social impacts anticipated during this stage, which remain of medium and above impact after mitigation. The full tables of impacts is presented in Table 7.2.8 (Operational Phase - Environmental and Mitigation Measures) and Table 7.2.9 (Operational Phase - Social Impacts and Mitigation Measures).

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Loss of biodiversity	Clearing of vegetation for progressive expansion of plantation areas	Clearing of vegetation, soil treatment and preparation before planting will cause local species within the project area to migrate to other areas, leaving the project site devoid of important species (mammals, reptiles, birds and insects) which contribute to the natural balance of the ecosystem within this area.	Certain	Moderate	Buffer zones will be established as well as botanical gardens/ ecological corridors. Environmental monitoring plots also established to monitor changes in the area. Some animals may return to the plantation areas, but natural processes may continue to be disrupted with operational activities including tree tending and treatment, harvest, etc.	Difficult	Medium
Soil erosion	Clearing of vegetation for progressive expansion of plantation areas	Exposure of loose soil resulting from vegetation clearing and soil preparation activities	Certain	High	Implementation of erosion prevention measures including in particular contour farming and cover cropping to minimise	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
					<p>excessive runoff and erosion, the installation of sediment traps and storm water collection channels/settling ponds.</p> <p>Vegetation will be left around rivers and surface water bodies to minimise sediment entry.</p> <p>The risk of soil erosion will reduce once planting commences; the same obtains once construction commences and the soil is no longer exposed.</p>		
Hydrology	Irrigation	Water abstraction from the Sewa River could affect the availability and hydrological regime within the project area.	Certain	Moderate	Agricultural practices such as no till or the use of cover crops/mulch and the incorporation of organic matter from plant residues are all	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
					<p>Good Agricultural practices which will mitigate excessive irrigation requirements.</p> <p>Pineapple is a drought tolerant plant so can withstand some water stress.</p> <p>Alternative water sources will be made available for the project and communities through the sinking of boreholes. These will supplement and reduce the water obtained from the Sewa River.</p>		
Water quality	Use of Fertilisers and Pesticides	Water pollution can occur due to the use of agro-chemicals such as fertilizers and pest control substances which can leach into the soil	Certain	High	Precision Agriculture methods will be employed with regard to crop nutrition, and fertilizer will only be applied in the required	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
		and get carried into water by runoff. Aquatic life would be affected.			amounts foliarly so contamination of surface water is greatly reduced. Chemicals i.e. Herbicides and pesticides will be used only as required or when particular pest thresholds are reached.		
Noise pollution	Daily operations	Elevated noise levels will result from movement and operation of vehicles and machinery.	Certain	Moderate	Modern equipment will be used which are fitted with noise suppression devices to prescribed standards. Noise generating agricultural activities such as land preparation (every 3 years), maintenance	Difficult	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
					(fortnightly) and harvesting (every 12 – 14 months) will occur periodically and not on a daily basis. Workers will be provided with noise protection PPE.		
Emergency Response and Disaster Management	Throughout project life	Poor management of emergencies or disasters will result in loss of life, damage to equipment and disruption of operations	Likely	Very high	Implementation of Emergency Response Plan, regular drills and training exercises for staff, monitoring of potential situations leading to disaster.	Achievable	Medium
Community Health and Safety	Daily Activities	Community residents will be exposed to increased vehicular traffic and other operational activities which being unfamiliar puts them at risk of safety hazards.	Likely	High	Regular community consultations and community health and safety sensitization will be conducted periodically throughout the life of the project.	Achievable	Moderate

Environmental and Social Management Plans

The Environmental and Social Management Plan (ESMP) for the Project identifies the principles, approach, procedures and methods that will be used to control and minimize the environmental, social, health and safety impacts of all project activities.

Environmental Health and Safety Plan

The Environmental Health and Safety Plan (EHS) Plan identifies the principles, approach, procedures and methods that will be used to control and minimize the adverse environmental and social impacts of all construction and operational activities associated with project development. It is intended to complement the project's Environmental and Social Impact Assessment (ESIA) and ensure that commitments made by Sierra Tropical Ltd. to minimize project related adverse environmental and social impacts are upheld throughout all project phases.

Waste Management Plan

The Waste Management Plan (WMP) describes the procedures, systems, equipment, and structures specific to waste management and disposal. Waste generation should be limited at all levels of the operation in order to decrease the volume of waste generated and make waste disposal more manageable. The WMP also defines who is responsible for developing and implementing the plan, and what records and reporting will be required.

Pesticide and Chemical Management Plan

The Pesticide and Chemical Management Plan (PCMP) outlines the general measures and approaches for prevention and mitigation of occurrences of pesticides and other agro-chemicals in ground and surface waters following their application, and recommended actions to accomplish appropriate goals. Risks associated with the use of pesticides during the period of managing a pest outbreak are outlined.

Emergency Response Plan

The Emergency Response Plan provides employees and managers with specific instructions that will enable them to respond quickly and efficiently to any foreseeable emergencies likely to occur at the Project. It is developed using recognized and accepted methods and practices and includes specific responses, protocols, and management contacts. The ERP essentially has the goal of protecting people, the environment, property and the operations. This document deals with typical emergency types that characterize the operation which include:

- a. Fire or Explosion;
- b. Pollution or Chemical Spills;
- c. Flooding, Landslides, Rain Storm (Natural Conditions);
- d. Medical Health Cases;
- e. Civil Unrest & Disturbances.

Resettlement Policy Framework

It is not envisaged that the project will entail any displacement of people and extensive damage to crops. However, it is possible that the infrastructure requirements of the project in terms of feeder road construction/rehabilitation and other project logistical requirements may result in damage to crops. It is impossible at this stage to determine and assess these, procedures should nevertheless be put in place to handle such matters if they arise.

Community Development Action Plan

Specific community development and social assistance programmes aimed at improving the living conditions of the local communities in a sustainable way are captured under the CDAP.

Public Consultation and Disclosure Plan

The PCDP is intended to define objectives and establish the framework necessary to provide understandable information to all parties involved. This plan will be implemented to ensure timely and effective communications with the project management and the affected stakeholders, throughout the lifetime of the project. The main objective of the PCDP is to establish a program for multi-directional communication between the management and stakeholders.

Closure Plan

After the project has reached its economic life span and can no longer be operated in an efficient, reliable and safe manner, the Project reclamation activities are implemented to re-establish a beneficial post-operation land use. All structures would have to be removed in order to return the site to its pre-construction phase. Monitoring programmes will be implemented to ensure that the site reverts to a natural and useable state.

Management, Mitigation, Monitoring and Implementation Measures

The Environmental Monitoring Plan (EMP) outlines the plans for monitoring within the plantation areas and project facilities; key personnel and their responsibilities are also identified.

Conclusion

Key Assessment Findings

Physical Environment

There are potential major impacts relating to hydrology and water quality, biodiversity, soil erosion, water & soil pollution due to agro-chemical and water abstraction. Mitigation measures to limit the extent of impacts have been highlighted and will be implemented.

Biological Environment

There will be some loss of floral and faunal species during land clearing and preparation, however it is not expected that any species will be wiped out from the region as a result. Mitigation measures have been presented to ensure that minimal clearing is carried out to limit the extent of biodiversity loss.

Socio-economic Environment

Perhaps the most critical aspect of the project is the potential reduction of farm land by the communities, potential conflict from issues related to labour, unrealistic expectations held by the communities with regard to benefits created by the project, vehicular traffic and safety risk. The selection of the project site has been judiciously done to avoid the need for community relocation.

The project will have positive impacts in the area of job creation, improving the quality of life of locals through payment of land lease rent and compensation, outgrower schemes, creation of secondary businesses, etc.

Conclusion

This Environmental and Social Impact Assessment has been professionally carried out to satisfy the Government of Sierra Leone's legislation and regulations. The study has achieved the following objectives for such an exercise, viz:

- Baseline Survey of the Project Area;
- Execution of an Environmental and Social Impact Assessment (ESIA) and development of Environmental and Social Management Plans (ESMP);

The investigations of impacts on the social environment are a crucial part of the study, since the operation may impact the communities which reside at close proximity to the project site.

The investigation of social impacts has involved the following:

- A baseline socio-economic study of the community envisaged to be impacted by the project activities in both the construction and operational phases;
- A Public Consultation and Disclosure Process (PCDP) undertaken to sensitize stakeholders.

The baseline environmental and socio-economic situations have been presented.

Environmental impacts of the project's activities have been identified for all phases of the project. The most significant of these impacts is the change of morphology and soil and water (surface and ground) pollution due to agro-chemicals utilization. Social impacts include loss of agricultural farm land and unrealistic expectation from the people regarding benefits as the project progresses.

Generally, the investigations reveal that environmental and social problems incurred by the project can be adequately managed and that there are no insurmountable problems that should stop the project from proceeding.

A monitoring system must however be put in place to ensure that management practices for mitigating negative impacts and enhancing those that are positive are affected. It must however be ensured that recommendations made in the Environmental and Social Management Plans are followed through.

1 INTRODUCTION

1.1 Project Background

Sierra Tropical Ltd. is a wholly owned subsidiary of Dole Asia Holdings Pte. Ltd., a Singapore based Agro-processing Company. A leader in growing fruits and vegetables, Dole Asia Holdings also markets or processes these fruits into fruit juice and healthy snack products throughout the world.

Sierra Tropical Ltd (STL) proposes to embark on an agro-processing project involving large scale planting and processing of tropical fruits in Sierra Leone using approximately 15,000 hectares. It is envisioned to be fully integrated from cultivating, growing and harvesting pineapple, mango, papaya and other tropical fruit and processing them into manufactured products. The project is proposed to be implemented in Sierra Leone as it has the necessary favorable conditions, such as ideal weather, plentiful land, duty concessions, and a business friendly government and administration particularly with regards agriculture.

The project will initially grow up to 4,335 hectares of leased land in Lugbu Chiefdom, Bo District, and will be implemented in phases, starting with the development of a pineapple nursery, spreading out in phases to eventually cover the maximum area desired and possible.

If the first phase of the project is successful and the quality/productivity of the fruits meet our requirements, a manufacturing facility will be constructed in the later stages of the second phase, to process the raw materials into manufactured products in various packages of cans, drums, plastics and boxes. An extension of this ESIA Study will be done at this point, to cover the manufacturing facility; however this will be dependent on the success of phase 1 and the early stages of phase 2.

1.2 National Perspective

The agricultural sector, comprising food crops, tree crops, fishery, livestock, and forestry sub-sectors is the backbone of Sierra Leone's economy (Ighobor, 2014) with the crop sector dominating the agricultural GDP with 33% in 2010 (Larbi, 2012). More than half of the country depends on agriculture for livelihood.

The major food crops are rice, cassava, maize, millet, sorghum, sweet potato, and groundnut which are produced by smallholders with an average land holding averaging from 0.5-2.0 hectares (Larbi, 2012).

Rice is the most important crop, cultivated by nearly 85% of farmers, while cassava is the second most important crop. Oil palm is the main tree crop and other perennial crops include citrus, cocoa, coffee, coconut and sugarcane (Larbi, 2012).

Crop yields are generally low due to: limited access to agricultural inputs, low levels of mechanization, pest and disease problems, lack of markets and market information, labour shortages and a weak private sector.

There has been a steady decrease in the amount of funds spent by the government on food importation due to its drive to promote increased agricultural capability within the country. Between 2007 and 2013, costs on food importations dropped from \$32 million to \$15 million, improvement which has been confirmed by the World Health Organisation (Ighobor, 2014). The government believes that the agriculture sector can be further improved, and aims to increase the national budget to more than 10% of the national budget. This increase will allow farmers to have access to farm implements, machinery, fertilisers and other materials that will enhance their work/yield.

1.3 Project Area

The project will start in Lugbu Chiefdom, Bo District which will primarily be the host community. It is however anticipated that as the project develops, more agricultural land will be required which will necessitate expansion to other areas within Lugbu Chiefdom and possibly into the neighbouring Tikonkor Chiefdom.

Since five of the six Sections in Lugbu Chiefdom qualify for the establishment of pineapple farms in terms of land suitability and environmental friendliness, sufficient land is available in the Chiefdom for the next phases of the project (three to ten years and beyond). There would also be adequate land within the settlements for food production.




Table 1.3-1: Lugbu Chiefdom

1.3.1 Brief History of Lugbu Chiefdom

According to Reed and Robinson's history of paramount chieftaincies in Sierra Leone (2012), it is believed that the father of the first Paramount Chief of Lugbu Chiefdom Samuka Nallo was a warrior and hunter who came from Sudan. In an account that has not been verified, he is said to have travelled through Liberia and entered Sierra Leone through Pujehun, settling in a place called Gikubu, a Mende ward meaning settlement under the tree.

This chiefdom is mostly descended from two related families – Nallo and Magawo. The Nallo family has however, has dominated the chieftaincy since the early 20th century.

1.4 Environmental and Social Impact Assessment Process

1.4.1 Stages of the ESIA Process

Prior to commencement of any project that may affect the environment and communities, it is mandated by legislation that an Environmental and Social Impact Assessment (ESIA) study be done, and, upon approval by EPA-SL, a licence is secured.

The Sierra Leone Environment Protection Agency Act, (SLEPAA) 2008 and the EIA Supplementary Acts, 2010 describe the requirements and process for securing an EIA licence, which is laid out in a "checklist" prepared by EPA-SL. In short, the client first applies to the local regulatory body, Environment Protection Agency, Sierra Leone (EPA-SL) for an EIA

licence. EPA-SL requires that a screening form be filled and submitted with the application letter, after which a decision is made on the category of the project; this is followed by a scoping report. EPA-SL will then decide on the terms of reference (TOR) to be drafted by the project proponent or an independent consultant hired by the proponent.

On the approval of the agency, the consultant carries out an assessment of the environmental and social impacts of their planned operations on ecosystems and communities in the project area. A report is prepared at the end of the study and submitted to EPA-SL for review. If approved, the proponent will then be requested to conduct public disclosure meetings with relevant stakeholders on the findings and recommendations of the study, and incorporate comments, suggestions and requests made during those meetings into a public consultation and disclosure report. Finally, all reports pertaining to the ESIA study are then forwarded to the Board of EPA-SL for a decision to be made on the issue of the licence.

1.4.2 Purpose of the ESIA Study

The purpose of the environmental impact assessment is to identify and mitigate potential negative and identify and enhance positive environmental impacts. This is done through the conduct of desktop and field studies to:

- Obtain secondary and primary biophysical and socio-economic data;
- Anticipate the potential impacts of the proposed project on the environment and communities;
- Propose an environmental management plan that mitigates adverse impacts whilst enhancing positive ones;

These are achieved by employing a methodology that *inter alia* consists of a literature review, field investigations and a social survey.

The purpose and findings of the study will be disclosed to project interested and affected persons in a series of stakeholder consultation and disclosure meetings to elicit community acceptance and participation that are imperative for the construction and operation of the facility.

1.4.3 Objectives of the ESIA Study

The objectives of the study were as follows:

- i. To assess the potential positive and negative impacts of the planned construction project and its operations on society and the environment;
- ii. To recommend mitigation measures to avoid or mitigate negative impacts and enhance benefits;
- iii. To prepare an environmental and social management and monitoring plan (ESMP) that integrates mitigation and monitoring measures;
- iv. To develop specific topic management plans which will include the following:

- a. Waste Management Plan (WMP)
- b. Environmental Health and Safety Plan (EHSP)
- c. Community Development Action Plan (CDAP)
- d. Public Consultation and Disclosure Plan (PCDP)
- e. Emergency Response Plan (ERP)
- v. To conduct public disclosure and consultation meetings on the findings and recommendations of the EIA study.

1.4.4 ESIA Consultants and Teams

A number of specialists were identified to undertake the investigations and address the issues outlined during the ESIA phase. A team was formed of the respective ESIA consultants and sub-teams (Table 1.4-1).

The terms of reference for each of these studies are outlined in the next section. The specialist studies were undertaken during the ESIA phase. This Environmental and Social Impact Assessment (ESIA) report summarizes all their findings and has been compiled using the information gathered during these studies. The recommendations and mitigation measures developed from these studies have also been pulled together to generate an Environmental and Social Management Plan (ESMP) (Volume 2 of this report) which will be adhered to during the various stages of the project.

Table 1.4-1 ESIA Consultants and Teams

Specialist	Specialist investigation
Andrew Keili	Project Director
Vanessa James	Project Manager/Infrastructure and Operations Assessment
Ralph Bona/ Josephine Turay	Hydrologist/water quality specialist
Leonard Buckle	Soils/air/noise specialist
Arnold Okoni Williams	Ecologist- Flora and Fauna
B. Bockarie	GIS/Land Use expert
J. Lappia	Socio-Economist/Resettlement expert/lease expert
/Rashidu Sinnah	Socio-Economist

Specialist	Specialist investigation
Joseph Gbassah	GIS/ Environmentalist

1.5 Description of the Terms of Reference (TOR)

The ESIA study conducted by CEMMATS Group Ltd, was primarily done to meet the local requirements for securing an Environmental Impact Assessment licence from EPA-SL.

This study commenced in August 2016 and consists of biophysical and socio-economic baseline data collection and impact assessments conducted in the project area by an inter- and multi-disciplinary team of professionals. The Terms of Reference as submitted to and duly approved by the EPA-SL is outlined in the following table.

Table 1.5-1 Terms of Reference of the ESIA Study

Field	Objectives	Methodology
Hydrology	<ul style="list-style-type: none"> Determine direct physical impacts of the project on the surface and groundwater. Recording baseline qualitative and quantitative data. Suggest mitigation measures to address the impacts noted in the surface and groundwater hydrologic assessment. 	<p><u>Desk study:</u> A review of existing data on this area will be undertaken together with additional information which may include:</p> <ul style="list-style-type: none"> A description of the site in terms of catchments characteristics such as site boundaries, Mean Annual Runoff (MAR), Mean Annual Evaporation (MAE), Mean Annual Precipitation (MAP) using available sources (National Reports); A review of land (water) uses and predicting the potential impacts that could arise from the proposed project; and Review of existing baseline water quality data (local/regional) and the national water quality targets applicable to the project. <p><u>Field surveys:</u></p> <ul style="list-style-type: none"> Verify catchment characteristics such as sub-catchments, vegetation cover, land uses, topography, and average slopes in order to describe the hydrological characteristics of the project site ; Collect additional samples to submit these to a laboratory for analysis Identify potential sources of pollution. Identify potential impacts that could result from the proposed project on the surface water resources.
Hydrogeology	<p>This phase will include a desktop study to acquire relevant hydrogeological information for studies performed in the area to date. A hydrocensus will be taken of the study area. During the hydrocensus important data pertaining to the current groundwater conditions and use will be collected. This will include:</p>	<p>The hydrocensus surveys will include interviews with landholders/managers, visits to individual wells. This data together with its spatial distribution will determine the current water resource and environmental status and serve as reference to any future environmental projects conducted in the area.</p>

Field	Objectives	Methodology
	<ul style="list-style-type: none"> localities of current groundwater abstraction points ownership current usage volumes and types, Equipment and groundwater levels. <p>Hydro chemical samples will also be taken from selected wells. The data collected will serve as a reference point against historical and future groundwater conditions in the area.</p>	
Water Quality	<ul style="list-style-type: none"> Provide basic characterization of the quality of surface and potable water sources. Assess the risk of contaminating ground water, springs and surface water. 	<ul style="list-style-type: none"> On-site Measurements of physico-chemical parameters such as, pH, Temperature, dissolved oxygen, turbidity and conductivity. Sampling for laboratory analysis of chemical (anions and cations) and microbiological (faecal and total coliforms) characteristics. Information from the original ESIA study of this area will also be utilised.
Soils Assessment/Agricultural issues	<ul style="list-style-type: none"> To characterize the soils, in the study area that are to be affected by the proposed project. It will provide an indication of the existing soil and land capabilities for the survey area and give a characterization of the land capability within the study area. To describe and assess the agricultural issues related to the project 	<p>Some areas within the lease will be surveyed by hand auger observations. Coordinates of the observation points will be loaded onto a GPS to locate points in the field.</p> <p>Auger observations will be made to the depth of the first restricting layer. The following attributes will be recorded at each observation point of the detailed survey:</p> <ul style="list-style-type: none"> Soil form classified according to an appropriate Soil Classification System; Soil depth; Estimated soil texture; Soil structure; Current land use; Land capability. Samples will be analysed for soil acidity, fertility and textural indicators

Field	Objectives	Methodology
		<p>as follows:</p> <ul style="list-style-type: none"> • pH (water); • Extractable cations and Na, K, Ca, Mg • Phosphorus • Cation exchange capacity (CEC); • Carbon content; • Soil texture (3 classes namely sand, silt and clay). • Describe and assess salient agricultural issues
Air Quality	<ul style="list-style-type: none"> • To identify key aspects that might have significant air quality impacts during all phases of the project. • Ambient and meteorological data will be sourced for the area under investigation. • Qualitative assessment will be undertaken which will evaluate the possible impacts of other polluting sources in the area. • Point and non-point sources of pollution 	<p>Identification and quantification of dust emissions sources; Development of emissions inventory</p>
Noise Assessment	<p>To assess the ambient noise levels in settlements in and around the project area in accordance with international guidelines.</p>	<ul style="list-style-type: none"> • Preliminary survey and identification of measuring points for day and night time readings. Monitoring measurements will be taken at 1.5m above ground level; • Sound pressure readings will be done at the closest residential area; • Noise survey at the identified measuring sites – Ambient noise measurements;
Land Use	<ul style="list-style-type: none"> • Engage with the social scientists to ensure that questions related to land use are asked during the social impact assessment; • Clarify the complexities associated with 	<ul style="list-style-type: none"> • Set up focus groups together with the social scientists to answer these questions; • A landuse GIS cover dataset will be digitised using visual identification classification method. The most recent colour aerial photography will

Field	Objectives	Methodology
	<p>rotational land use;</p> <ul style="list-style-type: none"> • Where possible, integrate findings with those obtained from soil surveys; • Develop strategies, to deal with issues related to income generation and food security; • Ensure that the study deals with any other issues related to land use raised during scoping; • Assess the environmental significance of the anticipated impacts; • Provide feasible recommendations on the mitigation of impacts related to land use. 	<p>augment our base maps for this analysis;</p>
Ecology-Fauna	<p>Amphibian, reptile and mammal surveys to include causal observations and refuge examinations (e.g. searching under rocks, logs, rotten tree stumps, leaf litter, old termite mounds and rodent burrows with consideration to Health and Safety protocols). Information will also be available through sightings from local people; however, these data will only complement the main survey. Favourable habitats to also be examined within the development area. Priority will be given to key indicator groups such as:</p> <ul style="list-style-type: none"> • frogs and toads • small reptiles such as lizards • other small mammals • monkeys <p>Methods for surveying the above groups will simply involve visual observations and photographs.</p>	<p>Results for the ecological survey for this area in the original ESIA study will be utilised as best as possible. Additional information in areas within the lease area will be collected. The collective information will cover the following issues:</p> <ul style="list-style-type: none"> • Consideration will be given to the location and whereabouts of each survey undertaken before arriving at site. • Surveyors will record the habitat types present in each of the survey areas • The location of all surveys undertaken will be recorded with the aid of a handheld GPS. • Photographs will be taken of the animals, where possible. • Sightings from people and local communities within the vicinity should be noted in the reports but remain separate from the field surveys above. Importance and relevance of species identified in respect of local stakeholders should also be included in the reports.

Field	Objectives	Methodology
Ecology-Flora/wetlands	<p>Sensitive and critical plant species (species of conservation importance) will be identified and habitats mapped.</p> <p>The position and locality, as well as species composition of sensitive areas inventory will be conducted in order to identify and map all wetlands in the area.</p>	<ul style="list-style-type: none"> • A floristic survey will be conducted to determine the species composition of the area of interest. This will give an indication of the actual species present on site and these will be discussed in context of plant communities (should the area support distinct communities) within the ecosystem of the area. • Assessment of rare species of flora and fauna will be carried out; this will identify the categories of biodiversity and give an indication of the offset required. Possible locations for biodiversity offset, if required will be suggested as the National reserves surrounding the project area are identified. <p><i>Wetlands</i> A baseline investigation involving a desktop assessment and limited ground truthing will be undertaken covering the proposed site. Wetlands, including riparian wetlands within the site boundaries will be identified using a range of tools, including topographical maps.</p>
GIS and mapping	<p>The most progressive means of mapping is through the use of Geographic Information Systems (GIS) as it enables sound cartography as well as a holistic analysis of an area. GIS is a digital cartographic tool capable of creating, integrating, storing, editing, analysing, sharing, and displaying geographically referenced information. It allows users to create queries, analyse spatial information as well as editing and managing data with a final product being the creation of maps to aid in visual interpretation.</p> <p>The use of mapping in EIAs is integral, as maps provide the means through which the proposed</p>	High resolution aerial surveys, satellite imagery will also be used

Field	Objectives	Methodology
	development can be viewed within a spatial context.	
Technical Risk assessment	Describe the process and assess the technical risks of the process including occupational health and safety issues	<p>Description and assessment of the technical processes. Technical risk assessment includes issues such as:</p> <ul style="list-style-type: none"> • Electrical installation • Fire safety consideration • Waste management system • Operations management (fuel transport, loading/offloading, storage, staff training) • Scheduled equipment maintenance and record keeping on site
Infrastructure Assessment		<ul style="list-style-type: none"> • Identify and describe various community infrastructure such as hospitals, health centres, <i>court barriers</i>, road networks etc; • Identify and describe the possible impacts related to the construction of access and maintenance roads; • Ensure that the study deals with the issues raised during scoping; • Assess the environmental significance of these impacts; • Assess the environmental impacts of other associated infrastructure.
Visual Assessment	<ul style="list-style-type: none"> • To inspect the project area and include a photographic survey with a focus on the project site. • Undertake a desk-top study analysis of maps, plans etc; • Ensure that the study deals with issues related to land use raised during scoping 	<ul style="list-style-type: none"> • Features of the project installation that are of particular relevance to the visual impact assessment will be reviewed; • GIS modelling tools will be employed to assess depth, elevation etc. • Review and assessment of photographs of the project area

Field	Objectives	Methodology
	<ul style="list-style-type: none">Assess the environmental significance of these impacts	

1.6 Scope of Work

The purpose of the environmental impact assessment is to obtain prevailing biophysical and socio-economic data, anticipate the potential impact of the proposed project on the environment and communities and propose an environmental management plan to mitigate adverse impacts whilst enhancing positive ones. These will be achieved by employing a methodology that consists of literature review, field visits and administration of questionnaires.

The ESIA study covers the start-up area of the project of 4,335 ha and related activities including nursery development, establishment of plantations, construction of administrative buildings and quarters.

1.7 Assumptions and Limitations of Study

1. The EIA study is done mainly to meet the local requirements for securing the EIA licence; relevant international guidelines will be taken into consideration and referred to during the study.
2. The initial ESIA will be done to assess the agricultural phase of the project. Additional studies for the processing phase of the project will be carried out closer to the implementation of this phase and included as an addendum to this ESIA document.
3. Consultation will be carried out and relevant information disseminated throughout the execution of the project which will keep to the tenets of a proper Public Consultation and Disclosure Process (PCDP) for an EIA. However the EPA-SL would normally require that the results of the EIA report are discussed with communities after submission of the report.
4. The timeframe within which the study is carried out will not allow for seasonal variations to be taken into consideration, however desk studies was done on historical climatic data and other records obtained at various times during previous years.

1.8 ESIA Study Boundaries

In carrying out an ESIA project like this one, various issues outside the immediate project boundary may have to be considered. The project spatial boundaries which have been considered for this project include:

- **Project Boundary** - Includes all areas of direct physical changes - the 'footprint' of the Project.

- **Environmental (Ecological) Boundary** - Encompasses all areas in which environmental impacts, including indirect impacts, may occur.
- **Socio-economic Boundary** - Includes all communities potentially affected by the Project. The Project area and the area of potential environmental, physical and biological impacts fall within the Socio-economic Boundary.
- **Administrative Boundary** - Includes the various jurisdictions with influence on the Project.
- **ESIA Study Boundary** - Includes source of the impacts as well as the area of potential environmental and socio-economic concern. Study was carried out during the rainy season.

1.9 Organisation of the ESIA Report(s)

The final report consists of two volumes. Below are brief comments on the contents.

1.9.1 The ESIA Report

Volume 1 – Environmental and Social Impact Assessment (ESIA) contains the Executive summary and main report. The Executive Summary presents a concise overview of the significant findings recommendations and actions contained in the ESIA. The Main Report contains the policy, legal and administrative framework under which the ESIA was carried out. There is an analysis of the feasible alternatives, including the “no project” alternative, and a description of the project in its geographic, ecological, social and temporal context. It includes baseline data describing the relevant physical, biological and historical conditions and the environmental effects associated with project implementation. Mitigation measures needed to control those effects to acceptable levels are presented.

Volume 2 – Environmental and Social Management Plan (ESMP) presents the environmental management, mitigation, monitoring and institutional measures to be undertaken during project implementation and operation to reduce adverse environmental and social effects to acceptable levels and to enhance potential benefits. It specifically defines what actions must be taken and who is responsible to reduce adverse project impacts and enhance benefits. The ESMP also includes action plans for waste management, emergency response, closure, community development, and public consultation and disclosure. The ESMP highlight the issues and concerns that are presented in the ESIA report and identifies reasonable and practical responses to address and mitigate potentially adverse effects/enhance benefits. It describes the specific actions that will be required to effectively implement those responses in a timely manner and describes the methods by which management will demonstrate that those requirements have been met. It also establishes the course that will be followed in complying with Government of Sierra Leone environmental laws and regulations as well as international policies and guidelines.

1.9.2 Management Plans featured in the ESMP

1.9.2.1 Environmental Health and Safety Plan

The Environmental Health and Safety Plan (EHS) Plan identifies the principles, approach, procedures and methods that will be used to control and minimize the adverse environmental and social impacts of all construction and operational activities associated with project development. It is intended to complement the project's Environmental and Social Impact Assessment (ESIA) and ensure that commitments made by Sierra Tropical Ltd. to minimize project related adverse environmental and social impacts are upheld throughout all project phases.

1.9.2.2 Waste Management Plan

The Waste Management Plan (WMP) describes the procedures, systems, equipment, and structures specific to waste management and disposal. Waste generation should be limited at all levels of the operation in order to decrease the volume of waste generated and make waste disposal more manageable. The WMP also defines who is responsible for developing and implementing the plan, and what records and reporting will be required.

1.9.2.3 Pesticide and Chemical Management Plan

The Pesticide and Chemical Management Plan (PCMP) outlines the general measures and approaches for prevention and mitigation of occurrences of pesticides and other agro-chemicals in ground and surface waters following their application, and recommended actions to accomplish appropriate goals. Risks associated with the use of pesticides during the period of managing a pest outbreak are outlined.

1.9.2.4 Emergency Response Plan

The Emergency Response Plan provides employees and managers with specific instructions that will enable them to respond quickly and efficiently to any foreseeable emergencies likely to occur at the Project. It is developed using recognized and accepted methods and practices and includes specific responses, protocols, and management contacts. The ERP essentially has the goal of protecting people, the environment, property and the operations. This document deals with typical emergency types that characterize the operation which include:

- a. Fire or Explosion;
- b. Pollution or Chemical Spills;
- c. Flooding, Landslides, Rain Storm (Natural Conditions);
- d. Medical Health Cases;
- e. Civil Unrest & Disturbances.

1.9.2.5 Resettlement Policy Framework

It is not envisaged that the project will entail any displacement of people and extensive damage to crops. However, it is possible that the infrastructure requirements of the project in terms of feeder road construction/rehabilitation and other project logistical requirements may

result in damage to crops. It is impossible at this stage to determine and assess these, procedures should nevertheless be put in place to handle such matters if they arise.

1.9.2.6 Community Development Action Plan

Specific community development and social assistance programmes aimed at improving the living conditions of the local communities in a sustainable way are captured under the CDAP.

1.9.2.7 Public Consultation and Disclosure Plan

The PCDP is intended to define objectives and establish the framework necessary to provide understandable information to all parties involved. This plan will be implemented to ensure timely and effective communications with the project management and the affected stakeholders, throughout the lifetime of the project. The main objective of the PCDP is to establish a program for multi-directional communication between the management and stakeholders.

1.9.2.8 Closure Plan

After the project has reached its economic life span and can no longer be operated in an efficient, reliable and safe manner, the Project reclamation activities are implemented to re-establish a beneficial post-operation land use. All structures would have to be removed in order to return the site to its pre-construction phase. Monitoring programmes will be implemented to ensure that the site reverts to a natural and useable state.

1.9.2.9 Management, Mitigation, Monitoring and Implementation Measures

The Environmental Monitoring Plan (EMP) outlines the plans for monitoring within the plantation areas and project facilities; key personnel and their responsibilities are also identified.

2 PROJECT DESCRIPTION

Through Sierra Tropical Ltd., Dole Asia Holdings will establish a new agro-processing project in Sierra Leone. It has elected to set up its project in Sierra Leone because there are the necessary favourable conditions, such as ideal weather, plentiful land, duty concessions, business friendly government and administration especially to agriculture.

The project is envisioned to be fully integrated from cultivating and marketing fresh pineapple papaya and tropical fruits as well as processing those fruits into manufactured products in various packages of cans, drums, plastics and boxes, thereby adding further value to the product. The manufacturing aspect of the project will involve the construction and installation of a cannery and a canning plant. The finished products will be exported to Europe, United States, Middle East and Africa, as product of Sierra Leone, primarily under the Dole label.

As part of its corporate responsibility, the project intends to develop a smallholder or out grower scheme in the areas where the project sites are located. Activities to be developed may include but not be limited to providing access to basic agricultural inputs, training, market access, etc. in line with the Smallholder Commercialization Programme (SCP) and the Food Security flagship programme of GoSL. It is expected that this program will create more local employment directly and indirectly and positively impact on the livelihood of the surrounding communities.

In addition, Sierra Tropical Limited will establish and implement appropriate training programs to all levels of its employees. Training programs will be developed for potential Sierra Leoneans who will be tapped to occupy key positions in the supervisory and managerial positions.

The acquisition of the entire 15,000 hectares required for the project will be done gradually, over a series of phases. Phase 1 will last approximately 2 years, and will utilise a minimum of 800 hectares. The second phase will be implemented in the third year and will last for 3 years, with a proposed expansion to a minimum of 4,000 hectares. For years 6 to 10, more than 2,000 hectares roughly may be acquired, and with the proven success of Phase 2, expansion will continue to increase steadily. Using this projection, and assuming success at each stage, the project is estimated to continue for over well over 50 years.




Figure 1.9.11.0-1: Proposed Location of STL Project Features Spanning 1st, 2nd and 3rd Phases of the Project

Details of each phase are as follows:

Phase 1—Technical Feasibility Study

Phase 1 will be a technical feasibility study phase in order to confirm the viability of establishing a pineapple plantation in the areas identified for the project site. While the Company estimates completion of this phase approximately two years from commencement, the company may need to extend this period in order to fully accomplish its feasibility study.

In this phase, the company intends to lease land, import planting material and other agricultural inputs, establish a nursery, and implement land preparation, set up generators and water supply, and other capital requirements necessary to establish a pineapple plantation. Sierra Leoneans, who will be adequately qualified and trained, will be employed to implement the project.

The main focus of Phase 1 in addition to initial commercial planting of pineapple is to propagate planting materials through development of a nursery. Sufficient planting materials need to be generated to achieve commercial planting of Phase 1 covering approximately 1,200 hectares and, if the feasibility study is confirmed, to expand to the end of Phase 2 covering up to or beyond 4,000 hectares and onwards. Planting materials will be initially imported as there are no materials available locally of the quality and variety that the company requires.

Sierra Tropical Ltd. will invest more than or up to a total of US\$4.5 million for Phase 1. This investment will cover cost of the start up processes, legal, consultations, importing planting materials, cost of preparing land for commercial and nursery plantings, establishing and maintenance of a nursery, purchasing of various agricultural equipment, construction of buildings, purchase of generators, construction and installation of water supply and water treatment, and construction of offices, first aid facilities and staff housing.

This phase would require about 500 or more direct, causal or indirect employees estimated to be broken down into 490 or more Sierra Leoneans and possibly up to 8 to 10 expatriates, also on a rotational basis will come from Dolefil, Agricultural and other Specialist to develop the farm the practices and to train the local supervisors and employees. From 2 to 4 persons every 3 to 4 months. However, the project will indirectly use hundreds more Sierra Leoneans as it contracts out several construction projects covered by Phase 1. As the project develops, Sierra Tropical Ltd. will begin looking for a new generation of supervisors and managers to be trained. The Dolefil training teams will assist in this as stated above. They will be appropriately trained to assume future positions in the project if the company confirms the viability of the project after Phase 1 and then expands its operations.

A complete growing cycle of pineapple is about 36 months from time of planting to harvest of first crop and second crop and allowing for up to 6 months fallow period to rest the land in preparation for the next cycle. Phase 1 commercial fruit planting produce less than 1,000 metric tons of fresh pineapple for testing of fruit quality.

Sierra Tropical Ltd. will spend more than US \$3 million in expense costs for the two-year coverage of Phase 1. This is in addition to the US\$4.5 million it will spend for capital investment. In all, Phase 1 spending by Sierra Leone Tropical Limited will amount to more than US\$7.5 million. Any obligations of the Company beyond the Phase 1 feasibility study is subject to the Company's decision whether to proceed with the project following such study. The Company has the right to terminate the project without any obligations at any time during or following the Phase 1 feasibility study if the Company determines it is not feasible to continue with the project.

Phase 2 -First Expansion

If Sierra Tropical Ltd. determines to continue with the project following Phase 1, it then intends to further expand its operations. In this phase, the company intends to lease more land and increase plantation capability. Furthermore, the Company intends to establish a cannery and also install additional equipment for IQF Frozen products and other processes for the expansion of this project.

The Company intends to produce Pineapple Juice, Concentrate and Canned Pineapple and expand to IQF Pineapple and other processed fruits for export to various countries of the world. Phase 2 will start the development of the smallholder or outgrower scheme for the production of different tropical fruits. This will be done in collaboration with the Paramount Chiefs of the Chiefdom Councils where the project sites are located. The extent of the smallholder or out grower scheme for pineapple will depend largely on the availability of planting materials and equipment and the capacity of the outgrowers. It is expected that this program will greatly improve the livelihood of the surrounding communities.

This phase will cover a three year period, during which Sierra Tropical Ltd. will spend about or more than US \$15.0 million in capital investment for this expansion. The capital investment will cover construction of a cannery and can plant, installation of frozen fruit equipment and other processes as seen possible, continuation of the nursery operations, preparing land of the expanded areas, purchase of more agricultural equipment and power generators and expansion of water supply, water treatment plant, offices, first aid facilities and staff housing.

This second phase of the project will see significant increase in areas to be leased for commercial plantings, in total workers to be employed, in total production of finished pineapple products and in total spending. The first year of Phase 2 will see an increase in preparing beyond the 1,200 hectares of phase 1 and will generate employment of about or more than 1,000 direct and indirect employees which will mostly be Sierra Leoneans. As in Phase 1, hundreds more of Sierra Leoneans will be employed by the company's contractors hired to continue the expansion of its capital projects.

For the second year of Phase 2, total area to be leased will increase to a minimum of 2,500 hectares. About 1,700 workers to be employed directly or indirectly of which will mostly be Sierra Leoneans. There will be about 45,000 metric tons or more of pineapple to be harvested

and processed. The third and last year coverage of Phase 2 will involve leasing a minimum of 3,000 hectares. Sierra Tropical Limited will generate employment of about or more than 1,900 direct and indirect employees, About 65,000 metric tons or more of pineapple will be harvested.

Training will continue to develop potential Sierra Leoneans who will occupy supervisory positions as the company continues to expand beyond Phase 2. Training programs will also continue in all areas of the company operations to develop new workers to be employed and improve skills of present ones.

During this second phase, company-hired contractors will be employing hundreds more of Sierra Leoneans for the completion of the expansion projects.

In summary, at the end of Phase 2, Sierra Tropical Limited will have 4,000 or more hectares planted to pineapple. It will produce pineapple juice, concentrate, canned pineapple, frozen pineapple and fresh pineapple to be sold to markets in Europe, United States, Middle East and Africa.

Phase 3 and beyond —Long Term Goal

Sierra Tropical Limited intends to expand its operations further in this phase, depending on the success of Phases 1 and 2. The Company's long term goal is to develop and establish a large scale agriculture and manufacturing operation in Sierra Leone. It aims to be the major exporter of high quality and competitive pineapple and other tropical fruit products in the major markets in Europe, United States, Middle East and Africa. It also aims to be a major source of employment in the country, continuously developing and improving the skills of Sierra Leoneans.

Lastly, the smallholder or out grower scheme will play a strategic role for Sierra Tropical Limited to achieve its long term goal. The company aims to source a good portion of its pineapple requirements from the growers. In this regard, Sierra Tropical Limited will continue to develop an effective growers program by providing growers with adequate training and appropriate technical assistance in cultivating and growing tropical crops. The company's growers program aims to continuously create more local employment and greatly improve the livelihood of the communities.

2.1 Specific Project Components

2.1.1 Phase 1

2.1.1.1 Nursery Development

The first phase of the project involves the establishment of new land to plant mother material to generate new planting material. There are generally several steps involved in this which is described below:

1. Survey

New land is identified for its suitability. Slopes, soil structure, chemical makeup etc. are all considered in the selection. Once an area is selected, the first important aspect is to survey the field and determine the slopes so that the water flow can be understood and necessary measures can be taken to improve drainage and reduce soil erosion. In Dole this process is usually done using an Unmanned Aerial Vehicle (UAV) or “drone”. Information from the images taken is used to create the necessary maps.




Figure 2.1-12.1-1: Field showing Natural contours of a selected field

2. Water Flow & Drainage

Several Maps are generated from the DOLE GIS system. The first is the water flow map which shows natural water flow and drainage.

Drainage channels are overlaid on the natural colour image or mosaic. This clearly shows the watershed which runs approximately in an arc from North to South, creating the drainage

pattern shown. The western half drains into the natural drainage gully on the western side of the field, while the eastern side drains into the eastern gully.




Figure 2.1-22.1-2: Water flow and natural drainage pattern

3. Field Design

Using this information and the GIS software the layout or intended direction of rows (or blocks) can be added taking into account the natural slope of the field. An accurate map indicating the field layout is then prepared for the land preparation team to rip, bed forming and make drains all respecting the natural contours and slope of the field as best as possible.





Figure 2.1_32.1-3: The field layout is adapted to the relief. The center road follows the watershed and the orientation of the field blocks is approximately parallel to the contour lines.

Once land preparation starts steps in the following flow diagram are followed. As the project will be starting on new land, certain practices are modified, for example knock down and shredding may be different. Table 2.1-1 is an indication of the type of process which will be adapted for local requirements.

Table 2.1-1: Land Preparation Flow Diagram



Nvandehun Nursery Site Sierra Rutile Mine

Background

As part of the feasibility studies and preparations for establishing this project, an agreement was reached between Sierra Rutile Ltd (SRL) and African Lion Agriculture (ALA) in 2015, for the establishment of a 2ha test plot for the Sierra Tropical proprietary variety of pineapples. The test plot aimed at testing the suitability of the Sierra Tropical variety to

conditions in Sierra Leone. The test was conducted by ALA within the scope of their EIA Licence, as part of a Corporate Social Responsibility (CSR) project for SRL.

In early 2016, a further agreement was reached between SRL and ALA to convert the existing pineapple farm into a nursery for the production of planting materials (using the proprietary variety) for the proposed site at Sumbuya. In 2017 however, SRL was purchased and taken over by Iluka Resources. The agricultural areas were subsequently handed over to Carmanor, a new company not covered by an EIA Licence. This plot of land is now to be included in Sierra Tropical's EIA Licence.

The nursery was planned and intended as a temporary alternative to grow and speed up the provision of planting material for the Sumbuya Project. As the plants' life cycle is completed the area will be cleared and returned to Carmanor for their continued use. It is expected that this will be achieved and the nursery cleared away by the end of 2019.

Description of Nursery Site

The following table highlights key features of the nursery site:

<u>Location</u>	<u>Nyandehun village, Pirri Chiefdom, Bonthe district</u>
<u>Point Coordinate (centre of land area)</u>	<u>7° 39'05.50" N, 12° 18'36.37" W</u>
<u>Land area</u>	<u>30 hectares</u>
<u>Work force size</u>	<u>133</u>
<u>Capital Outlay</u>	<u>USD 1.43million</u>

Commented [VJ1]: EPA will request for boundary coordinates; at least four points

Commented [VJ2]: The narrative states 2ha; an explanation needs to be provided on how the land area increased over time.




Figure 2.1-4: Pineapple Farm Plots (Previously owned by ALA)

Nursery Operations

The project and cultural process, such as field planning and layout, cultural practices, variety of plants, fertilizers and chemicals used are the same as have been described for the Sumbuya Project. This nursery is comparatively much smaller than the Sumbuya Project, and developed on a previously established farm/field.

The nursery involves 3 distinct stages as follows:

Stage 1 – Plant Crop (PC)

This refers to the first cycle of the plant's life. Sucker material is planted in raised beds in cultivated fields. There are on average 80,000 plants per ha. It takes 10 to 12 months for the plants to grow to a size large enough for them to be induced to flower. This is the stage at which some of the fields are at the Nyandehun nursery.

Once induced, the plants will produce a flower. A simple growth regulator (registered in the USA and suitable for this purpose) is used to stop the reproductive process (flowering) and change the plant back to vegetative growth. Instead of fruit, it forms propagules or suckers. These are allowed to grow to a certain size after which they are harvested. This stage lasts for 9 to 14 months depending on growing and environmental conditions.

Commented [VJ3]: Name would need to be provided and here and MSDS made available. Please let me know if it is among the Chemical lists already provided

Stage 2 – Ratoon Crop (R1)

Once all the suckers are harvested for plant production, one sucker is left behind to create a second mother plant. The process mentioned above is then repeated. This stage takes a further 8 to 10 months, depending on growing and environmental conditions.

Stage 3 – Knock Down/Clearing

After all the propagules/suckers have been harvested the plants are removed or incorporated into the soil and the process starts again following land preparation if required. Minimum tillage is practiced by Dole in order to keep cultivation to a minimum.

The full cycle of a pineapple plant takes 36 to 40 months depending on growing and environmental conditions.


Water Requirements

Water is sourced from the SRL lagoons based opposite the pineapple nursery area. This water is tested by the mine on a regular basis and is also suitable for use as irrigation water. Water requirements of the plants are as described for the Sumbuya Project, as the soil composition is very similar.

Chemical Requirements


Chemicals and Fertilizers used at Nyandehun are all registered and approved for use on Pineapples. They are also approved by Dole for use in their worldwide operations. The list is exactly the same as that provided for the Sumbuya Project.

2.1.1.2 Organizational structure



2.1.1.3 Construction Work

Construction work will be an integral part of the developmental activities during this first phase of the project. It will involve the development of offices, warehouse, support facilities, security posts, wells and water tower, the facility boundary wall and living quarters.



Different construction contractors will be selected through a bidding process following the acquisition of the EIA License for the different projects.

2.1.1.4 Water Supply

Water for the project will be sourced from boreholes and from the Sewa River. Boreholes will be sunk to provide water for the day to day running of the facility including domestic use. Water for irrigation and spraying will be obtained from the Sewa River; the following table shows the average calculated water requirements for the project.

Water required (in terms of precipitation)	7mm/day
Continuous water flow required in litres/second/ha	0.81lt/sec./ha
Total equivalent ha	500 ha
Total irrigation water requirements (maximum)	875,000 m ³ per year

2.1.1.5 Power Supply

Power supply will be used for running the offices and irrigation purposes. Several diesel generators will be used to supply the project's power requirements initially. It is estimated that a total of 1 – 1.5MW of electricity will be required to power the project initially; in the longer term as the project progresses, demand may increase to beyond 3MW.

There are however plans to look into other more environmentally friendly sources of energy as the project progresses. Options are Hydroelectric, Solar, Biomass for combustion and the use of pineapple/fruit waste to produce methane gas. For Methane gas production a sufficient amount of biomass would be required to be available on a regular basis to sustain energy production. This could be possible after 2023. All of these options will be investigated and pursued later, we look to reduce Diesel generation and consumption to a minimum and have it as a backup capacity. The following table highlights the total expected power requirements to supply irrigation needs assuming a pump rate of 50m³/hour (Pump head of 60m, pump and motor efficiency of 70%):

Estimated maximum water application over a season (in terms of precipitation)	1000 mm
Expected kW	12
Estimated running time	1500 hours
Estimated number of pumps	3
Total annual estimated Power requirements	36kW

2.1.1.6 Chemical Usage

The following list highlights the full range of chemicals approved for use in Dole's operations. It is not likely that most of these chemicals will be used, but at this stage it is difficult to identify exactly which ones will be required for the project. The Material Safety Data Sheets for these chemicals are compiled in Volume 3 of this report: List of DOLEFIL-Approved Agrochemicals and their MSDS.

Table 2.1-2: Fertilizers

Fertilizer Material	Element	%
Urea	N	46
Ammonium Sulfate	N	21
Monoammonium Phosphate	N	12
	P ₂ O ₅	60
Diammonium Phosphate	N	18
	P ₂ O ₅	46
Potassium Sulfate	K	42
Sulfate of Potash	K	50
Potassium Chloride	K ₂ O	60
Muriate of Potash	K ₂ O	22.1
Potassium Magnesium Sulfate	Mg	11.2
	S	22.5
	Ca	25
Dolomite	Mg	11
	Ca	11
CalMag	Mg	7
	Ca	27
Calcium Chloride	Ca	19
Calcium Nitrate	N	15.5
	Ca	25
Amino Acid chelated Calcium-Boron	B	12
	Ca	5
Magnesium Sulfate, Anhydrous	Mg	20
Magnesium Sulfate, Heptahydrate	Mg	10
Molybdenum Amino Acid Chelate	Mo	10-20
Zinc Sulfate Heptahydrate	Zn	20
Ferrous Sulfate Heptahydrate	Fe	20
Sodium Tetraborate Pentahydrate	B	15
Elemental Sulfur	S	99.9
Phosphorous Acid (Nutriphite P Soil 0-60-0)	H ₃ PO ₃	70

Table 2.1-3: Pesticides

Product Name	Active Ingredient	%
Insecticides		
Lorsban 40 EC	Chlorpyrifos	40
Vessel 400 EC	Chlorpyrifos	40
Pyrinex 40 EC	Chlorpyrifos	40
Diazinon 60 EC	Diazinon	58.8
Magister 200 SC	Fenazaquin	18.2
Siege Pro 73 G	Hydramethylnon	0.73
Cosavet DF	Sulfur	80
Fungicides		
Aliette 80 WP	Fosetyl- Al	80
Ridomil Gold 480 SL	Metalaxyl-M	48
Cuprofix WG	Metallic Cu	20
Tilt 250 EC	Propiconazole	25
Herbicides		
Fezpax 80 WP	Ametryne	80
Hyvar-X	Bromacil	80
Diuron 80 WP	Diuron	80
Onecide 15 EC	Fluazifop p-butyl	15
Pounce 480	Glyphosate IPA	48
Assure II EC	Quizalofop p-ethyl	10.3
Surfactants		
Enspray N	Paraffinic oil	100
Silwet 408	Polyalkyleneoxide Modified Heptamethyltrisiloxane	70-100

Table 2.1-4: Plant Growth Regulators

Product Name	Active Ingredient	%
Maintain CF125	Chlorflurenol methyl ester	12.5
Ethrel	Ethephon	48
Xtragro 480 PGR	Ethephon	48
Leadthrel	Ethephon	48
Ethylene	Ethylene	98.9
Gibberellic Acid, Technical grade	GA ₃	90
Fast Gro Cytokinin	Kinetin	0.04
X-Cyte		0.04
Ericin	Etephon	43.3

Table 2.1-5: Post Harvest Chemicals

Product Name	Active Ingredient	%
Aluminum Sulfate	Al ₂ (SO ₄) ₃ .14H ₂ O	100
Calcium Hypochlorite/ Hypochlorous Acid	Ca(OCl) ₂ .2H ₂ O	100
Ethyl formate	Ethyl formate	100
Decco Lustr 444	Oleic acid, Propylene glycol, Sodium monostearate, Ethoxylated	<5 <50 10-40

2.1.1.7 Waste Management

The main types of waste expected to be generated from the project are as follows:

- Agricultural waste - biomass
- Domestic Waste – generated by day to day activities of workers (paper, plastics, food, etc.)
- Hazardous Waste – left over/unused chemicals (fertilizer and pesticide), chemical packaging, etc.
- Sewage

During this initial phase of the project, there will be no processing of fruits; this will feature during the second phase following the introduction of the cannery.

Agricultural Waste:

This category of wastes will comprise of plant waste (from vegetation clearing, pruning, etc.) and will be shredded (using a wood chipper) and used as compost or the vegative material will be Mulched user a forestry mulcher and incorporated directly back in to the soil, improving the organic material content. Care will however be taken to ensure that these wastes do not become a haven for rodents and other pests. Burning of biomass will be avoided unless absolutely necessary.

There are proposals to be reviewed for the use of biomass generated by the project to produce electricity to supplement or replace the diesel generators. There are two options under consideration 1) is a dry biomass burner for steam generation, burning rice hulls or other materials this is not possible at this time, 2) the other system of Biomass is for the production of Methane gas for steam production and electrical generation- that is a separate investment to be reviewed at a later date.

This will however only become feasible when the project has developed and expanded to a point where sufficient biomass is produced to support energy production.

Domestic Waste:

Domestic waste will be produced in offices and day to day activities in other areas of the facility. Waste minimization, reuse and recycling will be employed as much as possible, and workers will receive training in these waste management methods. Waste receptacles will be labeled to collect different types of wastes which can then be disposed of appropriately.

Hazardous Waste:

Hazardous wastes will include items such as waste oils, fuel filters, pesticide/fertilizer containers and packaging. These will be disposed of in assigned and labeled waste receptacles and stored safely for transportation to and disposal at a hazardous waste management facility. Staff will be trained in the safe handling and disposal of these wastes.

Sewage:

Grey and black water will be directed into underground soak away pits which will be emptied by a licenced sewage management company when required. Such a company will be identified at a later stage. It will be ensured that sewage disposal regulations are adhered to.

2.1.1.8 Occupational Health, Safety and Security Issues

Dole is committed to providing its employees with a safe and healthy work environment. This includes providing appropriate protective equipment, as well as following good manufacturing practices and taking proper safety and sanitation measures. By complying with

applicable environmental and occupational health and safety laws and regulations, a safe working environment will be ensured.

In order to uphold the Company's commitment to a safe and healthy workplace, management and employees will be required to:

- Follow all safety laws and procedures
- Observe posted safety-related signs
- Use prescribed safety equipment whenever required

Management and staff will be required to work together to prevent hazardous or unsafe working conditions. Each staff will be responsible to follow prescribed safety and reporting procedures if any hazardous conditions or unsafe behaviour is detected.

Use of alcohol and illegal drugs will not be accommodated and employees will be prohibited from possessing, distributing, selling or using these items while on the Sierra Tropical Premises.

To further ensure a safe work environment, acts or threats of violence will not be tolerated. Any threatening behaviour will be addressed immediately with utmost seriousness. Weapons will not be permitted on the premises.

2.1.1.9 Environmental Health and Safety

It is STL's policy to comply with all applicable laws and regulations at all times and to take all practicable steps to promote health, safety and environmental protection. STL's parent company, Dole, manages operations worldwide that belong to different economic sectors – farming, food processing, manufacturing, research, transportation (including trucking, shipping and managing port facilities), distribution and sales. In all these operations, it is Dole's goal to prevent adverse effects on health, safety and the environment and subsidiaries are required to operate within the same standards.

Dole strives to develop and employ approaches that are most appropriate and effective under local conditions and are guided by: scientific research and knowledge; principles of risk analysis; public, community and worker concerns; and regulatory policies and standards of Japan, the U.S., the European Union and international organizations such as the World Health Organization.

Dole's farming operations utilise sustainable agricultural practices and integrated pest management methods that employ biological and agricultural approaches to controlling pests and plant diseases. Crop protection products are used only when and where necessary, and always with the proper care and in accordance with applicable laws. Dole and its subsidiaries will not use any product banned for reasons of unacceptable health or environmental risk by the United States Environmental Protection Agency, Japan or the European Union.

2.1.2 Phases 2 and 3

2.1.2.1 Outgrower Program

The Outgrowers scheme will play an important role for Sierra Tropical Ltd. in achieving its longer term goals. The company may source a good proportion of its pineapple and other fruit requirements from local growers in and around Sumbuya, the site of the new plantation.

Sierra Tropical Ltd will, like it has done in its Philippine operations, continue to develop an effective growers program. This will involve the training and appropriate technical assistance along with access to necessary equipment and material inputs to cultivate and produce quality fruit. Other tropical fruits such as a Papaya (pawpaw), Mangoes, Guavas and Passion fruit will also be considered as the project develops.

The outgrower's program aims to create more local employment opportunities, improve the livelihood of farmers and greatly improve the economic wellbeing of the local communities in the area.

Main Objectives:

- Increase tonnages for the cannery
- Improve global competitiveness
- Improve the farmer's income and develop farmer entrepreneurship skills
- Generate new employment opportunities for the community near to their homes
- Create a new social landscape in Sumbuya

Rational:

- There is an expanding world market creating additional needs for our products
- There are well suited growing land/soils in Sierra Leone
- There is well established production technology and technical knowledge available from the company
- The company has many years of experience working with small commercial farmers in different countries around the world.

Potential Benefits To The Growers and their Communities

- Out Growers will have a firm commitment for their contracted production
- Access to well tested and proven tropical fruit growing technology
- Access to low cost and a reliable supply of agricultural inputs as part of a contracted grower program.
- Access to quality training and disciplined systems/structures
- Higher earnings versus current crop alternatives (i.e. Cassava)
- Employment of family members and neighbors who meet the qualifications required.
- Sound company relationships with the growers and community through community outreach programs (i.e. Improvements to clinics and schools / infra-structure improvements, i.e. water system, sanitation and others)

- Improved farmers income, bring new opportunities for economic activity to the community
- The relationship between outgrowers and local government representatives, Chiefs and government line agencies will be enhanced.

2.1.2.2 Development of Cannery

The development of a cannery is anticipated to commence in 2019. The cannery will be used to process the fruit into various forms including canned fruit, fruit juice, juice concentrate, etc.

The development of this aspect of the operations will involve the following aspects:

- Fruit Receiving Area
 - Dumper Truck or trailers
 - Receiving Conveyor
 - Small flume (to wash fruit)
 - Fruit Elevator with Washer Spray
 - Feed Conveyor
 - Fruit Grader
 - Fruit Accumulator
 - Fruit Distribution merry-go-round (MGR)
 - Truck Scale System
- Four (4) Ginaca Pineapple Processing Machines intended for juicing
- Two (2) preparation lines with slicers, handguns
- Four-stage juice press system
- 25-GPM 6-effect tube -type evaporator and aseptic filling system expandable to 50 gpm
- Boiler at 20K pounds per hour, saturated steam.
- Reverse Osmosis water at 55,000 liters per hour - source would be deep borehole
- Cold storage for the juice concentrate
- 3 units 350 KVA continuous and stand by generators
- A laboratory
- Waste Water Treatment Plant
 - 2 settling ponds
 - 2 aeration ponds
 - 3 facultative ponds
- Fire Protection System

2.2 Timescale of Operations

Phase 1 will last approximately 2 years, and will utilise a minimum of 800 hectares. The second phase will be implemented in the third year and will last for 3 years, with a proposed expansion to a minimum total of 4,000 hectares. For years 6 to 10, more than 2000 hectares roughly maybe be acquired, and with the proven success of Phase 2, expansion will continue to increase steadily. Using this projection, and assuming success at each stage, the project is estimated to continue for over well over 50 years.

3 ANALYSES OF PROJECT ALTERNATIVES

In accordance with current ESIA good practice, it is appropriate for the study to review alternatives considered during planning of the project, and to explain why the proposed project activities have been selected, including potential environmental, social and advanced technological considerations. For the purpose of the agro-processing (plantation, product processing & marketing) activities for which this ESIA is being conducted, analysis of 'No Project Option' and choice of project site have been considered.

3.1 The "No Project Option"

The project area is virgin ground for the agro-processing industry and the Sierra Tropical Ltd project will be first large scale mechanized cultivation and tropical fruit processing project in the southern region and Sierra Leone.

The project will provide opportunities for socio-economic development and growth within the Lugbu Chiefdom. This will come about as a result of land lease payments, outgrower programs, job opportunities, business opportunities, community development activities etc. Infrastructural development within the project area will contribute towards improving the standard of living and improve the social conditions within the related communities.

National benefits will be realized through the socio-economic improvement of the project area as well as the through the exportation of the fruits and fruit products (juice, canned fruit and juice concentrate)

The socio-economic assessment carried out during this study indicated that currently in the absence of an Industrial Project such as Sierra Tropical Ltd's, most community residents (over 85%) depend mostly on small scale farming with the next most popular livelihood being artisanal mining (6%). On a district level, more than half of the population of Bo District depends on farming as their main source of livelihood. Thus economic growth and poverty reduction in the district will only be sustained with development in this sector as the District is endowed with cultivable land (Upland and Inland Valley Swamps) and adequate rain-fall.

The implementation of the Sierra Tropical agro-processing will therefore have far reaching benefits to the project area and Sierra Leone as a whole, which would not be realized if the project were not to be implemented. Sierra Tropical Ltd's decision to implement this project in an environmentally sustainable manner will have greater benefits for the local communities and Sierra Leone as a whole, than not implementing the project and maintaining the existing biophysical, socio-economic and physical conditions at the site.

3.2 The Project Site

Dole Worldwide Packaged Foods has projects in different parts of the world including the Philippines and Thailand. The company became interested in establishing a project in West Africa due to the suitable soil and climate, and the fact that this region is at minimal risk of typhoons or hurricanes. The proximity to the USA, Europe, South America and the Middle East markets was also taken into consideration.

In specifically selecting Sierra Leone for the project, the following factors were taken into consideration:

1. Suitable Climate and Soil for PA - Confirmed through exploratory Soil survey, Water and Climate assessment
2. Enough land for the project can be acquired at a reasonable cost.
3. The project can be established at a competitive running cost; labour is also available and competitive.
4. Political Stability and Public Safety
 - Experienced democratic election twice peacefully (2007/2012) and current government is stable.
 - No tribal/religious conflict in Sierra Leone.
 - Many Foreign Investors recognize political stability and Public Safety of Sierra Leone.
5. Attractive Incentives by Supportive Government

The Lugbu Chiefdom was chosen due to the proximity of the Sewa River, good rainfall and encouraging preliminary soil surveys.

3.3 Project Technology & Operations Option

Sierra Tropical has plans to adopt the most modern technology in the cultivation, processing and marketing of the tropical fruits. This will ensure that maximum yield and production output is obtained, resulting in a thriving local and international market.

During the second phase on the project, STL intends to install a fully mechanised Cannery where the fruit will be processed. The equipment utilised will provide an extraction efficiency of about 90% (this refers to the percentage extracted from the total juice present in the fruit).

The fully industrialised agricultural and processing methods to be used will be a vast improvement from the existing rudimentary farming methods used within the project area.

4 POLICY, LEGAL, REGULATORY and INSTITUTIONAL CONTEXT

4.1 Policies and Plans

4.1.1 National Environmental Policy, 1994

This National Environmental Policy seeks to achieve sustainable development in Sierra Leone through the implementation of sound environmental management systems which will encourage productivity and harmony between man and his environment. It also promotes efforts which will prevent or eliminate damage to the environment and biosphere and stimulate the health and welfare of nationals, and serves to enrich the understanding of ecological systems and natural resources important to the Nation. Thus the key objective of the policy is to secure for all Sierra Leoneans a quality environment that can adequately provide for their health and well-being.

The policy takes into consideration major sector goals and policies for enhancing sustainability in environmental management systems. The following sectorial policies are highlighted within the National Environmental Policy:

- Land Tenure, Land Use and Soil Conservation;
- Water Resources Management;
- Forestry and Wildlife;
- Biodiversity and Cultural Heritage;
- Air Quality and Noise;
- Sanitation and Waste Management;
- Toxic and Hazardous Substances;
- Coastal and Marine Resources;
- Working Environment (Occupational Health and Safety);
- Energy Production and Use;
- Settlements, Recreational Space and Greenbelts;
- Public Participation;
- Quality of Life;
- Gender Issues and the Environment;
- Institutional and Government Arrangements;
- Legal Arrangement.

Subsequent to this policy is the Environmental Protection Act of 2008

4.1.2 Agricultural Policy

The policy recognizes that growth in agriculture is crucial for achieving government's development and poverty reduction objectives. The policy deals with pertinent issues of crop production, crop post-harvest technology, livestock production, produce marketing and pricing, agricultural finance/credit, agricultural research, agricultural extension, mechanisation, land and water resources management, food security, natural resource management, biotechnology and genetic engineering as well as other policy issues. It also establishes significant guidelines for the participation of various stakeholders in the agricultural sector to promote sustained growth. The policy makes reference to land tenure situational factors and seeks to ensure compliance amongst different stakeholders by setting up a "Land Commission" to investigate and make recommendations on land tenure for accommodating the demands of commercial agriculture.

The government through this policy seeks to build, in the medium and long-term, technology pyramids to provide alternatives to the significant proportion of its foreign exchange used to purchase fossil fuels. The policy therefore provides for the use of existing or new crops for biomass production or conversion to biofuels to ensure sustainability in the agricultural processing sector and meeting the energy needs of the country

4.1.3 Draft National Lands Policy, 2015

This policy framework provides the vision, principles and policy components to give direction to and definition of the roles and responsibilities of various government and customary authorities, and other non-state actors, in land management. Specifically, it enunciates Policy Statements in respect of the key components of the National Land Policy such as access to land and tenure, land use, regulation and the management of special land issues, land administration structures, land laws and the Constitution.

The National Land Policy proposes to improve upon and strengthen the existing land administration systems and land laws, particularly by recognizing and working with the differentiated land tenure categories in the Western Area and the Provinces, and enhancing the capacities of relevant institutions on mobilizing sufficient national and international resources to ensure the implementation of the National Land Policy.

The specific objectives of this National Land Policy are:

- a. To clarify the complex and ambiguous constitutional and legal framework for sustainable management of land resources;
- b. To promote law reforms that will further harmonize the two separate jurisdictions of the current land tenure systems;
- c. To ensure the security of tenure and protection of land rights to all legitimate landholders, regardless of their form of land tenure;
- d. To promote equitable access to land

- e. To promote and enforce sound land use, regulation and management
- f. To streamline and decentralise land administration to be more efficient, transparent and effective
- g. To modernize and streamline land information system
- h. To promote the eradication and/or avoidance and efficient settlement of land disputes by rationalising and strengthening the capacity of traditional institutions, local and national courts in the speedy and effective resolution of land disputes;
- i. To ensure protection and security of Sierra Leone national boundaries in accordance with international conventions as enshrined in the International Law of the Sea, Anglo-Francophone Protocols and Joint Border Commissions;
- j. To build capacity for the effective monitoring and evaluation of the implementation and impact of the national land policy.

4.1.4 Forestry Policy, 2010

The Forestry Policy support relevant provisions of the Constitution which permits restrictions on activities within forests which is reasonably required in the interests of conservation of the natural resources, the respect for international law and treaty, obligations, as well as the seeking of settlement of international disputes by negotiation, conciliation, arbitration or adjudication.

This Forestry Policy also supports strategies outlined in the Framework for Effective Management of Natural Resources.

4.1.4.1 Biodiversity Strategic Action Plan, 2003

The Sierra Leone Biodiversity Strategic Action Plan comprises a series of measures and mechanisms intended to conserve and promote the sustainable use of the different components of the country's biodiversity. The actions proposed cover several key thematic areas under: terrestrial biodiversity, inland water ecosystems, forest biodiversity, marine and coastal biodiversity and agricultural biodiversity. In addition, actions are also proposed for key cross-cutting issues affecting the sustainable utilization of biodiversity, including: policy, legislation and institutional review, capacity building, identification and monitoring, sustainable use, incentive measures, research and training, public education and awareness, regulation of access to genetic resources, protection of indigenous knowledge and intellectual property rights of local communities, technology transfer and handling of biotechnology and exchange of information and technical co-operation.

This Action Plan is intended to:

- Provide a framework for setting priority policies and actions for the conservation and sustainable use of biological diversity in Sierra Leone;

- Catalyze and provide guidance for legal policy and institutional reforms necessary to achieve effective conservation and sustainable use of biological diversity;
- Enhance the planning and co-ordination of national efforts aimed at the conservation and sustainable use of biological diversity;
- Guide the investment and capacity building programmes for the conservation and sustainable use of bio-diversity;
- Facilitate information sharing and coordinated action among the various stakeholders at the national level and foster scientific and technical cooperation with other countries and international organisation.

4.1.5 Conservation and Wildlife Policy, 2010

The Conservation and Wildlife Policy (2010) was developed in recognition that the previous wildlife conservation policy was in need of modernisation. Current legislation based on the 1972 Wildlife Conservation Act (as was the case of the previous wildlife conservation policy) does not reflect the advances made in biodiversity conservation in the past four decades; it also does not take into account international obligations that arose after its entry into force, such as the Convention on Biological Diversity (CBD), the Convention on International Trade in Endangered Species (CITES) and the United Nations Framework Convention on Climate Change (UNFCCC).

The Conservation and Wildlife Policy identifies that challenges to biodiversity conservation in Sierra Leone result from a lack of knowledge due to “recent conflict, land use change, uncontrolled exploitation of natural resources, and a lack of recent comprehensive inventory”. The vision of the policy document is to establish “an integrated wildlife sector that achieves sustainable, rights-based management of wildlife resources for biodiversity conservation inside and outside wildlife conservation areas which benefits present and future generations of Sierra Leone and humankind in general.” The policy presents a plan for biodiversity conservation based on a set of “policy statements” outlining concrete policy goals and develops the necessary institutional arrangements for policy implementation.

4.1.6 Disaster Management Preparedness Plan, 2006

As part of its post-war recovery effort, the Government of Sierra Leone reviewed its National Security Structure to meet the demands of the 21st century. This led the Government to enact the National Security and Central Intelligence Act in 2002 thereby mandating The Office of National Security to be ‘the Government of Sierra Leone’s primary Co-ordinator for the management of national emergencies such as disasters both natural and man-made’.

The disaster management Plan, 2006 is a comprehensive approach that enhances increased political commitment to disaster risk management, thereby encouraging government agencies to take the lead and supported by non-governmental organisations. It also promotes public awareness and the incorporation of disaster risk management into development planning. The policy highlights the sources of funding and the reduction of bureaucracies in accessing such funds for effective disaster co-ordination.

The Policy document emphasizes the following:

- Ensure the integration of disaster risk management into sustainable development programmes and policies to ensure a holistic approach to disaster management;
- Ensure priority and requisite institutional capacities for disaster risk reduction at all levels;
- Enhance the use of knowledge, education, training, innovation and information sharing to build safe and resilient societies;
- Improve the identification, assessment, monitoring and early warning of risks;
- Improve effectiveness of response through stronger disaster preparedness.

4.2 Legislation

Laws and regulations governing environmental issues are found as Acts of parliament. The pieces of legislation of the various government line ministries or institutions include:

4.2.1 Environment Protection Agency Act, 2008/2010

The EPAA 2008 is the government of Sierra Leone's overarching legislation that deals with the protection of the environment. The Environment Protection Agency was established with a Board of Directors set up as its governing body. Subject to this Act, the control and supervision of the Agency is the responsibility of the Board, whose administrative functions as stipulated by the EPAA, 2008 include the following:

- i. Promoting effective planning and the management of the environment;
- ii. Coordinating and monitoring the implementation of national environmental policies relating to Sierra Leone;
- iii. Providing policy guidance and advice to ensure the efficient implementation of the functions of the Agency so as to enhance its overall performance;
- iv. Facilitating co-operation and collaboration among Government Ministries, local authorities and other governmental agencies, in all areas relating to environmental protection;
- v. Coordinating environmentally related activities as well as serving as the focal point of national and international environmental matters, relating to Sierra Leone.

4.2.2 The Forestry Act, 1988

The Forestry Act, which first came into effect on 1st July 1988, mandates the Forestry Department to take steps to ensure compliance with the provisions of the Act. It mandates the Director of Forestry or his representatives to enforce the legislation.

The following activities in a forest reserve are considered offences under Sierra Leone forest laws:

- Establishing or carrying on a forest industry in or with resources of the core forest: charcoal burning, wood cutting, hunting, stone or soil deportation
- Clearing, cultivating or breaking up land for any reason
- Removing soil, sand or gravel
- Erecting a building or shelter in the core forest or its buffer zone
- Lighting, keeping or carrying fire
- Carrying a firearm, pasturing cattle or permitting them to trespass
- Damaging, altering or removing any notice board, land-mark or fence
- Assaulting or obstructing any person carrying out his/her duty under the Forest Act e.g. Forest Guards
- Altering, defacing or obliterating any mark placed on timber by a forest officer

The maximum penalty under the Forestry regulations is Le 5,000,000 or one year imprisonment. Once convicted, an offender can also lose the equipment that was used to carry out the offence, as the court may order that it be forfeited to the state in addition to the imposition of a fine or term of imprisonment.

4.2.3 Land Tenure and Ownership

Land administration in Sierra Leone is governed by a dual system of law, dispersed in about twenty statutes and regulations.

In the Western Area of Sierra Leone, land tenure is governed by Property Statutes. Land is either State (publicly) owned or privately owned. The right of the state to public land is inalienable and indefeasible. Rights of occupation over public land may be granted under warrant. The state has the power, conferred by the Unoccupied Lands Act, Cap 117, to take possession of unoccupied land.

In the provinces, customary law co-exists with statutes. The recognition of the force of customary law in the provinces is established by section 76 (1) of the Courts Act 1965.

Through customary law, ownership of land is vested in the chiefdoms and communities; and can never be owned freehold. Land always belongs to the communities under the different forms of tenure under customary law. This principle is established by the Chiefdom Councils Act as well as by Section 28 (d) of the Local Government Act 1994.

4.2.4 Fisheries Act, 2007

The Fisheries Act of 2007 provides protection for both fresh and marine water species as classified by IUCN with the Sierra Leone water. It defines clearly where commercial vessels could harvest-Exclusive Economic Zone (EEZ) and where artisanal fisheries operations could exploit – Inshore Exclusive Zone (IEZ).

4.2.5 Wildlife Conservation Amendment Act, 1990

The Wildlife Conservation Act, 1972 and the Forestry Act, 1988 are the main legislations that deal with issues of Biodiversity Conservation in Sierra Leone. It provides for the establishment, conservation and management of National Parks, Game Reserves and other forms of Natural Reserves.

Specific provisions dealing with the protection, management and conservation of these areas and the limitations therein are highlighted in Part II of the Act and include the following:

- Prohibition of all forms of hunting, capture and other activities leading to the injury of wild animals;
- Destruction of any plant form by any means including fire;
- Fishing within these protected areas;
- Erection of structures, construction of dams, forestry, agriculture, mining or prospecting activities;
- Introduction of species from outside of the boundaries of the reserve.

The Wildlife Conservation Act of 1972 saw minor amendment in 1990 (known as the Wildlife Conservation Amendment Act), which included redefinition of terms, and other modifications and qualifications. For example, the prohibition of hunting of elephants which was limited to protected areas in the 1972 Act was extended to include all forests. The 1990 Amendment Act provided for change of name from Forestry Department to Forestry Division. Despite the minor amendment the Wildlife Conservation Act of 1972 along with the Forestry Act of 1988 continue to be the main legislature for biodiversity conservation in Sierra Leone.

The Wildlife Regulations of 1997 however makes provision for the acquisition of licences or permits for hunting in such designated areas and for other purpose as may be prescribed.

4.2.6 Factories Act – 1974

This Act became effective on the 30th May, 1974. It basically deals with health and safety measures as they concern the factory worker. It protects the worker through demands for all aspects of cleanliness, reports of all injuries, accidents, diseases and death.

Rules for the Implementation of the Act

As stated in section 16, the Minister may make rules for the effective implementation of this Act and such rules may provide:

- ❑ For the safety of persons employed in such trades and occupations as may be declared to be dangerous trades;
- ❑ For imposing obligations for the better safeguarding of persons against accidents from dangerous parts of any machinery;
- ❑ For the construction and maintenance of fencing to the dangerous parts of any machinery;
- ❑ For the proper maintenance and safe-working of raising and lowering machinery;
- ❑ For prescribing the qualifications to be possessed by engineers and other persons, for them to be placed in charge of, or entrusted with the care or management of any specified machinery;
- ❑ For the reporting of any occurrences at any works arising from, or in connection with, the use, maintenance or repair of any machinery;
- ❑ For the appointment of persons to hold enquiries under this Act, and prescribing powers and duties of such persons; and
- ❑ For the fixing of penalties not exceeding a fine of one hundred Leones or imprisonment for a term of six months or both such fine and imprisonment, for the contravention of any rule.

Safety, Security and Welfare of Employees

Part V of this Act, deals with the aspect of health and stipulates that every factory shall be kept in a clean state and free from effluvia arising from any drain, sanitary convenience or nuisance. This part of the Act also states that for overall safety of all employees, the factory must not be overcrowded, must be effectively ventilated, and provided with suitable lighting systems. Every care must be taken by the factory holder, to secure the health, safety and welfare of all employees.

Offences and Penalties

Part VIII of this Act emphasizes on offences, penalties and legal proceedings. Section 47, subsection 1 of this part, states that in the event of any contravention of the provisions of this Act or of any Regulation or Order made there under, the occupier or owner of the factory, shall, be guilty of offence under the Act.

Regarding offences for which there are no penalties provided, section 48 stipulates that, any person guilty of an offence under this Act for which no express penalty is provided by or under the Act, shall be liable to a fine not exceeding fifty Leones or to imprisonment for a

term not exceeding one month or both. If the contravention for which he was convicted continues, he shall be guilty of a further offence and liable to a fine not exceeding ten Leones for each day on which the contravention is continued.

Section 50 states that if anyone is killed, or dies, or suffers any bodily injury, in consequence of the occupier or owner of a factory having contravened any provision of this Act, the occupier or owner of the factory, shall, without prejudice to any other penalty, be liable to a fine not exceeding two hundred Leones or to imprisonment for a term not exceeding three months, or to both.

All offences committed under this Act shall, as section 56 states, be prosecuted in a magistrate court.

Powers of Inspectors

Section 14 of part IV of this Act states that an inspector shall, in executing this Act, have the power to do the following:

- ❑ To enter, inspect and examine a factory and its environs at any time, as long as he has reasonable cause to believe that explosives or any highly inflammable materials are stored or used;
- ❑ To take with him during an inspection, a police officer, if he has reasonable cause to expect any serious obstruction during the execution of his duty;
- ❑ To require the production of all documents and to examine and copy them in pursuance of this Act;
- ❑ To make necessary inquiries and examinations to ascertain whether the provisions of the Act are complied with; and
- ❑ To prohibit the use of any machinery, if he is reasonably of the opinion after examination, that it is not in good and safe condition.

If anyone wilfully delays or obstructs the Inspector in the exercise of any of his duties under this Act, then such a person shall be guilty of an offence and be liable to a fine not exceeding twenty Leones or to imprisonment for a term not exceeding one month or both. The occupier of the factory shall also be guilty of such an offence and be liable to punishment in like manner, even though he has not personally caused the obstruction.

Safety, Security and Welfare of Employees

Part V of this Act, deals with the aspect of health and stipulates that every factory shall be kept in a clean state and free from effluent arising from any drain, sanitary convenience or nuisance. This part of the Act also states that for overall safety of all employees, the factory must not be overcrowded, must be effectively ventilated, and provided with suitable lighting systems. Every care must be taken by the factory holder, to secure the health, safety and welfare of all employees.

As indicated in section 38, it is incumbent on the company to notify the District Inspector, in writing, of any accident or death in the factory. It is also stated in section 39 that all factory contracted diseases identified by a medical Practitioner, must be brought to the notice of the Chief Inspector in Freetown.

Section 40 states that: Where injury immediately results in death, the site of the accident must be left undisturbed after the removal of the corpse, until inspected by a police officer or an inspector.

On receipt of the report of an accident, the inspector shall if he considers it necessary or if directed by a higher authority immediately proceed to the scene of the accident, as indicated in section 41, and shall make enquiry into the cause of the death. This section further states that for the purpose of this enquiry, the inspector is free to use any one under oath, any document, and award fees for giving evidences, as may be fixed by the minister.

Any person, who, without reasonable cause, fails to comply with the terms of summons of the inspector, or refuses to be examined or to answer questions other than that which may incriminate him, or anyone who obstructs an Inspector or any person acting under his directions in the execution of his duty under section 41, shall be guilty of an offence.

The owner of every factory, according to section 45, must within 24 hours report in writing to an Inspector every dangerous occurrence caused by any machinery or electrical abnormality.

Section 26 of part VI stipulates that there shall be kept posted in a prominent position in every factory:

- ❑ The prescribed abstract of this Act;
- ❑ The address of the Chief Inspector and of the nearest Inspector; and
- ❑ Printed copies of any regulations made under any part of this Act which are for the time being in force in the factory; or the prescribed abstracts of such regulations.

4.2.7 Local Government Act, 2004

This Act deals with the establishment and operation of local councils around the country to enable meaningful decentralization and devolution of Government functions. It stipulates that a local council shall be the highest political authority in the locality and shall have legislative and executive powers to be exercised in accordance with this Act or any other enactment. It shall be responsible, generally for promoting the development of the locality and the welfare of the people in the locality with the resources at its disposal and with such resources and capacity as it can mobilize from the central government and its agencies, national and international organisations, and the private sector. The local council should initiate and maintain programmes for the development of basic infrastructure and provide works and services in the locality. A local council shall cause to be prepared a development plan which shall guide the development of the locality.

Many companies are bound to operate within areas controlled by one local council or another. There is also a relationship between the local council and the Chiefdom within which a company operates. It is advisable for companies to involve local councils in their development work. The schedules to the Local Government Act outline the activities of various MDAs that have been devolved to local councils.

4.3 Regulations

4.3.1 Forestry Regulations, 1990

These regulations are deemed to have come into force on the 1st July, 1990. The Chief Conservator holds the same responsibilities as he does for the Act of 1988.

As a method of environmental protection, it is stated in that no land between the high and low water marks, nor those above the high water mark on both sides of the bank of any waterway, covering a distance of one hundred feet (approx. 33m), shall be cleared of any vegetation except permitted by a clearance licence.

Sacred bushes are protected by the stipulated regulations of section 40, whereby clearance of vegetation from land designated as sacred bush, is prohibited except by clearance authority from the Chief Conservator.

4.3.2 Fisheries Regulations

National Fisheries Regulations such as the Fisheries Act 1988 and Fisheries Amendment Act 1990 respectively, have evolved over time in order to address specific matters relating to the conservation and management of natural resources within the marine environment.

The 1994 Decree, passed during the military government regime of NPRC, further established sufficient provisions for the conservation of Marine Resources. These range from monitoring, control and surveillance provisions, as well as those relating to enforcement.

Section 9 (1 & 2) of the Decree gives the government sovereign right over the Economic Exclusion Zone. They include rights for the exploitation, exploration, conservation and management of its natural resources. It further stresses the requirement for a written consent to be provided by government for any form of activities to be undertaken within this zone by states, international organizations or persons.

The Fisheries Act of 2007 provides protection for both freshwater and marine species as classified by the International Union for Conservation of Nature (IUCN), within Sierra Leonean waters. It clearly defines where commercial vessels are to harvest that is the Exclusive Economic Zone (EEZ) and where artisanal fishing is to harvest that is the Inshore Exclusive Zone (IEZ). It also stipulates the fishing equipment tolerable in Sierra Leone, and it also stipulates the quality and quantity of fish to be harvested.

4.3.3 Draft Wildlife Regulation, 1997

The Wildlife Regulation came in to force in 1997. It describes Wildlife Conservation Estate as areas described under the 1972 Wildlife Conservation Act as a National Park, Game Reserve, Strict Natural Reserve, Game Sanctuary or Non-hunting Forest Reserve. The regulation prohibits all unlicensed hunting with a Wildlife Conservation Estate, which includes the removal of honey. It prohibits the hunting of young and immature wild animals or birds; female wild animal accompanied by its young; and birds which are apparently breeding. It also prohibits hunting at night with lights to dazzle birds and animals.

The regulations stipulates that a license or permit should be sought before any form of hunting of game and bird can be done as required by Section 33 and 34 of the Act. The regulation also states that such licenses and permits can be revoked by the Chief Conservator of Forest if the holder fails to comply with the provisions of the regulations.

4.4 Institutional Context

4.4.1 Ministry of Water Resources

This ministry has the mandate for the development of policies and programmes for the provision of safe drinking water on a constant and sustainable basis to the entire population of Sierra Leone by carrying out activities under the following major headings:

- Guma Valley water company (GVWC);
- Sierra Leone Water Company (SALWACO);
- Development of Dams and other water supply schemes;
- Protection and management of water resources;

In relation with:-

- International Hydrological Association
- Collaboration with relevant Government Ministries and national and international organizations/Institutions

4.4.2 Ministry of Fisheries and Marine Resources

The Ministry of Fisheries and Marine Resources is the sole government agency with the legal mandate to promote aquaculture activities and to develop policies regarding the fishing sector, control and monitor fisheries and other aquatic resources within the territorial waters in Sierra Leone. The Ministry's overall responsibility is the control, development and conservation of all aquatic organisms, including:

- Protection of the marine and Fresh water environments;

- Prevention of the deposit of toxic and nontoxic waste in the territorial waters;
- Maritime Laws relating to territorial rights and fishing limits;
- Preservation of Marine Species.

4.4.3 Ministry of Agriculture, Forestry and Food Security

The Ministry of Agriculture Forestry and Food Security (MAFFS) is the central government agency responsible for promoting the development of appropriate policies and programmes geared towards the development of agriculture, animal husbandry and the attainment of national food security in the country.

4.4.4 Ministry of Lands, Country Planning and the Environment

This Ministry develops appropriate policies and programmes for lands country planning and the environment (role now limited with the formation of the EPA-SL) and carry out activities under the following major headings:

- Land and Land Tenure;
- State Lands;
- Surveys, Mapping and Triangulations;
- Relations with the Directorates outside Sierra Leone;
- Geodetic and Topographical Surveys;
- Enforcement of planning and building control;
- Demolition of unauthorized structures;
- Collaboration with relevant Government Ministries and with national and international organisations and Institutions.

4.4.5 EPA-SL

The Environment Protection Agency was set up to replace the National Commission for Environment and Forestry (NaCEF), which was mandated to oversee issues pertaining to the environment and forestry. The Environment Protection Agency was established with a Board of Directors set up as its governing body. This Board consists of a Chairman and representatives from the various line Ministries and a Unit as stated in section 3 of part II of the Environmental Protection Agency Act. Subject to this Act, the Board shall have the control and supervision of the Agency. The Agency shall act in liaison and co-operation with government agencies to control pollution and the general protection of the environment. The Agency, subject to this Act, shall promote effective planning in the management of the environment and coordinate and monitor the implementation of national environmental policies, relating to Sierra Leone.

4.5 International Conventions Policies, Codes, Protocols and Guidelines

4.5.1 International Conventions

Sierra Leone is a party to many international agreements, conventions, and protocols that seek to protect the environment and ensure sustainable development that are applicable to such a project. These include the following:

Table 4.5-1 : List of International Conventions and Agreements to which Sierra Leone is Party

International Commitments and/Agreements	Republic of Sierra Leone
The UN Framework Convention on Climate Change (New York, 1992)	Yes +
The Kyoto Protocol is a protocol aimed at fighting global warming. The Protocol was initially adopted on 11 December 1997 in Kyoto, Japan and entered into force on 16 February 2005.	Yes +
Ramsar Convention for the Internationally Important Wetlands Especially as Waterfowl Habitats (1971)	Yes+
The Basel Convention on the Control of Transboundary Movements of Hazardous Waste and Their Disposal (Basel, 22 March 1989)	Yes+
Convention on the Environmental Impact Assessment in a Transboundary Context (EPS, Finland, 1991)	Yes+
Stockholm Convention on Persistent Organic Pollutants (22 May 2001; has not come into force yet)	Yes+
Convention on Biological Diversity (Rio de Janeiro, 1992)	Yes+
Convention on the Protection and Use of Transboundary Watercourses and International Lakes (Helsinki, 1992)	Yes+
Convention on International Trade in Endangered Species of Wild Flora and Fauna (Washington, 1973)	Yes+
Aarhus Convention on Access to Information, Public Participation in Decision-Making and Access to Justice in Environmental Matters (1998)	Yes+
Rio Declaration on Environment and Development (UN	Yes+

International Commitments and/Agreements	Republic of Sierra Leone
Conference, 1992)	

The following sections briefly discuss the International conventions that have relevance to the STL Project.

4.5.1.1 United Nations Framework Convention on Climate Change

Sierra Leone ratified this convention on 22nd June, 1995. The objective of this convention is to regulate levels of greenhouse gas concentration in the atmosphere, so as to avoid the occurrence of climate change on a level that would impede sustainable economic development, or compromise initiatives in food production. The Parties are required to protect the climate system for present and future generations. Developing countries should be accorded appropriate assistance to enable them to fulfil the terms of the Convention. The Parties should work in cooperation, so as to obtain maximum benefit from initiatives in the control of the climate systems; The Parties are to prepare national inventories on greenhouse gas emissions, and on actions taken to remove them; formulate and implement programmes for the control of climate change; undertake cooperation in technology for the control of change in the climate system; incorporate suitable policies for the control of climate change in national plans; undertake education and training policies that will enhance public awareness in relation to climate change. The developed country Parties (and other Parties listed commit themselves to take special measures to limit their anthropogenic emissions of greenhouse gases, and to enhance the capacity of their sinks and reservoirs for the stabilization of such gases.

4.5.1.2 United Nations Convention on Biological Diversity

This convention, the main objectives of which is to preserve biological diversity and rehabilitate all degraded areas, was ratified by Sierra Leone on 12 December 1994. All signatory States are obliged to affect the prescribed undertakings which include:

- Development of national biological diversity strategy plan;
- Establishment of protected areas;
- Prevention, control and eradication of invasive and alien species;
- Provision of educational facilities.

4.5.1.3 Vienna Convention for the Protection of the Ozone Layer

The Vienna Convention, concluded in 1985, is a framework agreement in which States agree to cooperate in relevant research and scientific assessments of the ozone problem, to exchange information, and to adopt “appropriate measures” to prevent activities that harm the ozone layer. The obligations are general and contain no specific limits on chemicals that deplete the ozone layer. The ozone layer protects the earth against excessive ultraviolet radiation, which could cause damage and mutations in human, plant, and animal cells.

4.5.1.4 Montreal Protocol

The Montreal Protocol on Substances that Deplete the Ozone Layer (a protocol to the Vienna Convention for the Protection of the Ozone Layer) is an international treaty designed to protect the ozone layer by phasing out the production of a number of substances believed to be responsible for ozone depletion. The Treaty was opened for signature on 16 September 1987, and entered into force on 1 January 1989, followed by the first meeting in Helsinki in May 1989. Since then, it has been revised seven times, in 1990 (London), 1991 (Nairobi), 1992 (Copenhagen), 1993 (Bangkok), 1995 (Vienna), 1997 (Montreal), and 1999 (Beijing). It is believed that adherence to the international agreement will lead to the recovery of the ozone layer by 2050.

4.5.1.5 Rotterdam Convention

The Rotterdam Convention is a multilateral treaty to promote shared responsibilities in relation to the importation of hazardous chemicals. The Convention promotes the sharing of information and calls on exporters of hazardous chemicals to use proper labelling, include directions on safe handling, and inform purchasers of any known restrictions or bans. Parties can decide whether to allow or ban the importation of chemicals listed in the Convention, and exporting countries are obliged to ensure compliance by producers within their jurisdiction.

4.5.1.6 Convention on Wetlands of International Importance (RAMSAR)

The Ramsar Convention on Wetlands (Ramsar) was signed by Sierra Leone on December 13, 1999, and went into effect on April 13, 2000. Signatory countries to the Ramsar convention agree to:

- Include conservation of wetlands in land use planning throughout the country, including the promotion of “wise use” of wetlands;
- Establish nature reserves within wetland areas;
- Promote training in the fields of research, management, and gardening; and
- Consult with other signatory countries about implementation of the convention especially in areas of shared wetlands, shared water systems, and shared species.

As required by Ramsar, Sierra Leone identified and listed one wetland site for inclusion on the Ramsar wetland list. This non-contiguous wetland is located along the Sierra Leone River Estuary near Freetown. The three areas making up the wetland have a combined area of approximately 295,000 hectares (ha) and include mangrove swamps and upland coastal plains. The mangrove swamp included in this wetland makes up approximately 19% of all the mangrove swamp in Sierra Leone.

4.5.1.7 The Stockholm Convention on Persistent Organic Pollutants

The Stockholm Convention on Persistent Organic Pollutants is an international environmental treaty that aims to eliminate or restrict the production and use of persistent organic pollutants (POPs) - chemicals that are persistent bio-accumulates found in fatty tissues and are bio-magnified through the food chain, and adversely affect health and the environment. This Convention was adopted on the 22nd May 2001 in Stockholm and Sierra Leone became a signatory on the 27th August 2001. The convention came into force on 17 May 2004 with ratification by an initial 128 parties and 151 signatories. Co-signatories agree to outlaw nine of the dozen dirty chemicals, and curtail inadvertent production of dioxins and furans.

This Convention recommends the elimination or restriction of production and use of all internationally produced POPs (i.e. Industrial chemicals and pesticides), particularly, Aldrin, Chlordane, Dieldrin, Endrin, Heptachlor, Hexa-chlorobenzene (HCB), Mirixtexaphene, Polychlorinated Biphenyls (PCBs). The convention also seeks continuing minimization and, where feasible, ultimate elimination of the releases of POPs, such as Dioxins and Furans. Wastes containing POPs, must be managed and disposed of in a safe, efficient and environmentally friendly manner, with regards for international rules, standards and guidelines.

4.5.1.8 Convention on the International Trade of Endangered Species - (CITES)

The requirements of this convention became effective in Sierra Leone on the 16th January 1995. The convention seeks to eliminate and/or reduce trade in certain species inclusive of those that are considered endangered. By this convention, a list has been produced comprising of species that require protection against trade. The majority of the species listed in CITES are those also considered by the International Union for Conservation of Nature and Natural Resources (IUCN), as endangered and threatened. CITES also takes cognizance of species not necessarily threatened, but which require trade control to protect them from being threatened or endangered.

4.5.1.9 The UNESCO Convention for the Protection of the World Cultural and Natural Heritage (the World Heritage Convention), 1972

This convention is the foremost example of an international treaty developed to protect outstanding natural areas and resources. These sites, usually nominated for preservation by the state, are listed as part of "the world heritage". A 21-State elected committee of the treaty parties (the World Heritage Committee) decides which to list, and then States are obligated to protect their sites in perpetuity. Over 100 States are parties to the treaty, and 119 "natural" and "mixed" natural-cultural sites have been established as of 1996. Sierra Leone ratified this convention in 2005; however there are no sites currently listed in Sierra Leone.

4.5.2 International Lending Institutions Policies, Standards and Guidelines

Recognizing the political and resource differences among nations in their ability to set and implement environmental and social safeguards, the International Lending Institutions (ILI), development banks and private financial institutions, developed environmental and social safeguard policies and procedures. The purpose of these safeguards was to foster adoption and enforcement of the environmental and social assessment and protective measures required to implement the provisions of international agreements; to reinforce the intent of national legislation, and to foster sustainable development. Examples of these safeguards can be found in the following:

- World Bank Operational Policies and Bank Procedure
- International Finance Cooperation Performance Standards
- The Equator Principles

4.5.2.1 Roundtable on Sustainable Palm Oil (RSPO)

The RSPO, an association for the sustainable palm oil production, was formally established under Article 60 of the Swiss Civil Code on 8 April 2004 in Zurich, Switzerland.

The RSPO was established with the overall objective of promoting the growth and use of sustainable palm oil through cooperation within the supply chain and open dialogue with stakeholders. It has adopted and published a set of principles and criteria to help oil palm producers to be more sustainable. Within the overall framework of the document, practical advice is given to assist plantation managers in developing operational procedures towards identifying impacts and also to measure and monitor appropriate indicators that demonstrates a reduction of impacts over time.

The RSPO acknowledges that a key aspect of achieving sustainability is to identify the significant impacts especially those that are negative on the environment. The RSPO request producers to prepare:

- Documentation of the impacts and assessment of their relative importance;

- Development of strategic management plans which includes the results of such assessments;
- Development of operational procedures which identify impacts and the required changes in current practices to mitigate their negative effects; and
- Production of improvement plans, including a timetable for change

The RSPO undertakes the following principle tasks towards the fulfilment of its objectives:

- Research and develop definitions and criteria for sustainable production and use of palm oil
- Undertake practical projects designed to facilitate implementation of sustainable best practices
- Develop solutions to practical problems related to the adoption and verification of best practices for plantation establishment and management, procurement, trade and logistics
- Acquire financial resources from private and public funds to finance projects under the auspices of RSPO
- Communicate RSPO's work to all stakeholders and to the broader public.

5 BASELINE SURVEY AND CONDITION

The purpose of the present section is to establish an accurate baseline of the project area before the implementation of this project.

The baseline assessment was carried out on the physical, biological and social environments. Descriptions of the existing environment include primary and secondary data and information from relevant and available sources; text is illustrated with summary tables of data, maps, graphs, photographs and detailed written descriptions.

Noise, dust and wind speed measurements were taken in various locations as indicated in the following figure:




Table 5-4.5-1: Physical Site measurement points

5.1 Physical Environment

5.1.1 Climate

Sierra Leone's climate is characterized by two distinct seasons: the dry season (November – April) and the rainy season (May - October). The hottest months are March and April while the wettest months are July, August and September.

5.1.1.1 Methodology

Information and climatic data relating directly to the Freetown ports area was available from the Sierra Leone Meteorological Department. Field measurements, specifically wind speed, were used in conjunction with secondary data obtained, to represent climatic conditions within the project area and its surroundings. Climatic and other data for Bo area have been used to represent conditions within the project area.

The following climatic data represents the period 1976 - 2006 except for wind speed which was measured during the period of this study, August 2016.

5.1.1.2 Findings

The analysis could not be extended, with sufficient reliability because during the period of the baseline study, there was no climatic station in the area.

A summary of some climatic data for Bo was used and is presented in the following table:

Table 5.1-1: Summary of Some Climatic Data for Bo

Climatic Variable		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Monthly Mean Temp (°C) (1976-1980)	Max	31.9	33.6	33.9	33.1	31.8	30.3	28.5	27.9	29.3	30.7	31.0	30.
	Min	20.1	20.8	21.6	22.2	22.4	20.0	21.8	21.2	21.8	21.5	21.7	20.8
Average Relative Humidity (%) (1976 – 2005)	9 a.m.	78.7	79.1	78.1	78.9	81.4	86.0	89.4	91.1	88.7	86.3	83.6	79.8
	3 p.m.	62.4	63.4	64.0	65.9	70.5	75.5	81.0	82.8	78.8	76.0	74.1	67.9
Monthly Means of Rainfall (mm) (2007)		4.3	5.2	46.8	93.7	176.5	446.3	649.1	637	445.6	290.4	125.2	12.1

Rainfall

The project area lies within the Transitional Rain Forest Savanna Woodland Agro-climatic Region of Sierra Leone.

The project area lies in the south of the Transitional Rain Forest Savanna Woodland agro climatic region which is characterized by a high mean annual rainfall of 2500 -3000 mm and moderately low (290+/-30mm) water deficit spread over some 100 - 200 days (Kowal, et. al. 1980).

Generally, the climate of the project area is described as wet tropical monsoon with a single wet season each year. The average annual rainfall is about 2540 mm overall. The greater part of this rain falls between mid-April and mid-November and the wettest month is usually August, even though rivers attain maximum discharge in mid-September. About half of the annual precipitation (1460 mm) finds its way to groundwater or runoff resulting in stream and river flows (Dijkerman et al, 1964). The contribution of rainfall to stream flow is prolonged, lasting from the beginning of May to the end of November.

There is very little rain in December, January and February. River discharge is at its lowest in March and April, and begins to gradually increase in May. Groundwater levels do not rise significantly until late July.

The annual water budgets for this region (Bo) are shown in the following table.

Table 5.1-2: Annual Water Budgets (mm) for the Transitional Agro-Climatic Region and for BO

Component	Region	Bo
Precipitation	2737	2738
Evapotranspiration	1531	1372
Water Surplus	1671	1711
Water Deficit	465	345
Effective Precipitation	1166	1127
Growing Period Duration (days)	281	290

Source: UNDP/FAO- TR5, 1980)

Relative Humidity

Relative humidity in the morning (9.00 am) ranges between 87% and 94% while in the day (3.00 pm) ranging between 45% and 76%. The lowest in the morning hours were recorded in March and April at the peak of the dry season. During the day the lowest was also recorded in February and March in the peak of the dry season.

Temperature

Normal temperature range is 20.1°C to 33.9°C although it can drop below 20°C at night during the Harmattan season in January. Day temperatures average is 31°C in the dry season and 28°C in the wet season (Hall, 1966). Average short wave radiation of the season is 393 calories per cubic centimetre (cal/cm³). The duration of growing period is 271 - 298 days.

Evaporation

Evaporation is high during the dry season with highest in March/April, while it is relatively low in the rainy season with August/September recording the lowest. Evapotranspiration is very high in March been the hottest month and lowest in August which is peak of the rainy season.

Wind Speed

Wind speed measurements were recorded within and around twenty one (21) settlements in the STL study area. The measurements were recorded at different locations and times of the day using the portable anemometer vane probe.

Table 5.1.3 below shows the level of wind speed recorded during this period (10/08/16 to 13/08/16). The figures (0.03 m/s – 1.7m/s) indicate that wind speeds for this time of the year are almost nil in some settlement, while in other areas they are generally low to moderate depending on the time, elevation and open spaces.

Table 5.1-3: Wind speed measurements for settlements within and around the project area

Location	GPS coordinates	Date	Time	Average Wind speed m/s
Gelehun	0177169/0854025	10/8/2016	1.30 pm	0.03
Heima	0179944/0856603	10/8/2016	2.20 pm	0.8
Benduma Sewa	0178712/0855668	10/8/2016	5.00 pm	0.1
Kpatema	0182311/0853217	11/8/2016	10.40 am	0.3
Kpumbu	0180341/0853432	11/8/2016	2.00 pm	0.4
Benduma	0180962/0850991	11/8/2016	3.00 pm	1.4
Momandu	0830956/0842895	12/8/2016	10.55 pm	0.0
Komendi	0169318/0841307	12/8/2016	11.50 am	0.0
Baoma	0830575/0844190	12/8/2016	1.55 pm	0.4
Moforay	0171211/0845094	12/8/2016	2.52 pm	0.2

Location	GPS coordinates	Date	Time	Average Wind speed m/s
Mokombo	0172155/0842281	12/8/2016	3.55 pm	0.2
Gola	0172451/0843968	12/8/2016	4.45 pm	0.03
Mosogbo	0174984/0842056	13/8/2016	11.05 am	0.2
Tawamahehun	0174506/0843587	13/8/2016	12.10 pm	0.0
Garinga	0174256/0844568	13/8/2016	12.55 pm	0.0
Bamba	0176595/0844300	13/8/2016	1.55 pm	0.3
Sumbuya	0173447/0846516	13/8/2016	3.15 pm	1.7
Lower Saama	0175661/0847342	13/8/2016	4.05 pm	0.0
Yambama	0180728/0847114	13/8/2016	4.50 pm	0.0
Gendema	0179769/0843782	13/8/2016	5.20 pm	0.0
Tungie	0180326/0843466	13/8/2016	5.50 pm	0.0

Source: CEMMATS field measurements 2016

5.1.2 Air Quality and Dust

5.1.2.1 Methodology

The quantity of dust particles (PM10) in the air was recorded within and around twenty one (21) settlements in the study area. The measurements were recorded at different times and duration using a portable micro-dust pro aerosol monitoring system.

5.1.2.2 Findings

From the analysis of the data collected the dust levels at the time of the study were so low that the dust probe barely recorded levels of particulate matter. This was due to the period during which the measurements were taken (rainy season), with rainfall occurring during the time of data collection.

The WHO threshold for PM10 is 50µg/m³; the maximum values recorded range between 0.001mg/m³ (1µg/m³) and 0.011 mg/m³ (11 µg/m³) while the average values is 0.00 mg/m³ (0 µg/m³) in all the settlements. All values are within the WHO recommended standards indicating that the settlements have an environmentally friendly atmosphere during the rainy season which permits healthy living conditions. It is advisable to also carry out dry season measurements to determine dust levels during that period. Table 5.1.3 displays dust measurement results within the study area.

Table 5.1-4: Dust level within and around the project area

Location	GPS coordinates	Date	Duration	Average Values (mg/m ³)	Maximum Values (mg/m ³)
Gelehun	0177169/0854025	10/8/2016	27min 29 sec	0.000	0.000
Heima	0179944/0856603	10/8/2016	20min 38 sec	0.000	0.001
Benduma Sewa	0178712/0855668	10/8/2016	20min 33 sec	0.000	0.000
Kpatema	0182311/0853217	11/8/2016	30min 15 sec	0.000	0.000
Kpumbu	0180341/0853432	11/8/2016	30min 20 sec	0.000	0.001
Benduma	0180962/0850991	11/8/2016	30min 18 sec	0.000	0.000
Momandu	0830956/0842895	12/8/2016	30min 20 sec	0.000	0.001
Komendi	0169318/0841307	12/8/2016	30min 52 sec	0.000	0.000
Baoma	0830575/0844190	12/8/2016	21min 35 sec	0.000	0.000
Moforay	0171211/0845094	12/8/2016	21min 07 sec	0.000	0.001
Mokombo	0172155/0842281	12/8/2016	30min 25 sec	0.000	0.002
Gola	0172451/0843968	12/8/2016	30min 21sec	0.000	0.001
Mosogbo	0174984/0842056	13/8/2016	26 min	0.000	0.001
Tawamahehun	0174506/0843587	13/8/2016	31min 41sec	0.000	0.001
Garinga	0174256/0844568	13/8/2016	20min 16 sec	0.000	0.001
Bamba	0176595/0844300	13/8/2016	21min 12 sec	0.000	0.003
Sumbuya	0173447/0846516	13/8/2016	20min 39 sec	0.000	0.001
Lower Saama	0175661/0847342	13/8/2016	20min 11 sec	0.000	0.002
Yambama	0180728/0847114	13/8/2016	20min 38 sec	0.000	0.001
Gendema	0179769/0843782	13/8/2016	20min 16 sec	0.000	0.000
Tungie	0180326/0843466	13/8/2016	20min 04 sec	0.000	0.011

Source: CEMMATS Field measurements (August 2016)

5.1.3 Noise

5.1.3.1 Methodology

Noise levels were measured and recorded within and around twenty one (21) settlements in the project area. The measurements were recorded at different times and durations of the day using hand held Lutron sound meter.

5.1.3.2 Findings

The noise levels ranged from 38 decibels (dB) in Momandu to 71 decibels (dB) in Garinga. Levels were influenced by vehicular movements and ambient noise of normal community activities, normally referred to as community noise. Noise levels in some of the communities measured exceed 55db which is the average community noise level threshold. In terms of industrial environment (which the project area will soon become), noise levels are within the World Health Organisation (WHO) recommended limits, which stipulates that hearing protection gear must be used for noise exposure levels above 85 dB.

Table 5.1-5: Noise level data for settlements within and around project area

Location	GPS coordinates	Date	Time	Average noise level (dB) for 5 mins	Peak noise level range
Gelehun	0177169/0854025	10/8/2016	1:35 pm	41.5 – 59.30	
Heima	0179944/0856603	10/8/2016	2:27 pm	47.8 – 68.3	
Benduma Sewa	0178712/0855668	10/8/2016	5: 05 pm	38.8 – 58.5	
Kpatema	0182311/0853217	11/8/2016	10:45am	47.8 – 63.5	
Kpumbu	0180341/0853432	11/8/2016	12:20 pm	42 – 68.3	
Benduma	0180962/0850991	11/8/2016	3:05 pm	40.5 – 60	
Momandu	0830956/0842895	12/8/2016	11:00 am	38 – 56.8	
Komendi	0169318/0841307	12/8/2016	11:55 am	45.5 – 64.5	
Baoma	0830575/0844190	12/8/2016	2:00 pm	45.3 - 66	
Moforay	0171211/0845094	12/8/2016	2.57 pm	39 – 58.8	
Mokombo	0172155/0842281	12/8/2016	4.00 pm	40 – 60.8	
Gola	0172451/0843968	12/8/2016	4.50 pm	43.3 – 62.3	

Location	GPS coordinates	Date	Time	Average noise level (dB) for 5 mins	Peak noise level range
Mosogbo	0174984/0842056	13/8/2016	11.11 am	46.5 – 65	
Tawamahehun	0174506/0843587	13/8/2016	12.15 pm	42.5 - 67	
Garinga	0174256/0844568	13/8/2016	1.00 pm	48.5 - 71	
Bamba	0176595/0844300	13/8/2016	2.00 pm	44.3 – 65.3	
Sumbuya	0173447/0846516	13/8/2016	3.20 pm	57.3 – 70.1	
Lower Saama	0175661/0847342	13/8/2016	4.10 pm	51.3 – 63.5	
Yambama	0180728/0847114	13/8/2016	4.55pm	44.5 – 59.8	
Gendema	0179769/0843782	13/8/2016	5.30pm	44.8 - 56	
Tungie	0180326/0843466	13/8/2016	5.55 pm	43 – 59.8	

Source: CEMMATS Field measurements (August 2016)

5.1.4 Geology

The project area geological study was done mostly through desk studies on information gathered applicable to the site and its general surroundings.

The Project area geology is within the Basement Granite and Acid Gneiss Terrain of the geology of Sierra Leone as it stretches along the Sewa River alluvial diamond field and covers approximately 7.5 to 17 km of reach to the diamondiferous Sewa River. The floor-rocks to the wide-spread gravel occurrences comprise the Leonean Granite and Gneiss Terrain of the Man Craton of West Africa.

The project areas along the Sewa River hosts both modern and palaeo diamondiferous gravel deposits. A variety of alluvial facies, including high terrace, middle terrace, low terrace, swamp and river depositional environments types spread sporadically throughout the project area. As a result of the alluvial deposits, majority of the artisanal mining activities are confined to the lower terrace facies, swamp facies and the modern river system.

The area is composed of prospective gravel horizons made up of lower terrace, middle terrace, upper terrace and swamp facies. The middle and upper terraces are terrestrial, chemically weathered and relatively in-situ, regolith landforms that form a lateritic residuum comprising a ferruginous (haematite and goethite) gravel horizon, containing abundant pisoliths and nodules, as well as diamonds. Unlike the lower terrace gravels, a relatively thin overburden allows easy access to these gravels, with a considerable thickness.

The alluvial deposits comprise a range of environments of deposition, from down-wasted, residual regolith landforms, to fringing terraces of the Sewa River and elongate inland swamps systems, which comprise a large percentage of the total drainage network; in fact, the gravel horizons of the swamps have, in the past, constituted the second-most important source of alluvial diamonds

5.1.5 Landform

5.1.5.1 Methodology

The landform assessment study was done mainly through desktop review of secondary data and field observations.

5.1.5.2 Results

The project area is generally located on a gently undulating plain with isolated hill remnants and dissected by well-defined valley swamps. The hills are usually rising with an extremely low relief from fairly broad interfluves. The gently to moderately sloping interfluves side slopes have commonly dissected by broad gullies giving rise to narrow, uneven crests. There is usually a well-defined break of slope between the interfluves and valley swamps, except where gentle, narrow foot slopes and terraces occur. Adjacent to the Sewa River there is continuous alluvial plains of variable width. There are four main landforms occurring within the project area which is described below:

- **Isolated hills** - short, straight to convex, gentle to moderately sloping.
- **Interfluves** - medium length undulating, very gentle to gentle sloping.
- **Valley swamps** - level to nearly level with varying width and locally channelled
- **Floodplains** - level to almost flat with varying width and locally channelled and terraced.

5.1.6 Soils

5.1.6.1 Methodology

The soil baseline study was done by a random survey method. Soil descriptions were done using auger borings on the different landforms within and around the study area. At each observation point, the soil was augered up to at least 100cm where possible and described in detail using FAO Guidelines for soil profile descriptions (FAO 1990). The parameters described are shown on the soil/landform physical properties table.

The results of soil tests conducted by STL in 2014, in which 20 composite samples collected within Moyamba and Bo were sent to the Sierra Leone Agricultural Institute (SLARI) Laboratory for testing, are included in the analyses.

5.1.6.2 Results

Generally the soils can be classified as uplands and bottomlands. The upland soils are deep to shallow, gravelly and gravelfree and well drained while the bottomland is deep, generally gravelfree, poorly drained to waterlogged.

From observations the soils can be divided into four types according to textural differences. Texture is an important physical characteristic that plays an important role in the fertility and degradation of the land, provisions for the minimisation and control of soil erosion and degradation can be implemented by proper management strategies and operational techniques which will minimise erosion. Related impacts of the project on soils in the project area relate mainly to the land clearing phase which will result in loose exposed soil susceptible to erosion. These impacts and their mitigation measures are detailed in the impact table.

Physical Properties

The physical properties of the various soils are described below:

a) Gravelly soils

These soils occur on the interfluvies of the upland and are generally shallow with compacted ironstone gravels occurring at an average depth of 50cm.

These soils comprise of the following properties:

- Moderately deep to shallow;
- Texture vary from gravelly Sandy loam to gravelly sandy clay loam in the topsoil and very gravelly sandy clay in the subsoil ;
- Well drained with water table below 50 cm ; and
- Black to dark brown over brown to dark yellowish brown colour

b) Gravel free over gravel soils

These soils generally occur on the interfluvies of the uplands and are generally gravel free in the topsoil and very gravelly in subsoil.

These soils comprise of the following properties:

- These soils are generally shallow to moderately deep;
- Texture ranges between sandy loam and sandy clay loam in the topsoil and gravelly sandy clay subsoil;
- They are well drained with water table below 80 cm; and
- Colours are black to dark brown top soil and dark yellowish brown in the subsoil

c) Gravel free soils (interfluvies/terrace)

The soils generally are gravel free throughout the profile and occur on the floodplains/lower terraces

These soils comprise of the following properties:

- Generally deep;

- Textures are generally ranging from sandy loam to sandy clay loam overlying sandy clay to clayey subsoil;
- Well drained with water table below 100 cm; and
- Colours are very dark brown to dark yellowish brown overlying light yellowish brown to brownish yellow with prominent yellowish mottles.

d) Gravel free Soils of the valley swamp (uncultivated)

Soil observation was done on the uncultivated swamp whose properties comprise:

- Deep to moderately deep at the fringes;
- Texture is silty loam to silty clay loam in the topsoil overlying sandy clay to clayey sand with sand content and grain size increasing with depth;
- Poorly drained to waterlogged; and
- Colours are very dark greyish brown over grey

e) Gravel free Soils of the valley swamp (cultivation)

Soil observation was done on the cultivated swamp whose properties comprise:

- Deep to moderately deep at the fringes;
- Texture is silty clay loam in the topsoil overlying sandy clay to clay with few ironstone gravel in the subsoil;
- Poorly drained to waterlogged; and

Colours are very dark brown over yellowish brown to brownish yellow subsoil

Table 5.1-6: Typical representatives of Soil/Landform Physical Properties in Sierra Tropical Limited Project Area

Obs No	Physiography	Soil types/Land form	Mapping Units	Soil Depth	Texture	Consistence	Colour		Coarse Fragments		H ₂ O Table	Remarks
							Matrix	Mottles	Type	%		
ST - 1	Upland	Gravel soils/ Interfluve	A	0-20	GrSL- GrSCL	SS/NP	10 YR 2/1	-	Ironstone gravel	20 -50	Below 50cm	Gravel content increases with depth. The entire profile is gravelly. Auger blocked at 50cm due to compacted gravel. Soil moisture – 46%. Field pH - 6.5
				20-50	VGrSC	SS/SP	10YR 3/4	-	Ironstone gravel	>50		
ST - 2	Upland	Gravel free over Gravel soils/Interfluve	B	0-20	SL - SCL	SS/NP	10YR 2/2	-	-	-	Below 80cm	Auger blocked at 80cm due to compacted gravel. Gravel and clay content increases with depth. Soil moisture – 50%. Field P ^H - 6.2
				20-50	SC	S/SP	10YR 4/4	-	-	-		
				50-80	GrSC	S/SP	7.5YR 4/6	-	Ironstone gravel	> 50		
ST - 3	Upland	Gravel free soils/Interfluve/ Terrace	C	0-15	SL - SCL	NS/NP	10YR 2/2	-	-	-	Below 100cm	Gravel free profile Clay content increases with depth. Soil moisture – 50%. Field P ^H -6.5
				15- 40	SC	SS/SP	10YR 6/4	10YR 7/8	-	-		
				40 – 100	SC - Clay	S/P	10YR 6/6	-	-	-		

Obs No	Physiography	Soil types/Land form	Mapping Units	Soil Depth	Texture	Consistence	Colour		Coarse Fragments		H ₂ O Table	Remarks
							Matrix	Mottles	Type	%		
ST - 4	Lowland	Gravel free soils/Inland valley swamp (uncultivated)	D	0 – 50	SiL- Siel	SS/NP	2.5Y 3/2	--	-	-	Waterlogged	Sand content increases with depth. Soil moisture – 80%. Field P ^H – 5.6
				50 - 100	SC - CS	SS/SP	2.5Y 5/1	-	-	-		
ST - 5	Lowland	Gravel free soils/Inland valley swamp (cultivated)	D1	0 – 20	SiCL	SS/SP	10YR 2/2	-	-	-	Below 10cm	Very little ironstone gravel in the sub soil. Clay content increases with depth. Soil moisture – 59%. Field P ^H - 6.9
				20 – 40	SC - C	S/P	10YR 5/6	-	-	-		
				40 - 100	SC - C	S/P	10YR 6/8	-	-	-		

Note: CS – Clayey Sand, SL – Sandy loam, SC – Sandy Clay, VGr – Very gravelly, SCL –Sandy Clay Loam, Si - silt

Chemical Properties

Soil tests were carried out in 2014 on samples from Moyamba and Bo, using methods described jointly by the International Soil Reference and Information Center (ISRIC) and the FAO (ISRIC/FAO, 2002, Procedures for Soil Analysis. Technical Paper 9, 6th Edition).

Soil color was visually compared with the Munsell Chart. Soil pH was determined on 1:1 soil:water and 1:1 KCl extracts. Exchangeable cations (Na, K, Ca, and Mg) were measured on neutral 1N ammonium acetate extracts. Exchangeable K and Na were read on a Flame Photometer while exchangeable Ca and Mg were read on an Atomic Absorption Spectrophotometer (AAS). Exchangeable Acidity (Al + H) was extracted by 1M KCl and titrated with 0.025 M NaOH. Effective CEC was calculated as the sum of exchangeable cations and exchangeable acidity. Zn, Cu, Fe and Mn were measured by AAS on DTPA (Diethylenetriaminepentaacetic acid) extracts.

The results of these tests are highlighted in the following tables:

Soil Test	Results									
	Sample 1	Sample 2	Sample 3	Sample 4	Sample 5	Sample 6	Sample 7	Sample 8	Sample 9	Sample 10
Soil Color (Munsell)	10YR 5/3 Yellowish brown	7.5YR 6/2 Greyish brown		10YR 5/3 Dull yellowish brown			10YR 5/2 Greyish yellow brown			
pH (1:1 H ₂ O)	4.53	5.00	4.90	4.33	4.68	5.04	5.17	5.01	4.82	5.27
pH (KCl)	4.09	4.21	4.15	4.13	4.23	3.82	4.32	4.27	3.98	4.03
% Organic Carbon	1.6	1.1	2.7	1.4	1.3	1.7	1.4	1.3	0.6	2.5
Total Nitrogen (%)	0.12	0.09	0.15	0.11	0.09	0.12	0.14	0.1	0.07	0.15
P (Bray 1) mg/kg	3.32	2.36	1.96	5.09	2.49	2.46	2.58	3.17	3.20	1.74
Exch K cmol(+)/kg	0.65	0.85	0.11	0.85	0.82	0.82	0.90	0.90	0.18	0.25
Exch Ca cmol(+)/kg	1.10	1.18	1.89	1.50	1.18	1.18	1.25	1.89	1.95	2.10
Exch Mg cmol(+)/kg	0.35	1.40	0.47	0.45	1.25	1.25	1.40	1.55	1.05	1.54
Exch Na cmol(+)/kg	0.05	0.45	0.11	0.05	0.25	0.25	0.25	0.25	0.05	0.25
Exch. Acid (H+Al) cmol(+)/kg	5.49	4.65	4.79	4.77	4.62	4.91	4.62	4.90	4.83	4.89
Effective CEC cmol(+)/kg**	7.64	8.53	7.37	7.62	8.12	8.41	8.42	9.49	8.06	9.03
Zn (DTPA) ppm	1.75	1.35	1.75	1.50	2.05	2.05	1.75	1.35	1.75	1.35
Cu (DTPA) ppm	2.70	0.95	1.35	1.35	0.35	0.35	1.30	0.90	1.45	0.95
Fe (DTPA) ppm	6.05	5.54	6.00	4.55	6.05	4.55	7.17	6.70	4.00	4.55
Mn (DTPA) ppm	3.5	5.0	5.0	4.0	5.0	2.5	2.0	2.5	2.5	2.0

Soil Test	Results									
	Sample 11	Sample 12	Sample 13	Sample 14	Sample 15	Sample 16	Sample 17	Sample 18	Sample 19	Sample 20
Soil Color (Munsell)	10YR 5/3 Dull yellowish brown					10YR 5/4 Dull yellow orange				
pH (1:1 H ₂ O)	5.83	4.92	4.84	4.76	4.75	5.09	4.55	5.40	5.16	5.38
pH (KCl)	4.89	3.99	4.14	4.12	3.96	4.27	3.96	4.26	4.19	4.25
% Organic Carbon	2.6	3.0	1.9	2.9	1.6	1.9	1.5	1.6	0.8	2.4
Total Nitrogen (%)	0.13	0.15	0.11	0.15	0.09	0.10	0.11	0.16	0.07	0.15
P (Bray 1) mg/kg	3.17	5.77	1.99	1.37	1.53	3.35	3.91	1.16	2.18	8.35
Exch K cmol(+)/kg	0.21	0.18	0.35	0.15	0.95	0.83	0.90	0.18	0.95	0.15
Exch Ca cmol(+)/kg	1.28	2.45	2.05	1.89	2.15	1.95	2.10	2.05	2.35	1.90
Exch Mg cmol(+)/kg	0.58	1.15	1.50	1.45	1.45	1.25	1.35	1.45	1.45	1.35
Exch Na cmol(+)/kg	0.05	0.10	0.15	0.05	0.15	0.25	0.20	0.15	0.20	0.15
Exch. Acid (H+Al) cmol+/kg	5.49	4.65	4.79	4.77	4.62	4.91	4.62	4.90	4.83	3.4
Effective CEC cmol+/kg**	7.61	8.53	8.84	8.31	9.32	9.19	9.17	8.73	9.78	6.95
Zn (DTPA) ppm	2.63	2.25	3.08	3.08	2.63	3.68	2.63	3.00	3.08	3.08
Cu (DTPA) ppm	1.78	0.40	0.46	0.46	0.40	0.46	1.78	0.40	0.46	0.46
Fe (DTPA) ppm	5.45	5.30	5.46	5.90	5.30	4.96	4.96	5.30	5.96	5.90
Mn (DTPA) ppm	2.5	2.5	2	2.5	2.5	2	2.5	2.5	2.5	2.5

Summary of Analyses Results

	Minimum	Maximum	Mean	Median	Coefficient of Variability (%)
pH (1:1 H ₂ O)	4.33	5.83	4.97	4.96	6.80
pH (KCl)	3.82	4.89	4.16	4.15	5.07
% Organic Carbon	0.60	3.00	1.79	1.60	37.35
Total Nitrogen (%)	0.07	0.16	0.12	0.12	23.45
P (Bray 1) mg/kg	1.16	8.35	3.06	2.54	54.46
Exch K cmol(+)/kg	0.11	0.95	0.56	0.74	60.27
Exch Ca cmol(+)/kg	1.10	2.45	1.77	1.90	23.71
Exch Mg cmol(+)/kg	0.35	1.55	1.18	1.35	32.33
Exch Na cmol(+)/kg	0.05	0.45	0.17	0.15	58.75
Exch. Acid (H+Al) cmol+/kg	3.40	5.49	4.77	4.79	8.29
Effective CEC cmol+/kg**	6.95	9.78	8.46	8.48	8.77
Zn (DTPA) ppm	1.35	3.68	2.29	2.15	30.63
Cu (DTPA) ppm	0.35	2.70	0.94	0.68	67.43
Fe (DTPA) ppm	4.00	7.17	5.48	5.46	13.70
Mn (DTPA) ppm	2.00	5.00	2.90	2.50	34.31

•

5.1.6.3 Soil Survey, Water and Climate Assessment of Sierra Leone

An exploratory Soil Survey, Water and Climate Assessment of Sierra Leone was conducted by Dr. Carlos Gauggel, Mr. Carlos Mandujano and Mr. Romeo Patricio in 2011. The assessment resulted in the following conclusions:

9. There are about 2.8 million ha of suitable soils for pineapple production in the Lower and Inner Coastal Plains of Sierra Leone out of which 1.8 million can be irrigated using the major rivers and very close to roads and townships.
10. In general, in this study, soil quality for pineapple production, as the first priority, has been ranked A, B, B2 and C in this report. Availability for water has been considered for this ranking. The available number of ha per zone is as follows : Area A 630000 ha, Area B 720000 ha, Area B2 360000 ha, Area C 180000 ha
11. Irrigation has to be implemented from December to May, river water and water reservoirs are the main source of water for irrigation. The number of ha indicated above have access to irrigation.
12. Primary roads and bridges are in very good shape and can handle containers. Second grade roads are very varied in condition, the condition of unpaved roads is a major issue in the rainy season.
13. Surface drainage and sub-surface drainage, in some areas, are a must for successful pineapple production due to heavy rains during the rainy season.
14. All soils will require an initial cross-way deep sub soiling to loosen the soil matrix and to break up any laterite or iron-stone layers, followed by disk harrowing, this is a must for the success of the project and it is probably required once (deep sub soiling).
15. Due to water availability, sloping landscapes, and the high risk of soil erosion, drip or low discharge irrigation is the best option.
16. The low temperatures (December-March), limited sun-light hours during some months (July-August) constitute the major constraint for pineapple production in Sierra Leone. Soil and water-wise it is very good area for pineapple cultivation with vast land and water resources available.

5.1.7 Hydrology

5.1.7.1 Methodology

The hydrology study was conducted through literature review of relevant reports on the project area, collection of samples and laboratory analyses, conduct of a hydro census and general observational assessments of the project area.

5.1.7.2 Results and Discussion

5.1.7.2.1 Description of Climatic Condition

Sierra Leone falls within a tropical climate zone with peculiar wet and dry seasons splitting the year in equal halves:

- Rainy Season (May –November): Thunderstorms and squalls and Steady rains;
- Dry season (December – April): Dry weather with high humidity and Dry weather with low humidity.

These seasons could be subdivided into four (4) distinct types of weather. (i) Heavy squalls and thunderstorms, (ii) steady rains (iii) Harmattan season characterised by dry and windy weather with low humidity and (iv) Hot and humid weather.

Thunder Storms and Squalls

Heavy squalls and thunderstorms characterize two different times of the year: May and June, which marks the beginning of the wet season and October and November, the end of the rainy season. Thunder storms accompanied by heavy rains travel east to west against the general wind direction. They are usually preceded by squall of the easterly wind. These thunderstorms are responsible for most of the rains at those times of the year. Towards the beginning and the end of the rainy season, the thunderstorms decrease in frequency and intensity and the weather is very changeable. The relative humidity throughout the rainy season is very high.

Steady Rains

Steady rains occur from July through September which is about the middle of the rainy season. Rainfall is frequent and often heavy and generally, most of the annual rainfall occurs during this period, being brought in by the south-west Monsoon winds. The sky is mainly overcast, sunshine is rare and the relative humidity ranges from 95% to 100%. The temperatures are at their lowest during this period and the diurnal range of temperature is small.

Dry Weather with High Humidity

Dry Weather with High Humidity occurs mostly during the dry season. Skies are usually clear and therefore day temperatures are relatively high. The nights are also warm and very humid. Heavy dew and fog often occur during the night and early morning. Winds predominantly come from the west.

Dry Weather with Low Humidity (Harmattan)

Dry Weather with Low Humidity (Harmattan) usually occurs between late December and early February. The lengths of these periods vary from a few days to a number of weeks. The

weather is characterized by a sudden drop in relative humidity from almost 100% to sometimes as low as 20%. Evapotranspiration is high because of low relative humidity and high temperature.

Climatic Evaluation

Daily rainfall and temperature records from the station at Bo from 2003 to 2009 were analyzed. There are no records for Sumbuya, but because Bo Town is the closest station to the project area, data from this station were used for analysis (Reference Table 5.1-1 on page 83).

5.1.7.2.2 Sensitive Human and Ecological Receptors

With regards to hydrology, a range of human and environmental receptors are located within the Sierra Tropical Agriculture (STA) concession area and immediate vicinity as listed in Table 5.1-7. These receptors were defined during the baseline studies undertaken for this ESIA, and are used as part of an integrated approach in the analysis of water resources.

Table 5.1-7 summarizes these receptors in terms of the segments for the Project and its operations.

Table 5.1-7: Sensitive Receptors

Receptors	Project Concession Area and vicinity	Access/Haul Road	Water Resources Consideration
HUMAN-- Communities and settlements	Settlements in and around the concession and dependent on the Sewa River on the West flank of the concession area and tributaries that flow in the North-east to south-west direction into the Sewa. These settlements could be affected by changing quality of water and flow patterns in the surface water, especially in the Dry Season.	Interception of water flow and contamination of water from streams and wetlands over or close to which access/haul roads are constructed could affect the activity for which that water is used.	Low flows for water supplies; High flows for water/flood management; Water quality for use of agrochemicals and pesticides; Spills
Ecological	Sewa River, its tributaries and Wetland habitats		Water quality; contaminated sediment; spills

5.1.7.2.3 Water Users

Freshwater systems located within the project site and its surrounding areas are used extensively by local inhabitants for potable water needs, irrigation/farming, washing, livestock watering, fishing and navigation within the Lugbu Chiefdom or between the Bo and Bonthe districts.

5.1.7.2.4 Estimated Project Water Requirements

Estimation of Evapotranspiration in the Sumbuya Area

In the Sumbuya area, evaporation pan data is not available. It was therefore necessary to estimate the Evapotranspiration (Etp) for the pineapple crop, using the methods of Thornthwaite and Blanney-Criddle (See Tables 1 and 2, respectively). Additionally, a crop factor (Kc) of 0.75 (for this crop the value varies from 0.25 to 0.75 during the growth cycle), soil (Oxisols) data and a root zone depth of 20 cm, was considered for this calculation.

The highest values were obtained using the Blanney-Criddle method, during the period of March to August and reaching a total of 1,510.9 mm/year. In the case of the calculations by the Thornthwaite method the Etp is lower and the highest values occurred during the Months of March to July, with a total value of 1,273.2 mm/year.

Table 5.1-8: Calculation of Monthly Evapotranspiration Using Thornthwaite Method for Sierra Leone

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Mean T (°C)	26.0	27.0	27.5	28.0	27.5	26.5	25.5	24.5	25.0	25.5	26.5	26.0
I	12.1	12.8	13.2	13.6	13.2	12.5	11.8	11.1	11.4	11.8	12.5	12.1
E	131.7	141.2	145.6	149.8	145.6	136.5	126.7	115.9	121.4	126.7	136.5	131.7
Nd	31	28	31	30	31	30	31	31	30	31	30	31
Ni	1.03	0.93	1.03	1.00	1.03	1.00	1.03	1.03	1.00	1.03	1.00	1.03
HRS	12.1	12.2	12.4	12.7	12.9	13.0	12.9	12.8	12.5	12.3	12.1	12.0
Li	1.01	1.02	1.03	1.06	1.08	1.08	1.08	1.07	1.04	1.03	1.01	1.00
ETTho (mm)	137.2	133.9	155.4	158.5	161.7	147.9	140.7	127.8	126.5	134.2	137.7	136.1
ETTPine apple	102.9	100.5	116.6	118.9	121.3	110.9	105.5	95.8	94.8	100.6	103.3	102.1

Source of Temperature, Length day and Rainfall Data: (Sierra Leone Weatherbase).

Table 5.1-9: Calculation of the Monthly Evt using the Blaney & Criddle Method for Sierra Leone

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Mean T (°C)	26.0	27.0	27.5	28.0	27.5	26.5	25.5	24.5	25.0	25.5	26.5	26.0
Nd	31	28	31	30	31	30	31	31	30	31	30	31
Ni	12.1	12.2	12.4	12.7	12.9	13.0	12.9	12.8	12.5	12.3	12.1	12.0
Ii	375.1	341.6	384.4	381.0	399.9	390.0	399.9	396.8	375.0	381.3	363.0	372.0
Ii/I	8.23	7.49	8.43	8.36	8.77	8.55	8.77	8.70	8.22	8.36	7.96	8.16
ETBla&Criddle (mm)	164.6	153.4	174.5	174.9	181.5	173.1	173.5	168.2	160.8	165.5	161.2	163.3
ETPPineapple	123.5	115.0	130.9	131.2	136.2	129.9	130.1	126.1	120.6	124.1	120.9	122.5

Source of Temperature, Length day and Rainfall Data: (Sierra Leone Weatherbase).

Water Balance

The water balance is a technique, developed to evaluate the soil moisture, and allow the detection of deficits or excess water in the soil, which when it is quantified can be used for various purposes. By understanding the specific crop potential evapotranspiration (ETP) of a period, against the average rain or probable rain (R), it is possible to determine any deficit (irrigation needs) or water excess (drainage needs). However, since the soil can store water, its water availability capacity must be considered in the water balance (AWC).

With known water inputs (usually rain), the potential evapotranspiration and initial soil moisture content (ISM), it is possible to determine in a specific period (month, week, day, etc.) a balance that simply quantifies the water deficit or water excess.

If $(ISM + R) - ETP > AWC = \text{Excess}$; If $(ISM + R) - ETP < AWC = \text{Deficit}$

The depth of the roots and the water depth available for the crop will modify the water balance in terms of the excesses and deficits. Table 3 (Thornthwaite Method) presents a general water balance of Sierra Leone, which indicates that during the month from May to November, there potentially exists an accumulated excess of 1744.4 mm. Between December to April a deficit of 450.3 mm occurred and from May to November a soil water storage accumulation occurs (210 mm), which is depleted due to the evapotranspiration. In Table 4 (Blaney & Criddle Method) this shows a general water balance, indicating that during the months from May to November, there is an accumulated excess of 1588.7 mm. Between December and April a deficit of 532.4 mm occurred. From May to November, an excess or soil water storage accumulation occurs (210 mm).

Table 5.1-10: Monthly Water Balance for Sierra Leone using average data and ETP by Thornthwaite method

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Rainfall (mm)	7.8	19.0	48.0	108.1	233.4	338.2	475.6	541.9	447.6	326.4	143.5	27.8
Mean T (°C)	26.0	27.0	27.5	28.0	27.5	26.5	25.5	24.5	25.0	25.5	26.5	26.0
ETTho (mm)	137.2	133.9	155.4	158.5	161.7	147.9	140.7	127.8	126.5	134.2	137.7	136.1
ETPPineapple	102.9	100.5	116.6	118.9	121.3	110.9	105.5	95.8	94.8	100.6	103.3	102.1
SWAC (mm) ¹	0.0	0.0	0.0	0.0	30.0	30.0	30.0	30.0	30.0	30.0	30.0	0.0
Deficit (mm)	125.1	111.5	98.6	40.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	74.3
Excess (mm)	0.0	0.0	0.0	0.0	82.1	227.3	370.1	446.1	352.8	225.8	40.2	0.0

¹ Soil Water Available Capacity for an average Oxisol with up to 20 cm root zone depth.

Conclusion

1. The results of the monthly water balance, calculated by the methods of Thornthwaite and Blaney-Criddle, indicate that during the months of December to May an accumulated water deficit of 450.3 and 532.4 mm occurs, respectively. The only way to satisfy this deficit it is by the use of an irrigation system. (A drip or sprinkler systems are the likely considered options).
2. The highest monthly irrigation requirement corresponds to January with values of 125.1 and 145.7 mm/month, meaning that daily watering needs are 4 and 4.7 mm/day, according Thornthwaite and Blaney-Criddle methods, respectively.
3. In order to accurately characterize the soils of Sierra Leone for irrigation potential, it is necessary to acquire laboratory and field equipment to determine physical characteristics like bulk density, saturation point, field capacity, wilting permanent point, soil-moisture curves, etc. The services of SLARI at Njala University will be acquired to complete the proper and required soil physical/moisture curves and characteristics.
4. To program and monitor irrigation requirements in Sumbuya, it will be necessary to install 5-10 weather stations (Watch Dog 2000) five at very least , A type evaporation pans (according to specifications of the USDA) to determine the daily irrigation requirements for the pineapple crop. Also, additional equipment, to measure the soil-

moisture will be necessary to implement a monitoring system of irrigation and soil moisture or the Daily Water Balance.

Summary of Total Water Requirements for a Season

From the above assumptions and calculations 5mm/day at the driest time of the year, is the minimum quantity of water required. To allow a safety margin, as we are only generally estimating climate and soil data, calculations for this exercise are based on 7mm/day.

Note: after more detailed soil analysis, by SLARI, the adoption of conservation tillage and agriculture principles, as well as irrigation type, greater soil water conservation can be achieved and in practice, total irrigation water requirements maybe much lower.

Annual estimated crop irrigation water requirements:

Water required	7mm/day
Continuous water flow required in litres/second/ha	0.81lt/sec./ha
Total equivalent ha	4335ha
Total irrigation water requirements (maximum)	7,585, 000 m ³ per year

5.1.7.2.5 Regional Context

Sierra Leone has a complex drainage pattern that includes numerous rivers and smaller creeks and streams. The larger rivers originate in the Fouta Djallon highlands of Guinea and generally flow from northeast to southwest across Sierra Leone. The country's drainage system includes nine major rivers and minor coastal creeks and tidal streams. The river watersheds range in size from 14,140 km² for the Sewa River to less than 385 km² for the smaller watersheds.

5.1.7.2.6 Project Site

During the field work conducted in July, 2016, within the project concession area, plans to conduct stream velocities and stage (water elevation) measurements were thwarted by the high discharge of the Sewa River. The team considered it unsafe to carry out such measurements.

5.1.7.2.7 Water Quality

The following receptors have been identified:

- Residents who use surface water for drinking water purposes (largely outside of the mining concession area);
- Residents who are exposed to surface water through fishing, rice farming, or swimming. Although these may not be considered purely recreational activities, international recreational guideline values would apply; and
- Ecological receptors within the sediment and surface water.

In order to assess potential risks to these receptors the following screening values were used:

- World Health Organisation (WHO), 2011 Guidelines for drinking-water quality;
- IFC, 2007 Environmental, Health and Safety Guidelines for Mining.

The following table summarizes the numerical standards used to compare the baseline data.

Table 5.1-11: Numerical Standards

Parameter	Units	WHO Drinking Water Guidelines (2011)	IFC effluent guidelines	Sierra Leone Environment Protection Regulations 2013. Maximum at any moment (Annual average maxima) (unfiltered concentrations)
Field Temp	°C		<3°C differential	
Field pH	pH units	6.5-8.5	6 – 9	6 – 9
Field EC	uS/cm			
Field Turbidity	NTU			
Field DO	mg/l			
Field TDS	mg/l	600		
Total Alkalinity as CaCO ₃	mg/l			
Total Hardness Dissolved	mg/l	200		
Sulphate	mg/l	250		
TSS	mg/l		50	50 (25)
COD (settled)	mg/l		150	
Trace elements		Metals are total	Metals are total	
Aluminum	ug/l	100*		
Arsenic	ug/l	10	100	100 (80)
Barium	ug/l	700		
Cadmium	ug/l	3	50	50 (40)
Total Chromium	ug/l	50	100	100 (80)
Copper	ug/l	2000	300	600 (400)
Total Iron	ug/l	300*	2000	2000 (1600)
Lead	ug/l	10	200	200 (160)
Magnesium	mg/l			
Manganese	ug/l	100*		
Mercury	ug/l	6	2	2 (1.6)
Nickel	ug/l	70	500	

Parameter	Units	WHO Drinking Water Guidelines (2011)	IFC effluent guidelines	Sierra Leone Environment Protection Regulations 2013. Maximum at any moment (Annual average maxima) (unfiltered concentrations)
Zinc	ug/l	3000*	500	1500 (1200)

The WHO guidelines were used in this study to assess drinking water quality and the lower values of either the IFC EHS guidelines or Environment Protection Regulations were used to assess effluent quality.

It should be noted that the Sierra Leone Environment Protection Regulations and the WHO standards are the only values to which water/effluent discharges from the study area need comply

5.1.7.2.8 Project Area and Surroundings

This work includes the measurement of inorganic compounds and other physical and chemical parameters (i.e. metals, temperature, pH, dissolved oxygen (DO), turbidity, conductivity, and total dissolved solids (TDS)). **Error! Reference source not found.** provides a brief description of the key sites sampled in August, 2016.




Figure 5.1-15.1-2: Map of Water Quality Sampling Sites at or in the vicinity of the Project site

5.1.7.2.9 Analysis and Summary

Specifically, seasonal and spatial variation in water quality in the catchments can be driven by:

- Dry season stagnation, resulting in a reduction of oxygen levels.
- Elevation of TSS and turbidity as a result of rainfall runoff, which typically has high suspended sediment loads from disturbed areas, combined with lower dilution during the dry season.
- Low pH in surface waters related to low natural buffering capacity, and reactions associated with biological activity in the wetland areas. pH varies seasonally, with a lower average pH in the wet season.

Elevated levels of iron, manganese and other metals are natural to the geology though weathering and erosion of exposed areas may be a factor within the project's concession area.

5.1.7.2.10 Domestic Water Quality and Environmental Sanitation

Domestic Water Quality

Introduction

This study was aimed at collecting data on all water points and channels within the project site as well as the surrounding settlements that may be affected by the company's operations and the analysis of potential impacts of the proposed activity on the water resources.

Water is a very important resource that needs to be maintained and preserved to ensure its potability. Water is vitally important to every aspect of our lives. Monitoring the quality of surface water will help protect waterways from pollution by project activities. It is therefore essential for development practitioners to embark on activities that will protect water resources as well as water channels within local communities that solely depend on these sources for their survival.

Methodology

Field investigations were conducted within the study area and its immediate environs to obtain all relevant data. This involved identifying all water points and channels within the study area. The methodology used for the study includes the following:

- An initial reconnaissance visit was made to the study area by a field party;
- The field party travelled to the study area to identify sites for sampling and measurements and to acquaint residents and project officials with the purpose of the study;
- Questionnaires were administered to residents in the villages within the study area by CEMMATS Social team to obtain data on several issues regarding water supply and quality and sanitation issues;

- Samples were collected from the identified water sources. Some physical parameters were tested in-situ and the remaining samples were transported to a laboratory for further physical, chemical and biological tests;
- The World Health Organisation's drinking water guidelines were used as a benchmark for drinking water contamination levels; and
- Critical observations of the nature of water systems and sanitary conditions in and around the project site.

Samples were collected from receptacles in the project area for water quality determination. A total of 23 water sources were assessed through in situ measurements using a water meter and/or sampled (6 sources) for laboratory analyses within and around the project site (12 underground wells, 6 streams and 4 locations in the Sewa River; all from a total of 20 communities, Table 3). Measurement of physical parameters was done in all these locations, whilst only 6 were selected for chemical and microbiological measurements in the laboratory, due to the critical values obtained from the physical measurements relative to the end use. Water samples were collected in sterilized polyethylene bottles.

Physical Analyses: Portable water testing meter, the *Maji Meter™* was used to measure the temperature, pH, electrical conductivity, turbidity and total dissolved solids of each water sample immediately it was obtained from the source.

Chemical Analyses: Concentrations of iron, manganese, chloride, magnesium, calcium, nitrate, carbonate, bicarbonate, aluminium and hydrogen ions were determined on a standard sample obtained from each field sample.

Microbiological Analyses: The membrane-filtration technique was used to enumerate faecal indicator bacteria.

Results and Findings

The results are shown in **Error! Reference source not found.** and 5.

Table 5.1-12: Sampling locations and characterization

Village	Source	Sample #/ In situ measurem ent location	Date	Time	Lat	Long	Altitude (m)	Comment
Heima	Sewa River	1	10/08/16	14:43	N07° 44.4678	W011° 54.2877	25	
Heima	Well	2		15:37	N07° 44.3844	W011° 54.0651	35	Constructed by GTZ, Bo Pujehun rural dev. Project. Iron and mud is a problem hence it is not regularly used for drinking
Heima	Well	3		15:56	N07° 44.3180	W011° 54.1125	46	Constructed in 2003
Benduma Sewa	Well	4	10/08/16	17:04	N07° 44.8890	W011° 54.7509	45	Constructed in 2002 by world vision, usual dried in march
Kpumbu	Stream	5	11/08/16	13:15	N07° 42.6062	W011° 53.8934	49	Used for drinking
Kpatema	Stream	6		13:45	N07° 42.5365	W011° 52.6616	61	Used for drinking. There are graves 15m from the stream
Benduma 2	Well	7		15:08	N07° 41.3383	W011° 53.4666	55	Constructed in 2007, usually dry in m\March
Gelehun	Sewa River	8		16:35	N07° 43.1323	W011° 55.5814	33	Used for drinking
Mumadu	Stream	9	12/08/2016	11:05	N07° 36.8596	W012° 00.0147	12	Used for drinking
Kormende	Well	10		11:56	N07° 36.0361	W011° 59.8211	17	
Kormende	Sewa River	11		12:02	N07° 36.0417	W011° 59.8535	25	
Bahuma	Stream	12		12:05	N07° 37.4656	W012° 00.1918	29	Used for drinking
Moforay	Well	13		15:00	N07° 38.1341	W011° 58.8344	2	Usually dry in march
Morkombo	Riparian	14		16:00	N07° 36.5439	W011° 58.1639	8	Drinking
Gola	Sewa River	15		16:50	N07° 58.0705	W011° 58.0705	34	No sample was collected
Mosorgbo	Well	16	13/08/2016	11:20	N07° 36.5194	W011° 56.6810	75	
Tawamaheh un	Stream	17		12:30	N07° 37.0396	W011° 57.2538	25	The water is called Gboneh, it is about one and half kilometers from the village.
Garinga	Well	18		13:08	N07° 37.8330	W011° 57.1225	30	Constructed by Bo Pujehun in 1989

Village	Source	Sample #/ In situ measurem ent location	Date	Time	Lat	Long	Altitude (m)	Comment
Bamba	Well	19		14:05	N07° 37.6971	W011° 55.8627	41	No sample was collected
Tungei	Well	20		15:55	N07° 37.2654	W011° 53.8312	40	No sample was collected
Gbandama	Stream	21		16:20	N07° 37.4546	W011° 54.2966	49	
Sumbuya Town	Well	22		17:04	N07° 38.8512	W011° 57.5865	51	
Lower Sama	Well	23		17:15	N07° 39.3460	W011° 56.3713	32	No sample was collected

Table 5.1-13: Sample Chemical Data

Location	Heima Sewa River	Heima Well	Benduma Sewa River	Gelehun Sewa River	Kormende Well	Kormende Sewa River
Sample #	S1	S2	S4	S8	S10	S11
Turb (NTU)	5.8	6.4	19.12	0.8	0.6	8.5
Cond.	28.8	142.4	14.24	30.7	148.6	30.2
TDS (mg/L)	14.4	71.2	7.12	25.3	74.3	25.1
Residual chlorine (mg/L)	0					
		0.01	0	0.01	0	0
Al (mg/L)	0.03	0.02	0.28	0.01	0.02	0.03
NH4 (mg/L)	0.02	0	0.28	0.01	0.01	0.02
Ca-hardness (mg/L CaCO3)	12	6	0	5	10	15
Cr (mg/L)	0.2	0.1	4.48	0	0.1	0.2
Cu (mg/L)	0.14	0.23	0.12	0.11	0.12	0.18
F (mg/L)	0.56	0.62	0.35	0.28	0.66	0.58
Dissolved Fe (mg/L)	0.6	0.4	0.3	0.2	0.2	0.8
Mg (mg/L)	27	25	15	10	12	28
Mn (mg/L)	0.4	0.3	0.2	0.1	0.2	0.5
NO2- (mg/L)	0.04	0.06	0.2	0.02	0.01	0.08
NO3-N (mg/L)	0.86	0.88	0.65	0.58	0.62	0.84
K (mg/L)	8.1	10.3	6.7	4.2	5.5	8.8
PO43- (mg/L)	5.4	6.2	4.8	3.3	5.1	6.7
SO42- (mg/L)	4	6	3	2	4	5
S2- (mg/L)	0.12	0.16	0.11	0.12	0.15	0.17
HCO3- (mg/L)	12	16	7	6	10	14
Zn (mg/L)	0.02	0.03	0	0	0.01	0.02
E. coli (cfu/100 ml)	45	20	10	5	25	50
Faecal coliform (cfu/100ml)	80	35	15	10	20	60
Non-faecal coliform (cfu/100ml)	20	10	0	5	15	30

Faecal indicator Bacteria: Faecal coliforms are a group of intestinal tract microbes and their presence in drinking water sources is an indication of faecal contamination. According to the World Health Organization (WHO), these bacteria should not occur repeatedly in drinking water. They therefore recommend zero faecal coliforms counts per 100 ml water sample.

A total of six (6) samples were analyzed and the results show that all 6 had high amounts of both *E. coli* and faecal coliforms per 100ml of water sample; most of the values were in excess of 20 counts/100 ml. four of these samples were taken from the Sewa River where the respective communities claimed to be coursing water for drinking and domestic use, namely Heima, Benduma, Gelehun and Kormende villages (**Error! Reference source not found.**). It is therefore obvious that such waters are contaminated. The main surprises were wells in Heima and Kormende villages (**Error! Reference source not found.** and **Error! Reference source not found.**), which were both well protected and at least 10m from the nearest dwelling. Contamination could possibly be linked to handling and roaming livestock.



Figure 5.1-25.1-3: Location in the Sewa River at Heima where water is fetched for drinking and domestic use



Figure 5.1-35.1-4: Sampling at Heima Village



Figure 5.1-45.1-5: Hand-pump well in Kormende

Dissolved Chemicals

Iron (Fe^{2+}): Iron is mainly present in water in two forms. It is either the soluble ferrous iron or the insoluble ferric iron. Water containing ferrous iron is clear and colorless because the iron is completely dissolved. When it is exposed to air, the water turns cloudy and a reddish brown substance begins to form. This sediment is the oxidized or ferric form of iron that will not dissolve in water.

Concentrations of iron as low as 0.3 mg/l will leave reddish brown stains on fixtures, tableware and laundry that is very hard to remove. Dissolved ferrous iron gives water a disagreeable taste. When the iron combines with tea, coffee and other beverages, it produces an inky, black appearance and a harsh, unacceptable taste. Vegetables cooked in water containing excessive iron could turn dark and look unappealing.

The analyses show that three (3) of the water sources have concentrations of iron above the WHO recommended value (0.3mg/l) for drinking water. These were the Sewa River at Heima (0.6 mg/L), the Heima well (0.4 mg/L) and the Sewa River at Kormende (0.8 mg/L). This could be attributed to the geology of the base rocks. Local inhabitants claimed that the Heima well is not used in the Dry Season due to the reddish appearance and unpleasant taste of the water. High Fe^{3+} content could be linked to high oxidation and corrosion, and the destruction of hand pump equipment.

Manganese (Mn): This is present in groundwater as the divalent ion (Mn^{2+}). High concentrations of manganese in water can cause dark stains in laundry and plumbing fixtures. It imparts an objectionable taste to beverage and tea.

However, deficiencies in manganese are rare but would include poor bone growth, problems with the disks between the vertebrae, birth defects, and problems with blood glucose levels and reduced fertility. Serious deficiency in children can result in paralysis, deafness and

blindness. Manganese is not easily absorbed but since small amounts are needed, deficiencies are not very general.

The analyses show that all water sources had manganese concentration above the secondary maximum contamination level (SMCL) standard of 0.05 mg/l by an order of magnitude.

Iron and manganese are non-hazardous elements that can be a nuisance in a water supply. Iron and manganese are chemically similar and cause similar problems. Iron is the more frequent of the two contaminants in water supplies; manganese is typically found in iron-bearing water. Water percolating through soil and rock can dissolve minerals containing iron and manganese and hold them in solution. Occasionally, iron pipes also may be a source of iron in water.

Iron and manganese can affect the flavour and colour of water. Iron will cause reddish-brown staining of laundry, porcelain, dishes, utensils and even glassware. Manganese acts in a similar way but causes a brownish-black stain. A problem that frequently results from iron or manganese in water is iron or manganese bacteria. These non-pathogenic (not health threatening) bacteria occur in soil, shallow aquifers and some surface waters.

There are several methods available for removing iron and manganese from water. These include (1) phosphate compounds; (2) ion exchange water softeners; (3) oxidizing filters; (4) aeration (pressure type) followed by filtration; and (5) chemical oxidation followed by filtration.

Nitrogen-Nitrate-Nitrite: Nitrogen is required by all organisms for the basic processes of life to make proteins, to grow, and to reproduce. Nitrogen is very common and found in many forms in the environment. Common sources of nitrate - nitrite contamination include fertilizers, animal wastes, septic tanks, municipal sewage treatment systems, meat preservatives and decaying plant debris. Nitrate is highly soluble in water and is stable over a wide range of environmental conditions. Ten (10) mg/l is the standard maximum contaminant level (MCL) for nitrate-nitrogen and 1 mg/l for nitrite-nitrogen for regulated public water systems.

The analytical result for the samples collected is found within the range of 0.01 – 0.88 mg/l. These results are below the WHO standards of 10 mg/l. The potential sources of nitrate could have been runoff from fertilizer use within the surrounding farms, sewage, and possible erosion of natural deposits from increased surface runoff. Excessive concentrations of nitrate can be harmful to humans and wildlife. Symptoms include shortness of breath and blue-baby syndrome.

The primary health hazard from drinking water with nitrate-nitrogen occurs when nitrate is transformed to nitrite in the digestive system. The nitrite oxidizes iron in the hemoglobin of the red blood cells to form methemoglobin, which lacks the oxygen-carrying ability of hemoglobin. This creates the condition known as methemoglobinemia (sometimes referred to as "blue baby syndrome"). Infants below the age of six months who drink water containing nitrate in excess of the maximum contaminant level (MCL) could become seriously ill and, if

untreated, may die. Nitrite in high doses is also linked with cancer as it can make carcinogenic nitrosamine and N-nitroso compounds.

Chloride (Cl⁻) - Chlorine is a chemical used in industry and in household cleaning products. At room temperature, chlorine is a gas. It has a yellow-green color, and a pungent, irritating odor similar to bleach. Chlorine does not catch fire easily, but may combine with other common substances to form explosive compounds.

Chloride in drinking-water originates from natural sources, sewage and industrial effluents, urban runoff containing de-icing salt and saline intrusion. The main source of human exposure to chloride is the addition of salt to food, and the intake from this source is usually greatly in excess of that from drinking-water.

No health-based guideline value is proposed for chloride in drinking-water. However, chloride concentrations in excess of about 250 mg/litre can give rise to detectable taste in water. The sample analyses for chloride levels were within the range of 0.00 – 0.01 mg/L and these values are below the WHO recommended maximum contamination level (250 mg/l). Excessive chloride concentrations increase rates of corrosion of metals in the distribution system, depending on the alkalinity of the water. This can lead to increased concentrations of metals in the supply.

When chlorine enters the body as a result of breathing, swallowing, or skin contact, it reacts with water to produce acids. The acids are corrosive and damage cells in the body on contact. Most harmful chlorine exposures are the result of inhalation. Health effects typically begin within seconds to minutes. Following chlorine exposure, the most common symptoms may include:

- Wheezing
- Difficulty breathing
- Sore throat
- Cough
- Chest tightness
- Eye irritation
- Skin irritation

The severity of health effects depend upon the route of exposure, the dose and the duration of exposure to chlorine. Breathing high levels of chlorine causes fluid build-up in the lungs, a condition known as pulmonary edema. The development of pulmonary edema may be delayed for several hours after exposure to chlorine. Contact with compressed liquid chlorine may cause frostbite of the skin and eyes.

Bicarbonate and Total Hardness - Bicarbonate is a major element in our body. Secreted by the stomach, it is necessary for digestion. When ingested, for example, with mineral water, it helps buffer lactic acid generated during exercise and also reduces the acidity of dietary components. Bicarbonate is an alkaline, and a vital component of the pH buffering system of the body (maintaining acid-base homeostasis). About 70 to 75 percent of CO₂ in the body is converted into carbonic acid (H₂CO₃), which can quickly turn into bicarbonate (HCO₃⁻).

The Bicarbonate (HCO₃) ion is the principal alkaline constituent in almost all water supplies. Alkalinity in drinking water supplies seldom exceeds 300 mg/l. The sample analyses for bicarbonate levels show a range of 6-16 mg/l. These values are below the expected maximum contaminant level which indicates that bicarbonate levels in the water sources are very low to almost being negligible.

However, bicarbonate alkalinity is introduced into the water by CO₂ dissolving carbonate-containing minerals. Alkalinity neutralizes the acidity in fruit flavors as well as acting as a buffer and acid dying.

Hardness generally represents the concentration of calcium (Ca²⁺) and magnesium (Mg²⁺) ions, because these are the most common polyvalent cations. Waters with high hardness values are referred to as "hard," while those with low hardness values are "soft".

Total Hardness is also the concentration of dissolved salts in water, expressed as total parts of dissolved salts in a million parts of water. The analysis from all samples indicates values below the recommended 100 mg/l (i.e. Ca²⁺ and Mg²⁺ concentration).

Hard water usually affects the amount of soap that is needed to produce foam or lather. It requires more soap, because the calcium and magnesium ions form complexes with soap, preventing the soap from lathering. Hard water can also leave a film on hair, fabrics, and glassware.

Aluminium (Al³⁺): Aluminum is the third most common element in the earth's crust and is present in soil, water and air. Aluminium's physical and chemical properties make it ideal for a variety of uses in food, drugs, consumer products, and water treatment processes.

The concentration of aluminum as recommended by the United States Environmental Protection Agency should range between 0.05 – 0.2mg/l of sampled water. The analysis gives a range of values that are below the recommended limit for aluminum concentration.

Although most of our daily aluminium intake comes from food, aluminum in food appears to be bound to other food substances and thus is in a form that cannot be absorbed into the bloodstream. In fact, it seems that the body's main defense against aluminum in food is that it does not allow aluminum to pass through the intestinal wall.

However, the amount of aluminium absorbed from drinking water is usually very small. One reason for this is that the presence of food in the stomach reduces the absorption. Absorption then is greatest when water is drunk on an empty stomach. In spite of this, the high concentrations of aluminium in water can increase the toxicity of the water because it has the

potential to reduce the pH level. When this happens, the successful growth of aquatic life may slow down to significant levels.

5.1.7.2.11 Physical Parameters

Table 5.1-14: Water Analysis Result for water sources around the project's concession area

Village	Source	Sample #/ In situ measure ment location	Temp (° C)	DO (mg/L)	pH	Turb (NTU)	TDS (mg/L)	EC (µs/ cm)	Sal (ppT)
Heima	Sewa River	1	27	6	7.31	7.9	0	0	0
Heima	Well	2	28.6	3.08	6.91	0	100	155	0.07
Heima	Well	3	29.4	6.57	5.26	0	7	12	0
Benduma Sewa	Well	4	29	6.99	6.44	0	105	163	0
Kpumbu	Stream	5	27.1	1.25	6.03	0	7	11	0
Kpatema	Stream	6	26.8	5.12	6.25	3.6	0	0	0
Benduma 2	Well	7	28.7	6.09	7.21	0	63	98	0.04
Gelehun	Sewa River	8	27.3	7.68	8.18	22	0	0	0
Mumadu	Stream	9	27.1	5.39	8.26	9.7	0	0	0
Kormende	Well	10	29.3	6.21	6.5	0	92	142	0.07
Kormende	Sewa River	11	27	7.52	7.12	16.6	0	0	0
Bahuma	Stream	12	27.5	5.32	7.57	6.3	0	0	0
Moforay	Well	13	27.3	7.2	8.03	43.1	0	0	0
Morkomb o	Riparian	14	28.5	0.41	6.47	6.3	0	0	0
Gola	Sewa River	15	27.4	4.22	6.02	0	0	0	0

Village	Source	Sample #/ In situ measure ment location	Temp (° C)	DO (mg/L)	pH	Turb (NTU)	TDS (mg/L)	EC (µs/ cm)	Sal (ppT)
Mosorgbo	Well	16	27.3	4.53	5.77	4.1	14	23	0.01
Tawamahe hun	Stream	17	26.9	1.06	4.92	0	0	0	0
Garinga	Well	18	27.8	5.26	5.07	4	0	0	0
Bamba	Well	19	27.8	5.67	4.49	2.4	0	0	0
Tungei	Well	20	29.3	6	5.49	0	147	226	0.11
Gbandama	Stream	21	26.8	5.71	5.6	0	0	0	0
Sumbuya Town	Well	22	29.7	6.02	5.45	13.3	52	79	0
Lower Sama	Well	23	28.2	6.31	5.92	0	8	13	0

Electrical Conductivity (EC): This is a measure of the ability of water to conduct electricity. It is an indication of the total amount of dissolved ions in water and this is related to the taste of water. Taste is an important factor in user's acceptance of water. In Sierra Leone, groundwater with EC values up to 450 µS/cm is considered 'good' while ground water with EC values greater than 850 µS/cm becomes progressively less potable.

The EC values for all the water samples were found to be less than the recommended 450 µS/cm. The highest conductivity levels from the analyses were seen in well water from Tungei, Heima and Kormende. The only source of water in the Sewa River with high conductivity value was at Benduma Village. The rate of conductance may be affected by certain factors which have the potential to impair the potability of a water source. Runoff from farms for example can contain fertilizers, which contain phosphate and nitrate. In addition, runoff from roads can also contain leaked automobile fluids which may eventually flow into nearby water sources. These compounds may have the tendency to increase the amount and mobility of ions when they break down and when this happens, conductivity level increases because they are negatively or positively charged when dissolved in water.

Total Dissolved Solids (TDS) - TDS are solids in water that can pass through a filter. It is a measure of the amount of salts or compounds dissolved in water. These compounds can include carbonate, bicarbonate, chloride, sulfate, phosphate, nitrate, calcium, magnesium, sodium, organic ions, and other ions.

A certain level of these ions in water is necessary for aquatic life. Changes in TDS concentrations can be harmful because the density of the water determines the flow of water into and out of an organism's cells (Mitchell and Stapp, 1992). However, if TDS concentrations are too high or too low, the growth of many aquatic lives can be limited, and death may occur.

Total Dissolved Solids are classified as a secondary contaminant and a recommended maximum limit is 1,000 ppm. Concerns with secondary standards relate to aesthetic or cosmetic quality of the water rather than health concerns. TDS can give water a murky appearance and detract from the taste quality of the water.

Gastrointestinal irritation in some individuals can be caused by high TDS levels. Water is considered hard when it has a relatively high concentration of calcium and magnesium ions. Water is referred to as hard when it presents difficulty to get a good lather from soap. This situation makes washing generally difficult.

The samples all show TDS levels way below the maximum recommended limit for drinking water (1,000 ppm) as recommended by the WHO.

pH - The pH is a measure of how acidic or basic (alkaline) the water is. pH stands for “potential hydrogen” or “hydrogen power” referring to the amount of hydrogen ions dissolved in a sample of water. pH is measured on a scale that runs from 0-14. Seven is neutral, indicating there is no acid or alkalinity present. The normal range for pH level in ground water lies between 6.5 and 8.5 as recommended by WHO.

Water sources in the southern section of the chiefdom recorded low pH values outside of the recommended WHO range; the observed range was from 4.49 – 5.92 in this area. 9 out of the 23 sampled water sources fell in this region, and would therefore be characterized as slightly acidic.

Many chemical reactions inside aquatic organisms (cellular metabolism) that are necessary for survival and growth of the organisms require a narrow pH range. At the extreme ends of the pH scale, (2 or 13) physical damage to gills, exoskeleton, and fins of aquatic plants and animals occurs.

Water with a low pH can be acidic, soft and corrosive. This water can leach metals from pipes and fixtures, such as copper, iron, lead, manganese and zinc. It can also cause damage to metal pipes and pose aesthetic problems, such as a metallic or sour taste, laundry staining or blue-green stains in sinks and drains. Low pH drinking water can be treated with a neutralizer.

Water Temperature – Temperature of water is a very important factor for aquatic life. It controls the rate of metabolic and reproductive activities of aquatic plants and animals.

Temperature also affects the concentration of dissolved oxygen and can influence the activity of bacteria and toxic chemicals in water.

Temperature readings for the sampled water sources range between 26.8 and 29.7 °C. These values are within the recommended permissible limits for potable water.

The rate of chemical reactions generally increases at higher temperature, which in turn affects biological activity. An important example of the effects of temperature on water chemistry is its impact on oxygen. Warm water holds less oxygen than cool water, so the water may be saturated with oxygen but still not contain enough for the survival of aquatic life. Some compounds are also more toxic to aquatic life at higher temperatures. Also, high water temperature enhances the growth of micro-organisms and may increase taste, odour, colour and corrosion problems.

Turbidity - This is a measure of the cloudiness of water, that is, the cloudier the water, the greater the turbidity. Turbidity in water is caused by suspended matter such as clay, silt, and organic matter and by plankton and other microscopic organisms that interfere with the passage of light through the water.

Although turbidity does not have a direct effect on health, it reduces the effectiveness of any disinfection procedures. Highly turbid water can lead to user rejection of a water source purely for aesthetic reasons. According to WHO, the threshold at which turbidity can be detected in water by the naked eye is above 5 NTU (Normal Turbidity Unit).

The turbidity values for samples analyzed show that eleven (11) of the water sources have turbidity values above the WHO recommended limit (1.1 – 2.7 NTU). The most turbid were the well at Moforay, Sewa River at Kormende and a common well at Sumbuya in order of reducing magnitude. The possible causes for such high value may be the high rate of soil erosion and sedimentation from sloped terrain, as well as the decaying of plants and animals. These have the tendency to alter the natural clarity of the water.

Conclusion

Water is a very important resource that needs to be maintained and preserved to ensure its potability for consumption. It is vitally important to every aspect of our lives. Monitoring the quality of surface water will help protect waterways from pollution. It is therefore essential for development practitioners to embark on activities that will protect water resources as well as water channels within local communities that solely depend on these sources for their survival.

The proposed agro-based project may result in the emission of significant levels of dust particles through land clearing and preparation activities. These particles when emitted into the atmosphere may be carried away by wind and deposited on surface water bodies. This condition could increase the concentration of contaminants for surface water bodies as well as increase turbidity levels of the water thus altering the natural clarity of the water body.

The presence of total coliform bacteria was found in all the samples collected and their concentrations exceed the zero limits for drinking water. Total coliform concentration was more evident in the well at Heima village at the North-western angle of the project concession area.

However, due to the poor quality of drinking and domestic water within the settlements surveyed, it is essential for the company to incorporate in their community development program, a comprehensive water resource management system which will ensure access to adequate and safe drinking water. The following are however recommended:

- Settlements without wells be provided with at least one well which can be frequently monitored for impurities;
- Well protection should be improved with proper lining and a cap (cover);
- At least one well from each settlement be identified for treatment to eliminate contamination, and to prevent further contamination;
- A Water Management Committee be established, and members should be trained at cleaning and maintenance of hand pumps and continuous treatment of well water against bacterial contamination; and
- A Water Management Plan be developed and incorporated into the company's main operational activities.

5.2 Biological environment

5.2.1 Ecology

The Lugbu Chiefdom is located in the Bo District, Southern Province of Sierra Leone, and is biogeographically part of the western extent of the Guinea-Congo forest biome and the Upper Guinea Forest. Historic and current land tenure system and land use practices have however extensively modified the landscape resulting in only few patches of forest now remaining. The landscape is now dominated by vast areas of agricultural fallow land (farm bush) as a consequence of the widespread traditional cut and burn agriculture. Much of the remnant forests are kept as sacred groves (society bush and cemeteries) and occur as gallery forest along river courses. No forest reserves exist within the chiefdom, but some of the gallery forest and sacred groves are still healthy and can be protected. There is also a proposed game reserve called Bo Plains, but this is located in the Gbo Chiefdom in western part of the Bo District and it is yet to be constituted.

The Lugbu Chiefdom and environs have experienced long-term diamond mining both from the industrial and artisanal sectors, leaving large areas of land unproductive. No published information exists on the ecology of the area, and there is no record of existing environmental impact assessment report for any past or current industrial or artisanal mining project in the area. This ecological report details findings of a six-day survey of the proposed concession zone, against the backdrop of very little or no baseline ecological information of the area. The survey constitutes a rapid assessment of key ecological indicators and the vegetation and landscape features of the area, an assessment of impacts and recommended mitigation

measures for the project, which will also constitute the first ecological impact assessment report in the area.

5.2.1.1 Methodology

This work was conducted in and around 17 sites/settlements within the project concession zone (Table 1). Data was collected between 9th and 15th August 2016, during the rainy season (which cover the month of April to October) and coincides with the peak of the growing season in country Sierra Leone (Birchall et al., 1979). The general methodology applied is consistent with a rapid assessment exercise and so no detailed long-term experimental sampling was done. However, the data collected was adequate for the purpose of the assignment. The following methods were used for the respective ecological themes:

5.2.1.2 Results

5.2.1.2.1 Vegetation and Botanic assessment

The ecology team visited various locations to observe, identify and describe the vegetation types that occur in the proposed project area. Assessment of the size and nature of vegetation cover, tree species, tree density and the diameter at breast height of trees were done to provide appropriate indications of the vegetation status. No transact assessment was done, but a general identification was done for wood plants (particularly trees and shrubs), focussing on the possible occurrence of species of conservation interest in the area.

Table 5.2-1: Site number, names, GPS data and comments of areas visited during the survey.

Site No.	Village/Town	Eastings	Nothings	Comments
Site 1	Heima	175849	847395	Mainly farm bush, farms, community/sacred forest good; Swamp vegetation exists
Site 2	Kpetema	182305	853225	Mainly Farm bush and farms; small community/sacred forest exist
Site 3	Kpumbu	180345	853433	Mainly Farm bush and farms; small community/sacred forest exist
Site 4	Momandu	168960	842499	Mainly farm bush and farms; a small sacred forest; Stream and associated flood plains exists
Site 5	Komande	169317	841305	Mainly Farm bush and farms; small community/sacred forest exist
Site 6	Baoma	830567	844176	Mainly Farm bush and farms; small community/sacred forest exist

Site No.	Village/Town	Eastings	Nothings	Comments
Site 7	Mokombo	172327	842210	Riparian ecology near village; sacred forest exist. Some farm bush and farms
Site 8	Ngola	172488	843963	Mainly Farm bush and farms; small community/sacred forest exist
Site 9	Mogorbor	174968	842043	Mainly Farm bush and farms; small community/sacred forest exist
Site 10	Tohamahehun	174450	843451	Plantations of oil palm and citrus; secondary forest exist. Some farm bush and farms
Site 11	Gahinga	174252	844584	Mainly farm bush and farms; a small sacred forest; Stream and associated flood plains exists.
Site 12	Bamba	176659	844275	Mainly farm bush and farms; small sacred forest; oil palm and citrus plantations exists
Site 13	Moforay	171258	845089	Gallery forest degraded; small patch of community/sacred forest. Some farm bush and farms.
Site 14	Sumbuya	173477	846440	Small town, thin gallery forest, Good community/sacred forest in places. Some farm bush and farms.
Site 15	Lower Sama	176113	847527	Mainly Farm bush and farms; small community/sacred forest exist
Site 16	Gendema	179772	843776	Mainly farm bush and farms; large oil palm plantation and small sacred forest
Site 17	Tongei	180311	843474	Mature forest/sacred forest close to village. Oil palm plantation nearby. Some farm bush and farms.

Nb. The site number and names is consistent with those used in the table of results.

5.2.1.2.2 Mammals Assessment

The mammalian diversity present in the area was assessed through direct field observations and semi-structured interviews with local hunters and farmers. Direct field observations were

carried out along established transects or foot paths and in forest locations and involved first hand sightings, footprints and faecal deposits of the animal. Semi-structured interviews (which provided much of the data on mammals) were conducted for at least three respondents in each of the 14 villages and settlements visited. Information provided by respondents was normally verified through triangulation.

5.2.1.2.3 *Herpeto-fauna (Reptiles and Amphibians) Assessment*

Assessment of the herpetofauna was carried out mainly by a random check on locations that possibly hold such species. Since reptiles are more active, they were either directly observed whilst moving about or by shaking of tree logs and stems to scare them out of their hide-outs, during which identification was done. In the case of amphibians, these were sampled by search for them under small rocks, logs, foliage and general observations. Local field guides, who are mostly farmers, were also asked about the type of snakes existing in the area.

5.2.1.3 Results and Discussions

5.2.1.3.1 *Vegetation and Botanic characteristics*

A 2016 Google Earth image of most of the areas covered in the concession shows a landscape mosaic dominated by bush fallows (farm bush) at various stages of succession, farms, human settlements and patches of secondary and gallery forest (Fig 1).

On the ground, the vegetation found in all areas visited is an exact reflection of the satellite imagery shown, with vast areas of farmbush (agricultural fallow vegetation), farms with cassava cultivation, some plantations of oil palm around villages and other places and small isolated patches of forest, most of which are sacred groves. No protected areas are found within the vicinity of the potential areas to be leased, except that sacred groves are generally protected by traditional bye-laws. Farm bush, are found to be mostly between one and five years. However, agricultural practices mainly involving cassava cultivation is very active in the chiefdom. The cassava is mainly processed into garri, an activity that constitute the main local industry in the area and a major income earner for the local communities.

Swamps form a very important component of the ecology in the Lugbu Chiefdom. The network form by the Sewa River and its associated tributaries and streams, is the main source of swamp inundation. Swamps are very vital substrates for agriculture, particularly for rice production by the local communities. However, the proliferation of mining activities over the last two decades is destroying swamp ecology and rendering substantial portions of swamp unfit for rice production. In fact, based on anecdotal evidence corroborated by some local respondents, rice production dropped significantly in swamps affected by mining activities, although agricultural activities are gradually regaining momentum.

From observation, most forests are secondary, except for a few sacred groves and the tract of gallery forest along the river banks that appears to be dwindling because it is very thin in most areas visited. Some gallery forests are contiguous with riparian ecologies found in places and these are vital ecosystems for a number of reasons. From experience and

information obtained locally, the forests are the main hosts of most of the mammal species identified, although the forest extent is small and is under constant threat of degradation and depletion because of agriculture and artisanal diamond mining.



Table 5.2-2: Google Earth (February 2016) image of part of the main areas and photos of the vegetation and landscape features of the Sierra Tropical Limited proposed concession zone in the Lugbu Chiefdom, in the Bo District.

The riparian ecology includes all areas inundated by natural flooding cycles along water courses and includes in-land valley swamps and wetlands. It is affected by seasonality, precipitation levels and stream flow, and so its ecology is flood-dependent. Flooding cycles cause natural disturbance regimes in riparian ecology, thereby facilitating species diversity and dispersal of seeds and plant propagules. This mechanism ensures that species are distributed in regular fashion to effect optimal ecological functioning. Riparian ecologies that are contiguous with gallery forest are observed to be the most stable, because of the constant supply of nutrient and the protection from adverse conditions provided by these forests. By inference, riparian ecologies and wetlands are vital component of the hydrological system in the area and to a large extent associated with water supply and swamp agriculture. Some areas around Sumbuya (along the banks of the Sewa river, Moforay, Galunga, Mokombo and Heima have ecologically viable riparian zones. The form of wetland in the area is the river and associated streams and flood plains, particularly along the western edge of the project area.

Fifty-five species of trees were identified belonging to 25 plant families (Annex C of the Appendices). Table 5.2-3 shows species that are listed as vulnerable by IUCN (2016), all of which are used as timber for various purposes such as housing construction, dug-out canoes and some are processed and sold for income generation. The gallery forest and small patches of community forest are the key sources of valuable tree species. Many fruit trees occur in and around the village settlements, the most common of which is was Oil palm *Elaeis guineensis*. A number of oil palm plantations were observed adjacent to all villages and settlements visited. Oil palm is a vital component of the socio-economic setting of the area, as many other areas in southern and eastern provinces in Sierra Leone. The fruit is harvested and processed to produce palm oil, which is mainly sold and the income used for subsistence and other purposes. Oil palm plantations are actually one of the biggest long-term agricultural investments by local farmers and their families.

Table 5.2-3: Tree species of IUCN conservation status observed forest patches and gallery forests within the project's concession zones

Botanical name	Family	IUCN Status	Local status	Main use
Afzelia africana	Caesalpinaceae	VU	Rare	Valuable timber
Garcinia afzelii	Guttiferae	VU	Rare	Timber
Heritiera utilis	Sterculiaceae	VU	Rare	Valuable timber
Lophira alata	Ochnaceae	VU	Rare	Valuable timber
Milicia regia	Moraceae	VU	Rare	Timber
Terminalia ivorensis	Combretaceae	VU	Rare	Valuable timber

Some of the wild oil palm trees are tapped for palm wine (a major beverage in these local communities), whilst the disused stands are logged and used as bridge over small drainages. A few cocoa/coffee and citrus farms were observed, particularly around Tawahama Hehun and Bamba. Stands of fruit trees common in all settlements visited, some of which are as

follows: Coconut, Bread fruit trees, Mango, Pawpaw; Banana; are very common in and around all settlements in the project area.

5.2.1.3.2 Mammals

Twenty-four species of mammals were identified by both visual evidence and interviews to occur across all sections of the project area (Appendix B). These include the following species of global conservation interest according to IUCN (2016): two endangered species (Western Chimpanzee *Pan troglodytes verus* and Red Colobus Monkey *Procolobus badius*); two vulnerable (Pied Colobus *Colobus polykomus*, Zebra Duiker *Cephalophus zebra* and Tree Pangolin *Phataginus tricuspis*) and two near threatened (Sooty Mangabey *Cercocebus atys* and Bongo *Tragelaphus euryceros*). The rest of the species are actually considered least concern (LC), meaning that there is no global threat to their occurrence and distribution; however, some of these least concern species are locally getting rare.

Table 2 also includes both the species' IUCN (2016) status and the local status; the latter indicate the occurrence and distribution status for each of the species listed. Most of the species that are of global conservation concern are very rare and respondents indicated that encounter rates for these species are seldom. In fact, the endangered and vulnerable species mentioned require adequate forest cover to survive, but with the contracting and increasing fragmented forest cover, it is very likely that these species might be locally extinct in the near future. Monkeys and antelopes are the main target species for hunting in these communities and this activity is further driving the rare species to the brink of local extinction and depleting the population of once common species. Also, in nearly all farms and farm bush, trap fences are installed to trap Marsh Cane Rat *Thryonomys swinderianus* which is also a delicacy, in addition to the monkeys and antelopes. Bush meat constitutes one of the main protein sources for people in the project area.

5.2.1.3.3 Herpetofauna

The herpetofauna in the project area is not impressive based on the data obtained. This may have resulted from two main factors: (i) the time and duration of the survey, which did not allow the installation of traps; and (ii) the high level of degradation, which may have driven some of the species into areas where they are difficult to reach. However, a total of 21 species of amphibians and reptiles were identified through visual evidence across all sites visited (Table 3). In the case of amphibians, 12 species belonging to six families were encountered. Among these were three species of conservation interest (IUCN, 2016) - *Hyperolius macrotis* (NT), *Petropedetes natator* (NT) and *Ptychadena arnei* (DD) – the last two are endemic to West Africa and depend on gallery forest, which is dwindling in the area. These species of global conservation interest are actually rare, but appear to be widely distributed in moist forest environment in the country. The family Hyperoliidae accounted for the highest number of species among amphibians. Amphibians are partly water-dependent (their main breeding habitat) and much of their foraging is done on land; a good diversity is forest dependent, and

so they are vulnerable to the widespread habitat characterising the forest degradation in the project area.

Table 5.2-4: Amphibians and reptile species encountered, their IUCN status and habitat distribution

Herpes group	Family	IUCN status	Habitat seen
AMPHIBIANS			
<i>Hyperolius concolor</i>	Hyperoliidae	LC	Forest
<i>Hyperolius lamtoensis</i>	Hyperoliidae	LC	Forest
<i>Hyperolius macrotis</i>	Hyperoliidae	NT	Forest/farm bush
<i>Leptopelis occidentalis</i>	Hyperoliidae	LC	Farm bush
<i>Hyperolius fusciventris</i>	Hyperoliidae	LC	Forest/farm bush
<i>Ptychadena pujoli</i>	Ranidae	LC	Farm bush
<i>Ptychadena arnei</i>	Ranidae	DD	Gallery forest
<i>Phrynobatrachus alticola</i>	Petropedetidae	LC	Forest
<i>Petropedetes natator</i>	Petropedetidae	NT	Forest
<i>Arthroleptis bivittatus</i>	Arthroleptidae	LC	Forest swamp
<i>Chiromantris rufescens</i>	Rhacophoridae	LC	Forest
REPTILES (Lizards)			
<i>Agama agama</i>	Agamidae	LC	Farm bush
<i>Trachylepis quiriquestaeniata</i>	Scinkidae	LC	Forest/farm bush
<i>Mabuya unimarginata</i>	Scinkidae	LC	Mature farm bush
<i>Varanus niloticus</i>	Varanidae	LC	Forest/farm bush
<i>Chamaeleo chamaeleon</i>	Chamaeleoninae	LC	Farm bush
REPTILES (Snakes)			
<i>Dendroaspis polylepis</i>	Elapidae	LC	Mature farm bush
<i>Dendroaspis viridis</i>	Elapidae	LC	Mature farm bush
<i>Naja nigricolis</i>	Elapidae	LC	Forest/farm bush
<i>Philothamnus irregularis</i>	Colubridae	LC	Farm bush



Species of herpes found in the project concession zone. Left to right - *Ptychadena arnei*, *Trachylepis sp* and *Agama agama*.

Nine species of reptiles of six families were recorded, among which were four species of snakes, two species of skinks, one species each of agama lizard, monitor lizard and chameleon (Table 3). The Rainbow Lizard *Agama agama* was most frequently encountered reptiles. Two of the snake species (Family Elapidae) are among the most venomous and are

responsible for most of the death from snake bite in rural communities in Africa (www.tlcafrica.com/tlc_snakes). Incidences of snake bites occur in the area, but not officially recorded; however, these incidences rarely result in death because, according to local people, most victims recover fully after treatment with traditional herbs.

5.2.1.4 Discussion of Impacts and Mitigation

The establishment of any large-scale agricultural venture means that vegetation would have to be modified to a monoculture that empirically has associated ecological and environmental problems (Altieri, 1995). With reference to the project concession area, the key impact would be the conversion of vast areas of land into a new ecological paradigm through loss of original vegetation cover including forest cover, which distorts ecological processes, causes ecosystem instability, increase pest problems and loss of species. Such negative ecological state has apparently been set in motion by the establishment of large cassava farms.

Although much of the original forest in the area has been modified into farm bush and farm lands, yet the remaining patches provide many ecological functions including stabilization of local climatic conditions, store of valuable genetic resources, store of carbon that would otherwise be injurious to the environment (the main culprit in the climate change phenomenon) and store of water associated with proper functioning of the hydrological system. With large-scale monocultures it is expected that the nutrient, energy, water and waste cycling will be distorted, whilst some naturally available resources (such as wood, timber and medicinal herbs) for use by local communities may be depleted. .

Upland and gallery forests support a diversity of both vascular and non-vascular plant species including epiphytic plants, lianas, macro-fungi, bryophytes and pteridophytes that are equally vital to the proper functioning of forest ecosystems. The loss of plant cover and possible key-stone species within the forest ecology, resulting from possible removal of some gallery forest and sacred groves may have irreversible consequences, if not controlled.

Large-scale monocultures also reduce the ability for natural enemies to control pest and plant disease, thus creating serious ecological imbalance on the food web and attendant environmental problems. According to Altieri (1995), pest explosion in monocultures results from the concentration of resources for specialist crop herbivores, which also increase areas available for pest immigration. Such pest problems could creep into local agro-ecology and farming systems, with devastating effects such as reduction in crop yield and loss of soil nutrient. Pest problems in Sierra Leone has mainly involved grasshopper invasion that tend to increase in open agriculture environment, increased rodents pests on local farming system, and viral plant pest on cassava and oil palm.

The main impact on fauna is related to the reductions in vegetation cover to be replaced by more open farm ecological setting, dominated by monocultures. This situation negatively affects the habitats for wildlife and biodiversity in general. The loss of vegetation and the remaining forest patch will displace wildlife species, exposing them to hunting, predation and other threats that would lead to population reduction and eventual local extinction. In addition, foraging habitats and breeding areas for species will be destroyed, whilst refugia for

juveniles and less aggressive wildlife are cleared, rendering these areas ecologically redundant to adequately support wildlife. The extent of habitat and home range size of some of the large mammals that exist in the area is expected to decline and so species that may not be able to find alternative habitats could go extinct locally, whilst others will migrate into nearby habitats and increase inter-specific and intra-specific competition. The implication to wildlife is that, whereas degradation-tolerant species such as rodents may increase in number, forest dependent species including amphibians and large mammals may decline significantly, distorting the predator-prey dynamics in the natural food chain, with potentially deleterious ecological consequences, including loss of biodiversity.

Increased human presence and equipment use in the area as people engage in various operational activities of the establishment will create new ecological problems. New types of waste will be produced in large volumes never experienced previously in such environments; some of these wastes may be injurious to wildlife, particularly herpeto-fauna. Increased tramping of substrate may adversely effect on soil organisms and soil structure and this may be injurious to ground-dwelling fauna such as amphibians. Increased human presence could also mean increase hunting of wildlife for subsistence, particularly large mammals and primate and may render them susceptible to local extinctions.

5.2.1.4.1 Recommendations

It is overly important that areas that are important for biodiversity such as sacred groves, gallery forests, swamps and riparian zones be reserved and protected. In fact, the lay out of the project zone must take into consideration the importance of the hydrological system. Care must be taken to avoid the deposition of debris, silt and waste into these sensitive areas in order to avoid ecological distortions that may put further stress on the already degraded ecosystem that has undergone many years of mining-related perturbations..

Biodiversity conservation and conservation management areas must be established within the concept of co-management with local authorities and their respective communities. This is to ensure regulated and controlled use of resources such as timber and wood which is required by the local people for various purposes. If practicable, based on collaboration with relevant institutions, a botanic reserve could be created for areas with reasonable forest cover that may not be under any traditional management system. Such reserves could serve as natural plant genetic store for genetic diversity and as a repository for rare or less common flora, including some of the endemic species.

Some patches of vegetation must be allowed between defined rows of the fruit crops and some connectivity must be established with forest patches where possible. This is necessary in order to enhance some level of ecological integrity of the modified landscape. Such a system will provided ecological corridors and allow movement of species between habitats and serve as refugia for a host of small mammals and juveniles of many species.

The historic and current state of deforestation by the local communities has, by default, created serious shortfall in the biomass resource needs of the community, in terms of fuel

wood and timber. As part of the environmental offset, the company could collaborate with local communities to establish and manage woodlots and/or agroforestry systems as a means of addressing the expected loss of wood biomass and other naturally available resources, once vegetation is cleared for industrial agriculture.

A proper waste management system should be put in place, to address all types of waste particularly, industrial waste water, industrial refuse and domestic refuse. This will ensure that viable ecological systems such as wetlands, riparian zone and community water supply systems are not contaminated or polluted, and people and wildlife are not exposed to environmental and health risks.

The data shows that dangerous snakes exist in the area encompassed by the project concession. With bush clearing and reduced forest cover, there is an increased chance that people will be exposed to snake bites more frequently.

5.2.2 Creation of Buffer Zones

In any large monocultures established a glaring effect is the modification of the vegetation and ecology in such way that once heterogeneous ecosystems and habitats at the macro-level and micro-level are reduced to a almost homogeneous state, which is injurious to biodiversity. In situations where the entire area is convert to an agricultural monoculture, there is very slim chance that any species not tolerant to disturbance would survive. In fact, the status and distribution of species of flora and fauna in such monocultures is expected to be drastically changed from its original state prior to conversion. To guard against the impact of large monocultures on biodiversity, there is need to reserve areas that are sensitive to change and are most important in maintaining some level of ecological functionality within concession zone. A number of sensitive sites have been identified (Table 5.2-5) which together would consitute a good representation of the original ecosytems and habitats that characterised the concession zone.




Table 5.2-5: Sites identified for the creation of buffer zones within concession zones. The sites are marked by yellow polygons.

Table 5.2-5 is a google map of the concession zone on which sensitive sites have been identified and mark by yellow polygons. These areas encompass a wide diversity of ecosystems and habitats which have been practically assessed during the field work, as sites that hold the largest complement of species associated with the concession zone. They include gallery forests, sacred groves, community forests, wetlands, streams, swamps and riparian zones. It is therefore important that these areas are reserved as buffer zones and remained untouched in order to provide functional habitats for species and to maintain some viability in the local gene pool, thereby sustaining some degree of species diversity. The sites identified with their central coordinates given in Table 5.2-6. The justification for the conservation of key ecosystems and habitats are explained below.

Table 5.2-6: Sites identified for buffer zone establishment within the Sierra Tropical Limited Concession zone

Site Number	Key habitat	Central coordinate
1	Wetland, floodplain and forest	171456 E & 841269 N
2	Gallery forest and community forest	172376 E & 847685 N
3	Gallery forest and community forest	172705 E & 846417 N

Site Number	Key habitat	Central coordinate
4	Community forest	174329 E & 843595 N
5	Community forest and swamps	175719 E & 847946 N
6	Community forest and swamps	176194 E & 846914 N
7	Community forest	180199 E & 843332 N
8	Community forest	182599 E & 852843 N

5.2.2.1 Habitat Corridors

Habitat corridors are areas of vegetation cover or wetland connectives which are used by animals to traverse large areas and to move across their home range during foraging and mate searching activities. These corridors include road site vegetation, streams, flood zones, and swamps. The corridor must also include everything that a species needs to live and breed, such as soil for germination, burrowing areas, and multiple other breeding adults (Beier & Loe 1992). At least 10 – 50 meters of vegetation cover should be allowed along the entire road network and around wetlands, swamps, riparian zones and gallery forest. This is necessary to create areas of activities for animals and to provide refugia for a diversity of fauna, whose habitat has already been depleted. In the case of road side vegetation, the cover also provides a source of ecosystem service for the local people, including wild fruits, shade against the hot burning sun when people walk along these roads and medicinal herbs. Habitat corridors (especially roadside vegetation) are also important for the survival and maintenance of the population of small mammals (squirrels, giant rats, cane rats) herpes (amphibians and reptiles) and a significant diversity of birds. Dispersal of seeds and the mixing of the gene pool through pollinations and mating is facilitated by the existence of passage for species, thus creating heterogeneity in populations and maintaining species diversity.

5.2.2.2 Sacred groves and community forests

Community forests including sacred groves are very important to biodiversity and the maintenance of the culture and traditional practices to the local communities. Sacred groves are among the most protected forest network in the country and these area's are held in trust by the chiefs and traditional heads for the observance of cultural and traditional rights of the people. Entry to sacred groves by non-indigenes is strictly prohibited except by special permission, usually preceded by a traditional ceremony. These sites by virtue of their protection level constitute some of the most important biodiversity concentrations in rural communities. By all means, it is important that sacred groves be protected and some buffer be allowed between cultivated zones and these very important community forests. Other types of

community forest need to be reserved to provide source of wood, timber and bush meat for the community. One of the key problems with the Socfin Project in the Pujehun District is the limited availability of wood, timber and other forest resources for use by the local inhabitants of most of the villages. Lessons should be learnt from these experiences to avoid discontent among the local communities.

5.2.2.3 Wetlands and water catchments

According to the Ramsar Convention, wetlands are “includes all lakes and rivers, underground aquifers, swamps and marshes, wet grasslands, peat lands, oases, estuaries, deltas and tidal flats, mangroves and other coastal areas, coral reefs, and all human-made sites such as fish ponds, rice paddies, reservoirs and salt pans”. The convention further describes wetlands as “indispensable for the countless benefits or ‘ecosystem services’ that they provide humanity, ranging from freshwater supply, food and building materials, and biodiversity, to flood control, groundwater recharge, and climate change mitigation”. As Sierra Leone is the 108th signatory to the Ramsar Convention it is incumbent on the country to protect and conserve wetlands, including areas within the Sierra Tropical Limited concession zone. These areas included the riparian zones and gallery forest along the main river course and associated with Njalated streams and tributaries, swamps and floodplains. In fact, the swamps and floodplains are vital components of the agro-ecology of the area and so are required to be reserved for maintaining food production through traditional agricultural activities of the local people.

5.2.3 High Conservation Value Assessment

Prior to any land clearing and development, a comprehensive HCV Assessment/Biodiversity study will be initiated. The study will be led by an environmental scientist from the Dole Philippines Research department with assistance from local students at the nearby Njala University.

The study will focus on the inter-relationship between agro-diversity conservation and sustainable use and development practices in Dole Plantations, including the proposed plantations Lugbu Chiefdom with emphasis on biodiversity.

5.2.3.1 Objectives

- To identify all existing natural ecosystems within the new Lugbu Plantation and immediate surroundings.
- To document the existing collection and count of flora (vegetation), fauna (animals), micro flora and fauna (plants and animals) among the newly identified Plantation areas and immediate surroundings.
- Establish biodiversity monitoring stations within the Lugbu Plantations and immediate surroundings.

- Using the baseline data collected, develop a possible program of Restoration, Protection, Conservation and the development of a Botanical garden within the new plantation areas.

5.2.3.2 Methods

1. Selection of Uncultivated areas
 - Low Elevation, Medium Elevation and High Elevation
 - 10 sampling sites (20-30 sq. m) distanced at 10-20 m apart
 - Located with GPS Coordinates
2. Count, Identification and Photographs
3. Data Gathering and Collection

For Flora:

- Leaf samples and other plant parts of unidentified plants will be collected for further identification.

For Fauna:

- Counting of animals (mammals, birds, insects, spiders etc.)
- Shaking of trees
- Sweep Netting will be used to capture low vegetation fauna (not visibly seen).
- Installation of trapping baits (improvised malaise traplight traps, water trap).
- As an update for the procedure for the year 2016 surveys and onwards, only Light traps will be installed in the survey sites and count of animals observed on site will also be recorded.

For micro flora and fauna:

- Samples will be gathered from soil pit, boulders underneath, soil-root area, tree trunks/ barks/ stem/ leaves, soil horizon O and others.
- Isolation of fungi and bacteria (Simple Plating Technique)
- Extraction of nematodes using the Baerman technique.

6 DESCRIPTION OF THE SOCIAL ENVIRONMENT

6.1 Introduction

The implementation of the STL Agricultural project will have both positive and negative social impacts throughout its various phases. This section highlights the baseline situation as it existed during the time of the ESIA study to serve as a benchmark for future reference purposes after the project would have been implemented.

6.1.1 National Socio-Political Context

Sierra Leone covers a total area of 71,740 km² and had an estimated population of 4.9 million in 2004 (Sierra Leone Population and Housing Census, 2004) but according to the 2015 Housing and Population Census result, the population has increased to 7,075,641. Political instability and poor economic growth led to the brutal and destructive 10 year civil war which officially ended in 2002.

The 2015 Global Human Development Report (HDR) titled Work for Human Development shows that Sierra Leone continues to make steady progress in human development despite the recent devastating Ebola epidemic in 2014-2015, but that deep poverty and various forms of inequality in Sierra Leone are among the highest in Africa.

The country moved one position up the Human Development Index (HDI) placing the country in 181 out of 188 countries with an HDI value of 0.413, but still below the 0.518 average for Sub Saharan Africa. Liberia is positioned at 177, Guinea at 182 and Ghana at 140 out of 188 countries. This means that Sierra Leone has overtaken its bigger neighbour Guinea for the first time while Liberia continues to lead its two bigger Mano River neighbours on the Human Development Index.

The report further shows that 77.5% of the population of Sierra Leone (about 4, 724,000 people) are multi-dimensionally poor even though income poverty (i.e. \$1.2 per day) is 56.6%.

Sierra Leone's gender inequality remains very high with only 12.4 percent of parliamentary seats held by women and only 10% of adult women have reached at least secondary level of education compared to 21.7% for their male counterparts. Gender Inequality Index (GII) reflects gender based inequalities in the areas of reproductive health, empowerment and economic activity. Sierra Leone's GII value in 2014 is 0.650 (rank 145 out of 155 countries). This implies that there is 65.0% loss in human development as a result of gender inequalities in reproductive health, empowerment and economic activity (UNDP, 2016)

Table 6.1-1 presents some of these gloomy socio-economic indicators.

Table 6.1-1: Information on National Social Indicators

Key Social Indicators	Rate	Source
GDP per capita	\$497.89 in 2015	http://www.tradingeconomics.com/sierra-leone/gdp-per-capita
Economic growth rate	13.8% in 2014	World Bank, 2015
Infant mortality rate	87/1000	World Bank, 2015
Life expectancy at birth	46 years	HDI, 2015
Maternal Mortality ratio	1410/100,000	World Bank, 2014
Population Growth rate	2.19%	(World Bank, 2014)
Adult literacy	44.46	(World Bank, 2012)
Illiteracy rate	55.54%	(World Bank, 2012)
Primary school enrolment	1,353,723	(MEST), (WAEC, 2008/2010)
Net primary enrolment rate	78%	(MEST, 2014)
Gender parity in primary school enrolment for boys and girls	1.1 : 1	(MEST, 2014)

Basic water and sanitation facilities for the majority of Sierra Leoneans is extremely limited due to the limited functional infrastructure for water supply as well as the increase in population in Freetown and provincial cities over the past decade as a result of the civil conflict (PSRP II, 2008).

From surveys done from the PRSP II document prepared by the Government of Sierra Leone, about 70% of the population live in absolute poverty, with expenditure below US\$ 1/ day. The average person's total consumption falls short of the minimum consumption level, by 27.5% of the poverty line. (PRSP II, 2008)

The most recent poverty profile prepared by The World Bank and Statistics Sierra Leone (2013) puts the incidence of poverty at 52.9 percent in 2011. The rural population which is about 70% (GoSL, 2012) is hardest hit with poverty headcount of 66.1 percent in 2011 (Ibid, 2013). Agriculture is the largest economic sector in the country. Nearly two-thirds of the population depends on it for their livelihood and it is responsible for almost half of the country's GDP. There has been a steady increase in domestic food production. For instance,

for rice, which is the staple food and the most common crop cultivated by majority of Sierra Leoneans, production increased at an annual rate of 17.8% between 2000 and 2010 compared to -7.1% between 1990 – 1999 (FAO, 2013). Nonetheless, the living conditions continue to be difficult especially for rural villagers who struggle to remain at subsistence levels. Poor health indicators reflect the lack of access among the population to basic service notably - health. Endemic diseases, especially malaria and HIV/AIDS, loom as a threat; in 1997, UNAIDS estimated the HIV prevalence among adults to be 3.2%. In 2002, a national prevalence survey estimated the rate at 5% while the survey in 2010 revealed an increase of 1.5% (UNDP, 2013).

Sanitary conditions are very poor as sewage and refuse disposal systems do not function effectively in most places. Urban living conditions are extremely difficult; (PRSP II, 2008).

Less than 10% of Sierra Leone's total population has access to electricity, compared to 49% in Ghana, 46% in Nigeria, 96% in North Africa, 73% in Asia, 99% in China and 76% global average. Only around 1% of the rural population in Sierra Leone has access to electricity.

Of the 11,300km of classified roads in the country, 8,148km are classified in the national road system. The remaining roads consist of urban roads, community roads, local roads and farm tracks. With respect to the regional distribution of roads, the Northern Province accounts for 41% of the roads followed by the Southern Province with 33% and the Eastern Province with 23%. The Western Area accounts for only 3% (PRSP II, 2008).

6.1.2 Local Governance Structure

Administratively, Sierra Leone is divided into four distinct areas: the Northern Province with its headquarters in Makeni, the Southern Province with Bo as its headquarters, the Eastern Province with Kenema as its headquarters and the Western Area comprising the Freetown Peninsular with Freetown as its headquarters.

Freetown is the capital city where most of the government ministries are located. District councils were established in the year 2000, with the appointment of management committees. The government is committed to decentralization and has held three Local Government elections in 2004, 2008 and 2012 and the elections scheduled for 2016 postponed to 2018. The elected councils constitute representative bodies with delegated powers and funds for local governance.

Councils are operating and the government is slowly devolving power and functions of various ministries to these devolved bodies.

6.1.3 Local Regional Socio-Economic Context

The information in this section was obtained from the Bo District Development Plan, 2014-2016.

Bo District is the largest of the four administrative Districts in the Southern Region of Sierra Leone. It is bounded on the north by Tonkolili District, to the west by Moyamba District, to the south west and south by Bonthe and Pujehun Districts, and to the East by the Kenema District. The District is sub-divided into 15 Chiefdoms, 94 sections and 1,654 Villages, each of which has long established traditional Chieftaincy for Local Administration.

Bo District occupies a total area of 7,003km² and a population of 574,201 people (Sierra Leone 2015 Population And Housing Census, 2016) of which about 173, reside in the cosmopolitan city of Bo. The district population is ethnically and culturally diverse, particularly in the city of Bo, however, over 60% of the population belongs to the Mende ethnic group; other ethnic groups include Temne, Limba, Susu, Mandingo, and Fullahs forming large settlements around Bo and the mining towns. Bo Town serves as the District and Regional headquarters and commercial centre for Bo District and Southern Region. The District is well fed by the Sewa River that flows east – centrally through Baoma, Tikonko, and Lugbu Chiefdoms, the Wanje River that flows south through the Wunde Chiefdom into the Pujehun District and the Taia that flows through Valunia and Gbo Chiefdoms into Moyamba District. Other big rivers include the Tabe and Dematoo in the Bumpeh Ngao and the Bagbo Chiefdoms.

6.1.3.1 Economic Activities

The major economic activities of the district population are gold and diamond mining, other activities include trading, agricultural production of rice and root crops, cash crops such coffee, cacao and oil palm plantation. Trading is also a livelihood means for many residents as the district serves the important trade route and business hub for the south west of the country. Traditional farming is a common livelihood and family income source for the majority of the population in the country, however, less than half (49%) of Bo residents are engaged in farming activities.

Agriculture

Agriculture is the largest economic sector in the district with more than half of the population of Bo District depending on it as their main source of livelihood. Thus economic growth and poverty reduction in the district will only be sustained with development in this sector as the District is endowed with cultivable land (Upland and Inland Valley Swamps) and adequate rain-fall. The expansion of this sector in the district will stimulate growth in other sector like the supply of input, job creation, marketing, processing etc.

There has been a steady increase in agricultural production - crops, livestock and even fisheries. Thus household food consumption has increased with regards to specific food like rice, cassava and potato.

In Bo District the consumption and production of rice has increased as the staple food crop due to progress in the expansion of land cultivated, increase in the farormation is availablen

families, donor and government input assistance to farmers through the ABCs, mechanized farming services etc. However rice production is still insufficient to meet the total consumption rate of the people in the district and thus there is a need to relay on imported rice. Root tubers such as cassava and potato have being the key substitute for rice in the district. Also the production of livestock such as (goats, sheep and poultry) and cash crops such as coffee, cocoa has also increased in the district.

Despite the progress made so far, the sector still face challenges in the district, among them are:

- Limited access to credit and saving facilities hereby limiting farmers to access loans for input.
- Inadequate budget allocation from centre Government.
- Limited irrigation system thus farmers depend on rainfall.
- Farmers depend on manual labour instead of the use of machinery to carry out their agricultural activities.
- Weak extension system due to low capacity of personnel.
- Productivity is hinder due to bad roads, transport, markets, energy and power.
- Weak coordination among partners in the sectors.

Housing

The quality of construction materials used in housing construction are often used as proxy indicators for assessing household poverty.

The housing condition in the district is poor with overcrowded houses and lack of minimal basic facilities. Foundations of most houses are also being destroyed by erosion because of poor drainage system.

Some residents have water facilities in their houses, while only a very few have bathrooms and toilet facilities in the house. Most have no electricity.

Mining

Mining activities in Bo District are concentrated in the Baoma, Tikonko, Lugbu, Kakua, Jaicama Bongor and Badjia Chiefdoms for diamonds and in Valunia, Selenga and Niawa-Lenga Chiefdoms for gold and Bumpe Ngao for Bauxite. The full mineral potential of the District is still to be explored even though illegal mining activities along the riverine banks and valley swamps fed by the Sewa, Moa, Taia, Tabe, and other large streams is a challenge in the district.

Industry

Industrial development in the District is poor. However, strides have been taken to promote cotton industries and small enterprises, particularly in construction, transportation, carpentry, clothing and blacksmithery. Development of industries for food processing is still poor and primitive.

Fishing

Fishing activities are limited as Bo District is land-locked. The District heavily depends on the weekly fish markets from Gbondappi (Pujehun District), Gbangbatoke and Shenge (Moyamba District), and Mattru Jong (Bonthe District). The District also heavily relies on fresh-water fish harvested from the streams and rivers that flow through the District. The result, therefore, is a high rate of malnutrition, especially among children and lactating women in villages.

Trade

The Lebanese and Fullah communities largely control Trade in Bo District. Indigenous trading is still at the petty-trader level because indigenous entrepreneurship is poorly developed, and credit facilities are not easily extended to locals with demands for high collaterals that are difficult and somehow impossible to meet. Most of the trading in Bo District is done at the weekly markets or “Ndorwei” currently organized at Barthurst (Wunde Chiefdom), Koribondo (Jaiama-Bongo Chiefdom), Gerihun (Baoma Chiefdom) and Mani Junction (Bagbo Chiefdom), however, the existing weekly markets will be improved and daily markets constructed. Also the private sector will be encouraged to take the lead for a rapid economic growth.

6.1.3.2 Education

There are 433 Primary schools and 28 Junior Secondary Schools in Bo District; the pre-schools are largely private owned and are not well-established. The participation rates decrease from primary to Junior Secondary Schools as well as the female participation when compared to males. The teacher to pupil ratio stands at 1:50 in primary and 1:35 in Junior Secondary School, but looking at these rates they are no better considering the number of out of schools.

The sector experiences a number of challenges that needs to be addressed

- Overcrowding in schools, limited school infrastructure including WASH
- Deplorable existing school structures
- Limited libraries/reading facilities
- Inadequate school furniture and school material-Teaching Learning Material

- Low capacity and morale of teachers-untrained and unqualified
- Inadequate facilities for Non-formal education for parent and vocational institution for drop out.
- Non-functional school feeding programme.
- Inadequate or damage staff quarters
- Exclusion of Inclusive Education-girls education disables ECD.
- Lack of science equipment and labs in JSS schools.
- Effect of Sporting in schools to academic work.

6.1.3.3 Health and nutrition

Health services in Bo District are provided by government institution, NGOS, Private Hospitals and clinics. Access to health services in Bo District has increased from 40% to 65% as per the Ministry of Health & Sanitation report due to the “Free Health Care” Policy. The Periphery Health Units, the smallest units for health delivery, are the major facilities in the district providing health services to the rural communities. The district has three hospitals that support the PHUs as they serve as referral points for complicated cases.

There is still high infant and maternal mortality rate even with the Free Health Care initiative. Poor sanitation in the District (especially in the rural areas) is also a concern. The District’s health status poses a major challenge mainly because of:

- High illiteracy and poor understanding of the spread of diseases;
- Large household population and congested community layout;
- Traditional norms about diseases and refuse disposal
- The high poverty level of people.
- Low awareness on the “Free Health Care” Policy

1.1.1.1 Water and Sanitation (WASH)

The Ministry of Energy and Water Resources (MEWR) comprehensive mapping of water points report 2012 indicated that the major drinking water sources for the district residents are wells, hand pumps, public water supplies (piped) and other sources (streams and untreated sources). There are some 3,656 functional water points of different sources, majority (2,412) of which are wells without a pump. During the time of the mapping exercise in 2012, 22% (797) water points were found not functioning, 275 of these sources need repairing. The Ministry of Education, Science and Technology (MEST) census for the 2012-13 school year indicated that 344 (47%) schools have safe drinking water sources (piped supplies and boreholes) inside the school compound while other schools are using wells, streams and other

untreated sources. 69% (499) of schools have access to toilet facilities within the school premises. (UN Office for the Coordination of Humanitarian Affairs, 2016)

6.1.3.4 Infrastructure

Roads and Works

The condition of major roads especially feeder roads in the District have deteriorated and suffered seriously from neglect and require extensive rehabilitation. All the bridges and culverts on the roads inspected require rehabilitation or reconstruction.

6.1.3.5 Information and Communication Technology Sector

Postal services do not function within the district. Mobile telecommunications companies, Africell and Airtel operate within the district as well as 5 radio stations. These two means of communications have contributed immensely to the communication systems in the district.

6.1.4 Socio-Economic Condition of Project Area (Lugbu Chiefdom)

Lugbu chiefdom is one of 15 chiefdoms in the Bo District. The general occupations of the inhabitants are agriculture, diamond mining and fishing. Agriculture, engaged in by more than two thirds of the inhabitants, is dominated by crop farming; this is mostly done on subsistence basis and employs rudimentary production techniques with the main crops being rice, cassava, groundnut, potatoes, yam, cocoyam and palm oil plantations. Fishing is also done in the Sewa River for both subsistence and commercial purposes.

Politically, Lugbu chiefdom is in ward 249 and 250 of the local council and part of constituency 75. The chiefdom is divided into 6 sections (Yarma, Gbo, Magbao, Kemo, Kamba and Kargbevu sections) and 44 villages. The main ethnic group in the chiefdom is the Mende, with smaller proportions of Sherbro, Temne, Fullas and Loko. These all have their own traditions, but there are broad cultural similarities among them, including the traditional secret societies, use of the Krio language, and their religions (traditional Christianity and Islam). The sacred sites of cultural importance include male and female secret society groves, ancestral cemeteries, shrines etc.

Education levels in the Lugbu chiefdom are low and illiteracy rates are high with many families unable to afford formal or informal school fees. The chiefdom has a total of 30 primary schools (4 of which are unapproved schools), 3 junior Secondary School and 1 Senior Secondary School (Sir Albert Senior School- Sumbuya).

The health sector is not well established in the chiefdom. Peripheral Health Units (the smallest units for health delivery) are the main facilities in the chiefdom providing health services. The chiefdom has 9 health centres with about 5 villages per health centre. The Mattru Jong Hospital and Government hospitals in Bo serve as referral points for complicated cases. There is still high infant and maternal mortality rate even with the Free Health Care

initiative (Development Planning Department Bo District Council, 2016). Poor sanitation in the chiefdom is also a concern.

Table 6.1-2: Population of Lugbu Chiefdom

Chiefdom	Male	Female	Total	Sex ratio
Lugbu	12,525	12,822	25,347	0.977




Table 6.1-3: Social Facilities in Lugbu

6.2 Socio-economic Status and Living Conditions

6.2.1 Methodology Used for Socio-Economic Baseline Survey

The social baseline study involving a review of available data and appropriate literature materials on the project area of influence, was followed by a reconnaissance visit in mid June 2015, followed by field investigations in late June to August, 2016 by various social experts to both ascertain ground-truth facts contained in the literature, and to obtain primary data for this report.

The social study was carried out using participatory techniques and aimed at facilitating and enhancing awareness, mutual understanding, trust and capacity building. Information was

collected on the following issues:

- Socio-economic conditions and land use patterns of affected communities,
- Formal and informal governing structures,
- Local infrastructure (transport, housing, health),
- Farming, Fishing, mining and other socio-economic activities and relevant statistics, and
- Income and expenditure trends.

Data analysis was carried out for both primary data collected at the focus group meetings and household surveys for landholders as well as the secondary data collected by means of a desktop review of existing data sources to gather relevant socio-economic baseline information at a national, district, and local level.

6.2.1.1 Project Area of Influence

The socio-economic baseline survey covered twenty-two (22) towns and villages in the Lugbu Chiefdom as outlined in the following table:

Sumbuya Town	Gelehun Village
Moforay Village	Bamba Village
Garinga village	Momandu Village
TawamahehunVillage	Mosomgbo Village
Komende Village	Gola Village
Makombo Village	Baoma Village
Upper Saama Town	Heima Village
Yambama Town	Tungie Village
Bendumasewa Village	Gbaloahun Village
Gelehun Village	Kpumbu Village
Kpatema Village	Lower Saama Village

6.2.1.2 Primary Data Collection

The 22 affected villages in the Lugbu Chiefdom constituted the sample frame from which a total of 121 randomly selected landholders were drawn and targeted for questionnaire survey. These villages were accessed by often narrow and unpaved roads with palm log bridges in

many instances. In some cases certain villages could only be accessed by motor bikes. The questionnaire was administered to the head of the land holding family, or - in the rare event that the head was absent - their spouses or a responsible member of the household. The survey generated information about households' demographic and socio-economic characteristics, health, and sanitation practices and experiences, sources of livelihood and agricultural activities, economic activities and income, social services such as access to health facilities, disease prevalence, and educational infrastructure and facilities. Data was also collected on their perceptions on the possible impacts the proposed agricultural project will have on their lives and their communities. MS Excel was utilised to capture the data and SPSS for analysis. The key findings from the analysis of the data derived from the survey are presented in this section.

The fieldwork was conducted using three key data collection methods:

- **Focus Discussion Group Meetings**

One venue was selected in each of the 6 sections for this meeting and stakeholders from the 22 villages likely to be affected were invited together with chiefdom authorities.

- **Administration of Household Questionnaires.**

Landholder questionnaires were administered at random to land holding heads in each of the 22 villages that will most likely be affected by the implementation of this project.

- **Meetings with Local and National Authorities.**

In order to identify key socio-economic activities in the affected chiefdoms, meetings with the appropriate authorities also took place.

6.2.1.3 Survey Tools

Inventory checklists and a landholders questionnaire developed for this survey aided the gathering of relevant data used for analysis. As stated in the methodology above and as part of the action-research approach, landholder questionnaires were administered randomly to 121 land holding heads in 22 towns and villages in the Lugbu chiefdom. The survey instruments included:

Landholder Questionnaires

The questionnaires solicited data from landholder/household heads on selected socio-economic and demographic information as stated above.

Landholder listing was done in the field by the social team and random sampling techniques

were used to select the landholder to be interviewed. The enumerators interviewed the landholder/household heads or a responsible person (preferably the household head's spouse).

The questionnaires were administered by trained enumerators in both Krio and Mende Languages. These interviews lasted for about 30 - 35 minutes per landholder and were conducted during the daytime. The enumerators carefully crosschecked responses in order to reduce observational errors arising from the use of this method. At the end of each day, the team met to share their field experiences as well as to identify ways of overcoming potential problems.

Farmers' Operations Checklist

The farmers' operations checklist also aided in learning the type of farming practised (subsistence or commercial farming), whether the villages have farmers' organisations; what rules and laws govern such an organisation; the kind of labour employed; how their farm products are sold and if they pay taxes or receive help from the government or non-governmental organisations.

Miners' Operations Checklist

The miners' operations checklist also aided in learning the type of mining practised (artisanal or mechanised mining), whether the villages have miners' organisations; what rules and laws govern such an organisation; the kind of labour employed; methodology employed in the mining process, how/where their gold and diamonds are sold and if they pay taxes on their revenue

Focus Group Discussion Checklist

The checklist of questions for the focus group discussion (FGD) solicited information mainly on the perceptions of the likely impacts the agricultural project might have on the economic activities especially the fishing activities and the communities as a whole. Mitigating measures to reduce perceived negative impacts from the project were discussed. Also discussed were the necessary community needs.

6.3 Results of Questionnaires administered to Landholders

6.3.1 Status of Respondents

Questionnaires were successfully administered to 90% of targeted participants (landholder/household head). The remaining 10% of participants included household head's son/daughter (5%), spouse, parent or brother/sister.




Figure 6.3-16.3-1: Status of Respondent

6.3.2 Gender of Landholder

Results revealed that about 92% of landholders are male.




Figure 6.3-26.3-2: Gender of Landholders

6.3.2.1 Age of Landholders

The following bar chart shows the age ranges of landholders interviewed. The majority (40%) fall into the 30 to 49 year age bracket.




Figure 6.3-3: Age Distribution of Landholders

6.3.3 Marital Status of Landholder

Landholders were asked their marital statuses, and in the case of respondents which were not the landholder/household head, what the landholder/household head's marital status was. 95% were reportedly married. Male landholders were asked how many wives they had (female landholders were not asked as polyandry is not practised in Sierra Leone). Monogamy was found to be prevalent in the communities (Figure 6.3-5).




Figure 6.3-4: Marital Status of Landholder




Figure 6.3-56.3-5: Number of Wives per Male Landholder

6.3.4 Number of households living in a dwelling unit

One household per dwelling unit is not common in these communities, with only 10% reportedly in this living condition. Over 45% of respondents lived in houses with two or more households.




Figure 6.3-66.3-6: Number of households per dwelling unit

6.3.5 Educational Level of Landholders

The educational level of landholders was investigated over a range of various academic levels of formal and informal education. 59% of landholders had never had any kind of education as shown in the following figure.




Figure 6.3-7: Educational Level of Landholders

6.3.6 Size of Landholders' Land

Landholders were asked to indicate the size of their land. The results are shown below.

Table 6.3-1: Size of Landholders' Land

Size of Land (hectares)	Frequency	Percentage
1 – 20	12	9.9
21 - 40	17	14.0
41 - 60	19	15.7
61 - 80	17	14.0
81 - 100	31	25.6
101 - 120	5	4.1
121 – 140	4	3.3
141 – 160	5	4.1
161 - 180	0	0
181 – 200	4	3.3
Over 200	7	5.8
Total	121	100.0

6.3.7 Availability of Sufficient Land for food production

Landholders were also asked to indicate whether their land was big enough to cater for their food production requirements, to which the majority reported that it was.




Figure 6.3-86.3-8: Size of Land Sufficient for Food Production Requirements

6.3.8 Income from Land Cultivation

Landholders were asked how much income they were able to generate annually from the cultivation of their land; the majority (62%), indicated between Le 1 million and Le 4 million per year.




Figure 6.3-96.3-9: Income Generated from Land Cultivation

6.3.9 Proportion of upland to be leased to Sierra Tropical

The following bar chart shows the result of landholders' responses to the question of what percentage of their upland they were willing to lease to STL. 45% indicated one third, while 37% indicated half of their land.




Figure 6.3-10: Proportion of upland land to be leased to Sierra Tropical

6.3.10 Sufficiency of Remaining Land for Family's Food Cultivation Requirements

When asked whether the remaining land left over after leasing a portion to STL would be enough for their family food production, an overwhelming majority (94%) reported that it would.




Figure 6.3-11: Sufficiency of Remaining Land for Food Cultivation Requirements

Landholder

6.3.11 Preferred Land Lease Payment Method

Landholders were asked how they would prefer the lease rent payments to be made. The following table highlights the results. The majority (57%) indicated 'by cash to each landholder'.

Table 6.3-2: Preferred Lease Rent Payment Method

	Frequency	Percent
By cash to every landholder	57	47.1
By cheque through community Bank	33	27.3
Through village Authority	31	25.6
Total	121	100.0

6.3.12 Intended use of lease rent

Landholders were asked what they planned to do with the lease rent money; 47% responded that they would build a new house.




Figure 6.3-126.3-13: Intended use of lease rent to landholder

6.3.13 Main Occupation of Landholder

The overwhelming majority of landholders are reportedly farmers.




Figure 6.3-136.3-14: Occupation of Landholders

Incomes reportedly generated from these primary occupations are detailed in the following table:

Table 6.3-3: Income from Primary Occupation

Amount	Frequency	Percent

Not Stated	2	1.7
Up to Le100,000	31	25.6
Le 101,000 - Le150,000	3	2.5
Le 151,000 - Le 200,000	8	6.6
Le 201,000 - Le 400,000	5	4.1
Le 401,000 - Le 600,000	7	5.8
Le 601,000 - Le 800,000	10	8.3
Le 1m - Le 2M	45	37.2
Le 2.1M to Le 3M	5	4.1
Over Le 4M	5	4.1
Total	121	100.0

6.3.14 Secondary Occupation of Landholder

Landholders were asked whether they engaged in any other income generating activities in addition to their primary occupations. Responses are highlighted in Figure 6.3_14 and income generated monthly from these secondary occupations outlined in




Figure 6.3_14: Secondary Occupations




Figure 6.3-156.3-16: Income from Secondary Occupation

6.3.15 Landholders' Awareness and Perception of STL Project

Respondents were asked whether they were aware of the STL agricultural project in their communities, and whether they thought it is a positive or negative initiative. 98% felt it would be a good project while the remainder did not have any particular perception about whether it would be good or bad.

Table 6.3-4: Landholder's Awareness of Project

	Frequency	Percent
Not stated	1	0.5
Yes	188	95.4
No	8	4.1
Total	197	100.0




Figure 6.3-166.3-17: Perception about Project

6.3.16 Perceived Impacts of Project

Positive outcomes identified by respondents include community development and job opportunities (Figure 6.3-17). Fears about possible negative impacts were also investigated and are outlined in




Figure 6.3-176.3-18: Perceived Positive Outcomes of Project

Table 6.3-5: Perceived Negative Outcomes of Project

	Frequency	Percent
Not Stated	37	30.6
Corruption in recruitment	2	1.7

Inadequate land for farming	4	3.3
Poor Management	34	28.1
Land grabbing	13	10.7
Inadequate job facility	28	23.1
Lack of security	3	2.5
Total	121	100.0

6.3.16.1 Recommended Remedies for Perceived Negative Outcomes

When asked what they felt should be put in place by the company to prevent the negative outcomes which they felt would occur, most indicated the implementation of proper systems and planning.

Table 6.3-6: Proposed Remedies for Perceived Negative

	Frequency	Percent
Not Stated	37	30.6
Proper land lease arrangements	16	13.2
Proper paying arrangements	7	5.8
Proper Management System	53	43.8
Good security network	8	6.6
Total	121	100.0

6.3.17 Food Security

Landholders were asked whether their households had experienced food shortage in recent years. 70% reported that they had.




Figure 6.3-186.3-19: Experience of Food Shortage

Months during which food shortages occurred in spite of a relatively good harvest were investigated, with the following result:

Table 6.3-7: Months during which Food Shortage is Experienced

	Frequency	Percent
July	2	1.7
August	11	9.1
July to September	91	75.2
August to September	14	11.6
May to July	2	1.7
November	1	.8
Total	121	100.0

Months during which households experienced food shortage following a particularly poor harvest are highlighted in the following table:

	Frequency	Percent
From June to September	26	21.5
From Feb, to the next harvest	67	55.4
July to September	14	11.6
Throughout the year	14	11.6

	Frequency	Percent
From June to September	26	21.5
From Feb, to the next harvest	67	55.4
July to September	14	11.6
Throughout the year	14	11.6
Total	121	100.0

Respondents were asked what they did to cope during food shortages:

Table 6.3-8: Coping Strategies during Food Shortages

	Frequency	Percent
Sale of animals	29	24.0
Sale of other household assets	4	3.3
Obtain loan to purchase food	82	67.8
Others	6	5.0
Total	121	100.0

6.3.18 Health

The various types of illnesses experienced by each landholder's family are documented in Figure 6.3_19.




Figure 6.3-196.3-20: Illnesses experienced by Landholders' Families

Sources of health care treatment listed by respondents are primarily from the local hospitals and Community Health Centre.




Figure 6.3-206.3-21: Sources of Health Care

6.3.19 Education

6.3.19.1 Primary School Enrolment

Landholders were asked whether they had any children of primary school-going age in their households.




Figure 6.3-21: Children of Primary School age (6 – 13) in Landholders' Households

91% of these children of primary school age were reportedly attending school.




Figure 6.3-22: Children in Landholders' Households of Primary School Age, Enrolled in School

Reasons given for the 10% not attending school are as follows:

Table 6.3-9: Reasons why children of primary school age are not in School

	Frequency	Percent
No Primary school in community	4	3.3
Cost of schooling not affordable	3	2.5
Not Applicable	114	94.2

	Frequency	Percent
Total	121	100.0

In spite of the encouraging percentage of children enrolled in and attending school, respondents reported facing difficulty meeting school fee requirements and other expenses.




Figure 6.3-236.3-24: Ease of Meeting Financial Requirements of School

6.3.19.2 Secondary School Enrolment

Landholders were asked whether they had any children of secondary school-going age (14 – 23) in their households.

76% reported that they did




Figure 6.3-246.3-25: Landholders with Children of Secondary School Age in their Households

When asked whether the children attend secondary school, 59% responded that they did.




Figure 6.3-25/6.3-26: Children of Secondary School Age attending School

The following reasons were given by those with children (14 – 23) not enrolled in secondary school:

Table 6.3-10: Reasons Children for Children not in School

	Frequency	Percent
No secondary sch. in community/school too far away	64	52.9
Cost of schooling not affordable	29	24.0
Early pregnancy	9	7.4
Unwanted pregnancy	7	5.8
Commercial bike riding	6	5.0
Marriage	6	5.0
Total	121	100.0

6.3.20 Water and Sanitation

Respondents were asked about their households' primary water source. Most indicated surface water sources:




Figure 6.3-26.3-27: Households' Primary Water Sources

They were further asked to describe the quality of the water obtained from these sources:

Problem with water	Frequency	Percent
None Stated	52	43.0
Colour	7	5.8
Taste	7	5.8
Both taste and Colour	29	24.0
Poor smell and taste	14	11.6
Colour, taste and smell	12	9.9
Total	121	100.0

Respondents were also asked what kind of toilet facilities they had and alternatives used if none. Most, unfortunately, reported using the bushes for this purpose:




Figure 6.3-276.3-28: Types of Toilet Facilities or Alternatives

Methods of waste disposal were also identified; again the bushes were indicated as the primary means of disposal:




Figure 6.3-286.3-29: Waste Disposal Methods

7 IDENTIFICATION OF POTENTIAL IMPACTS

7.1 Introduction

This chapter identifies and describes the potential environmental and social impacts of the project components on the biophysical and socioeconomic conditions of the environment and communities. Where applicable, it also identifies mitigation measures that will reduce adverse impact and enhance positive ones. The assessments carried out in this chapter are on potential impacts on overall environmental and social receptors caused by the project activities in the construction/development, operational and decommissioning phases with mitigation measures recommended accordingly.

7.2 Environmental and Social Impact Assessment

7.2.1 Methodology

As the method of assessment of project impacts may be subjective and sometimes difficult to quantify, the experience of the project team was relied upon to assess such impacts. The knowledge of the project area by team members was invaluable in analyzing impacts. A number of steps were followed:

- i. A matrix of important project-specific impact categories was prepared;
- ii. The level of significance, achievability of mitigation steps measured against practicality and cost-effectiveness were discussed in a workshop/meeting setting;

An impact assessment scale was then developed.

Table 7.2-1: Degree of Certainty of Impact

Certainty of Impact	Description
Certain	The impact will occur.
Very Likely	It is expected that this impact will occur,
Likely	On balance the impact may occur
Unlikely	The possibility of this impact occurring is remote, however it must be considered

Table 7.2-2: Environmental and Social significance scale

Significance scale	Description
Very High	Major or permanent alteration of environmental or social dynamics, with severe or very severe consequences, or (in the case of benefits), beneficial or very beneficial effects.
High	Long term effect on the social or natural environment. This category should be treated with a significant degree of importance at the project decision making stage.
Moderate	Medium to long term effects on the social or natural environment. This category should also be taken into cognizance in decision making as constituting a fairly important degree of threat.
Low	These would have medium to short term ramifications on the social or natural environment; these are relatively unimportant and pose very little real threat.

Table 7.2-3: Impact Assessment Scale

Mitigation Potential	Impact Significance			
	Low	Moderate	High	Very High
Very Difficult	Medium	Major	Extreme	Extreme
Difficult	Minor	Medium	Major	Extreme
Achievable	Minor	Minor	Medium	Major
Easily Achievable	Minor	Minor	Minor	Medium

Table 7.2-4: Degree of Difficulty to Mitigate

Degree of Difficulty	Description
Very Difficult	The impact can be mitigated in theory, but the extent of financial or technical involvement militates against its application or effectiveness

Degree of Difficulty	Description
Difficult	The impact can be mitigated, but there is a significant degree of difficulty in implementing the proposed measures.
Achievable	The impact can be mitigated without much technicality or cost.
Easily Achievable	The impact can be easily and effectively mitigated

Table 7.2-5: Categories of Impact

Impact	Description
Extreme	Such impacts would prevent the action or option concerned from being taken or approved; and alternatives would have to be considered.
Major	These impacts are significant, meaning that if effective mitigation measures are not taken, a project may be hindered from commencing or continuing. Such option would require effective management and monitoring, or abandonment altogether for other options.
Medium	These impacts though important, are of less serious nature; in such a case, the Best Available Technology (or Practice) Not Entailing Excessive Cost (BATNEEC) should be employed. Such impacts alone are usually not significant enough to prevent a project from commencing or proceeding.
Minor	These impacts fall within the acceptable limits of the impact of a project on the environment, and mitigation is desirable but not necessary. This does not preclude 'Best Practice' as a means of avoiding cumulative impacts.

7.2.1.1 Land preparation and Construction Stage

Impacts at this stage are often temporary. The main concerns at the construction stage will be environmental and occupational health and safety. Risks would be reduced by strict adherence to best construction management practices. In relation to land preparation for nursery development, the main concerns are loss of biodiversity and erosion. The impacts anticipated their recommended mitigation measures and residual impacts are shown in the following table.

Table 7.2-6: Land Preparation and Construction Phase - Environmental and Mitigation Measures

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Loss of habitat	Clearing of vegetation for nursery development and construction	Clearing of vegetation in the concession area will result in loss of vegetation cover resulting in loss of habitat for various species. This may cause interruption of natural cycles including breeding and mating. Different species of flora may be lost which will not be reinstated when the area is reforested.	Certain	Moderate	Buffer zones will be established as well as botanical gardens/ ecological corridors. Although certain species lost during clearing may not be replaced, the cleared areas will eventually be re-vegetated with trees. A new thriving and balanced ecosystem will develop	Achievable	Medium
Loss of biodiversity	Clearing of vegetation for nursery development and construction	Clearing of vegetation, soil treatment and preparation before planting will cause local species within the project area to migrate to other areas, leaving the project site devoid of important species (mammals, reptiles, birds and insects) which contribute to the natural balance of the ecosystem	Certain	Moderate	Buffer zones will be established as well as botanical gardens/ ecological corridors. Some animals may return to the plantation areas, but natural processes may continue to be disrupted with operational activities including tree tending and treatment, harvest, etc.	Difficult	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
		within this area.					
Generation of Biomass	Clearing of vegetation for nursery development and construction	Vegetation clearing will result in the generation of biomass which would need to be disposed of safely	Certain	Moderate	Biomass will be used as far as possible as compost. It can also be used as firewood sources for local villages to cut and remove. Biomass will be reincorporated into the soil and turned into compost. It can also be used as firewood sources for local villages to cut and remove. Burning will only be resorted to as an absolute last option.	Achievable	Minor
Soil erosion	Site preparation for nursery development and construction	Exposure of loose soil resulting from vegetation clearing, soil preparation activities and pre-construction activities.	Certain	High	Implementation of erosion prevention measures in particular contour farming and cover cropping to minimise excessive runoff and erosion. The installation of sediment traps in drainage lines and storm water	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
					collection channels/settling ponds. Vegetation will be left around rivers and surface water bodies to minimise sediment entry. Fields will also be surrounded by vegetation to create buffers. The risk of soil erosion will reduce once planting commences; the same applies once construction commences and the soil is no longer exposed.		
Air Quality	Daily operations	Air quality deterioration due to dust emission resulting from the movement of vehicles along dirt roads, and fumes from machinery/equipment used in the development areas.	Certain	Moderate	Dust suppression measures will be employed such as site watering; regular maintenance of vehicles and machines to ensure that emissions are within normal operating limits	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Hydrology	Site preparation for nursery development and construction	Clearing of vegetation will result in an alteration of the natural hydrological cycle as evapotranspiration in the area will be reduced. Similarly without vegetation to break the volume of rainfall reaching the soils, there will be an increase in surface runoff which has direct impacts on soil erosion and sedimentation in water bodies.	Certain	High	This will be a short lived impact which will be reversed once the fruit trees begin to grow. Ground cover crops and mulching will be used to reduce the impact of the rainfall on the soil.	Achievable	Medium
Water quality	Site preparation for nursery development and construction	Surface water runoff from the site may contribute to the total suspended solids and turbidity of the receiving water bodies which may affect the aquatic life. Runoff may also carry off construction related debris, spills or waste not	Likely	High	Implementation of erosion prevention measures including ground cover (grass and other ground cover plants), the installation of sediment traps and storm water collection channels/settling ponds. Chemical spills in the field will not occur as	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
		properly handled.			all chemicals will be mixed at mixing stations which will be designed to contain spills during mixture and loading. Waste will be stored in secure waste receptacles where they will not be exposed to runoff.		
Noise pollution	Daily operations	Noise levels within the project area will increase particularly as a result of construction and other mechanised activities.	Certain	Moderate	<p>Modern equipment will be used which are fitted with noise suppression devices to prescribed standards.</p> <p>Noise generating agricultural activities such as land preparation (every 3 years), maintenance (fortnightly) and harvesting (every 12 – 14 months) will occur periodically and not on a daily basis.</p>	Difficult	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Waste Management	Daily activities	Poor waste management will result in an environmental health and safety hazard with effects on soil, air and water within the vicinity.	Certain	High	Implementation of an effective Waste Management System taking into account the various kinds of waste to be generated during this phase. Institution of monitoring system to ensure the efficacy of the handling, storage and disposal measures put in place	Achievable	Minor

Table 7.2-7: Land Preparation and Construction Phase - Social Impacts and Mitigation Measures

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Air Quality	Soil preparation and construction activities	Dust generating activities and movement of vehicles and machinery along dirt roads, as well as exhaust emissions from vehicles and machinery may have effects on the respiratory health of workers and nearby community	Certain	Moderate	Dust suppression measures will be employed including site watering; Regular maintenance of vehicles and machines to ensure that emissions are within normal operating limits	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
		residents.					
Noise pollution	Daily operations	Noise levels within the project area will increase particularly as a result of construction and other mechanised activities which may serve as a source of disturbance to nearby communities and effect workers directly operating noise generating machinery.	Certain	Moderate	<p>Modern equipment will be used which are fitted with noise suppression devices to prescribed standards.</p> <p>Noise generating agricultural activities such as land preparation (every 3 years), maintenance (fortnightly) and harvesting (every 12 – 14 months) will occur periodically and not on a daily basis.</p> <p>Workers will be provided with noise protection PPE.</p> <p>Construction and other activities will be restricted to the daylight hours in order to minimise effect on communities.</p>	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Waste Management	Daily activities	Poor waste management will result in an environmental health and safety hazards for workers and the communities with far reaching effects beyond the immediate waste storage or disposal areas.	Certain	High	Implementation of an effective Waste Management System taking into account the various kinds of waste to be generated during this phase. Institution of monitoring system to ensure the efficacy of the handling, storage and disposal measures put in place	Achievable	Minor
Pest infestation	Clearing of vegetation for nursery development and construction	Vegetation clearing will result in the generation of biomass which may present a haven for pests if not properly handled. Pests can wreak havoc on local communities through their destructive and disease carrying tendencies.	Likely	Moderate	Biomass will be used as far as possible. Biomass will be reincorporated into the soil and turned into compost. It can also be used as firewood sources for local villages to cut and remove, or burnt if no other options for reuse or safe disposal are available	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Loss of land	Project development	Loss of land may be an issue even after land lease agreements have been reached.	Likely	High	<p>Intensive community consultations and additional sensitization meetings will be conducted prior to the commencement of this stage of the project to ensure that landholders' views and opinions are heard and taken into consideration. Locals will be made aware of the implications and potential impact of a large project starting in their communities, including the pitfall of giving away too much land.</p> <p>Land will be leased in phases as the project progresses in order to prevent leased land remaining unused over an extended period.</p>	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Visual Impact	Clearing of vegetation for nursery development and construction	There will be a sharp contrast between areas cleared for project development and the rest of the surroundings; this may create some apprehension or cause for concern in residents.	Likely	Low	The cleared areas would soon be replaced with trees and modern structures. However, this may still be cause for concern to some residents. During initial sensitization meetings conducted pictures of the Philippines sister operation were shown to the residents to give them an idea of the buildings and fields in a large agricultural operation.	Difficult	Minor
Occupational Health and Safety	Daily activities	Workers will be at risk of injury from the various machinery, equipment, processes and operational methods; this risk is particularly heightened in the case of locally employed workers who may not have been previously exposed to the working conditions.	Likely	Moderate	STL intends to maintain the safety of its workers through the following means: 1. Use of modern, low risk equipment 2. Training to ensure that only qualified and licenced operators are allowed to operate	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
					machinery and equipment. 3. Provision of PPE 4. Conducting risk assessments 5. Maintenance of machinery and equipment 6. Implementation of Management and Monitoring Plans.		
Community Health and Safety	Daily Activities	Community residents will be exposed to increased vehicular traffic and other operational activities which being unfamiliar puts them at risk of safety hazards.	Likely	High	Regular community consultations and community health and safety sensitization will be conducted periodically throughout the life of the project.	Achievable	Moderate

7.2.1.2 Operational Stage

Once best practices have been observed during the initial stages of the project - planning/design set up and construction stages - much of the threat to the safety and integrity of the environment and society will be reduced to levels defined by legislation and best practices.

Table 7.2-8: Operational Phase – Environmental Impacts and Mitigation Measures

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Loss of habitat	Clearing of vegetation for progressive expansion of plantation areas	Clearing of vegetation in the concession area will result in loss of vegetation cover resulting in loss of habitat for various species. This may cause interruption of natural cycles including breeding and mating. Different species of flora may be lost which will not be reinstated when the area is reforested with Fruit Tree.	Certain	Moderate	Buffer zones will be established as well as botanical gardens/ ecological corridors and environmental monitoring plots also established to monitor changes. Although certain species lost during clearing may not be replaced, the cleared areas will eventually be re-vegetated with fruit trees. A new thriving and balanced ecosystem will develop	Achievable	Minor
Loss of biodiversity	Clearing of vegetation for progressive expansion of plantation areas	Clearing of vegetation, soil treatment and preparation before planting will cause local species within the project area to migrate to other	Certain	Moderate	Buffer zones will be established as well as botanical gardens/ ecological corridors. Environmental monitoring plots also	Difficult	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
		areas, leaving the project site devoid of important species (mammals, reptiles, birds and insects) which contribute to the natural balance of the ecosystem within this area.			established to monitor changes in the area. Some animals may return to the plantation areas, but natural processes may continue to be disrupted with operational activities including tree tending and treatment, harvest, etc.		
Generation of Biomass	Clearing of vegetation during plantation expansion, pruning, etc.	Vegetation clearing will result in the generation of biomass which would need to be disposed of safely	Certain	Moderate	Biomass will be used as far as possible as compost. It may also be made available as a source of firewood for community members. Burning will only be resorted to as an absolute last option.	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Soil erosion	Clearing of vegetation for progressive expansion of plantation areas	Exposure of loose soil resulting from vegetation clearing and soil preparation activities	Certain	High	<p>Implementation of erosion prevention measures including in particular contour farming and cover cropping to minimise excessive runoff and erosion, the installation of sediment traps and storm water collection channels/settling ponds.</p> <p>Vegetation will be left around rivers and surface water bodies to minimise sediment entry.</p> <p>The risk of soil erosion will reduce once planting commences; the same obtains once construction commences and the soil is no longer exposed.</p>	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Soil acidification	Use of fertilizers and pesticides	Soil acidification is accelerated by the use of fertilizers containing ammonium salts or urea. Acidic soil can restrict the roots of plants' ability to reach nutrients and water.	Likely	Moderate	Natural fertilizers and compost will be employed as far as possible; only approved and non-persistent chemicals will be used where necessary. Regular soil testing and analysis will be carried out to monitor chemical changes in the soil.	Achievable	Minor
Air Quality	Daily operations	Air quality deterioration due to dust generation resulting from the movement of vehicles along dirt roads, and fumes from machinery/equipment	Certain	Moderate	Dust suppression measures will be employed including site watering; Regular maintenance of vehicles and machines to ensure that emissions are within normal operating limits	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Hydrology	Irrigation	Water abstraction from the Sewa River could affect the availability and hydrological regime within the project area.	Certain	Moderate	<p>Agricultural practices such as no till or the use of cover crops/mulch and the incorporation of organic matter from plant residues are all Good Agricultural practices which will mitigate excessive irrigation requirements.</p> <p>Pineapple is a drought tolerant plant so can withstand some water stress.</p> <p>Alternative water sources will be made available for the project and communities through the sinking of boreholes. These will supplement and reduce the water obtained from the Sewa River.</p>	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Water quality	Use of Fertilisers and Pesticides	Water pollution can occur due to the use of agro-chemicals such as fertilizers and pest control substances which can leach into the soil and get carried into water by runoff. Aquatic life would be affected.	Certain	High	Precision Agriculture methods will be employed with regard to crop nutrition, and fertilizer will only be applied in the required amounts foliarly so contamination of surface water is greatly reduced. Chemicals i.e. Herbicides and pesticides will be used only as required or when particular pest thresholds are reached.	Achievable	Medium
Noise pollution	Daily operations	Elevated noise levels will result from movement and operation of vehicles and machinery.	Certain	Moderate	Modern equipment will be used which are fitted with noise suppression devices to prescribed standards.	Difficult	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
					Noise generating agricultural activities such as land preparation (every 3 years), maintenance (fortnightly) and harvesting (every 12 – 14 months) will occur periodically and not on a daily basis. Workers will be provided with noise protection PPE.		
Waste Management	Daily activities	Poor waste management will result in an environmental health and safety hazard with effects on soil, air and water within the vicinity.	Certain	High	Implementation of an effective Waste Management System taking into account the various kinds of waste to be generated during this phase. Institution of	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
					monitoring system to ensure the efficacy of the handling, storage and disposal measures put in place		
Emergency Response and Disaster Management	Throughout project life	Poor management of emergencies or disasters will result in loss of life, damage to equipment and disruption of operations	Likely	Very high	Implementation of Emergency Response Plan, regular drills and training exercises for staff, monitoring of potential situations leading to disaster.	Achievable	Medium

Table 7.2-9: Operations Phase - Social Impacts and Mitigation Measures

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Air Quality	Daily operations	Dust generating activities and movement of vehicles and machinery along dirt roads, as well as exhaust emissions from vehicles	Certain	Moderate	Workers are provided with dust protection PPE. Dust suppression measures will be employed including	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
		and machinery may have effects on the respiratory health of workers and nearby community residents.			site watering; Regular maintenance of vehicles and machines to ensure that emissions are within normal operating limits		
Noise pollution	Daily operations	High noise levels as a result of vehicular and machinery movement may serve as a source of disturbance to nearby communities and affect workers directly operating noise generating machinery.	Certain	Moderate	Modern equipment will be used which are fitted with noise suppression devices to prescribed standards. Noise generating agricultural activities such as land preparation (every 3 years), maintenance (fortnightly) and harvesting (every 12 – 14 months) will occur periodically and not on a daily basis. Workers will be provided with noise protection PPE.	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Waste Management	Daily activities	Poor waste management will result in an environmental health and safety hazards for workers and the communities with far reaching effects beyond the immediate waste storage or disposal areas.	Certain	High	Implementation of an effective Waste Management System taking into account the various kinds of waste to be generated during this phase. Institution of monitoring system to ensure the efficacy of the handling, storage and disposal measures put in place	Achievable	Minor
Pest infestation	Clearing of vegetation for nursery development and construction	Vegetation clearing will result in the generation of biomass which may present a haven for pests if not properly handled. Pests can wreak havoc on local communities through their destructive and disease carrying tendencies.	Likely	Moderate	Biomass will be used as far as possible as compost. It may also be made available as a source of firewood for community members. Burning will only be resorted to as an absolute last option.	Achievable	Minor
Occupational Health and Safety	Daily activities	Workers will be at risk of injury from the various machinery, equipment, processes and operational methods; this risk is particularly	Likely	Moderate	STL intends to maintain the safety of its workers through the following means: 1. On the job training	Achievable	Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
		heightened in the case of locally employed workers who may not have been previously exposed to the working conditions.			2. Provision of PPE 3. Conducting risk assessments 4. Maintenance of machinery and equipment 5. Implementation of Management and Monitoring Plans.		
Community Health and Safety	Daily Activities	Community residents will be exposed to increased vehicular traffic and other operational activities which being unfamiliar puts them at risk of safety hazards.	Likely	High	Regular community consultations and community health and safety sensitization will be conducted periodically throughout the life of the project.	Achievable	Moderate

7.2.1.3 Termination/Decommissioning Stage

It is now common practice that closure operations are integrated into the planning/design, and operational stages, as the environmental gains of implementing certain mitigation measures may take much longer to be realized. This approach reduces the burden and cost of an end of project rehabilitation, reclamation or clean-up. Closure activities are generally performed to stabilize the site, and remove fixed and moveable surface and sub-surface structures.

Table 7.2-10: Decommissioning Phase – Environmental and social impacts

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Air Quality	Demolition	Air quality deterioration due to dust generation from demolition and related activities.	Certain	Moderate	<p>This will be temporary, and will normalise following completion of closure activities.</p> <p>Post closure monitoring will be implemented for at least 3 years following closure.</p> <p>Dust suppression measures will be employed including site watering; workers will be provided with respiratory PPE.</p>	Achievable	Minor
Hydrology	Post Closure	The pre-project hydrological regime of the Sewa River will be restored over time with water no longer being abstracted for irrigation and other purposes.	Certain		Implementation of post closure monitoring		Positive

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Water quality	Post closure	The risk of pollution from agrochemicals will be eliminated following closure.	Certain	High	Closure activities will include taking water quality tests on surface water bodies and implementing treatment if required. Post closure monitoring will be carried out for at least 3 years.	Achievable	Positive
Noise pollution	Demolition and other closure activities	Elevated noise levels will result from movement and operation of vehicles and machinery during closure.	Certain	Moderate	Temporary impact; following closure, elevated noise levels resulting from project activities will be reversed. Vehicles and machinery will be fitted with silencers or mufflers to help minimize the noise generated. PPE will be provided to workers.		Minor

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Waste Management	Closure	Poor waste management will result in an environmental and community health and safety hazards with effects on soil, air and water within the vicinity.	Certain	High	<p>Implementation of an effective Waste Management System taking into account the various kinds of waste to be generated during this phase (construction waste – from demolished structures, domestic waste, sewage, etc)</p> <p>Soil tests will be taken from storage and disposal areas in order to treat any contamination.</p> <p>Post closure monitoring will include checks and tests in waste disposal/storage areas.</p>	Achievable	Medium

Issue	Location/ Project Activity	Impact	Certainty of impact	Significance	Mitigation measure	Degree of difficulty	Impact Category
Occupational Health and Safety	Demolition and other closure activities	Workers will be at risk of injury from the various machinery, equipment and processes related to closure.	Likely	Moderate	Professional and experienced decommissioning contractors will be contracted for this phase.	Achievable	Minor
Community Consultations	Before and during closure phase	Lack of or poor community consultations may result in the establishment of a post closure land use not beneficial to or appreciated by the communities	Likely	High	Community consultations will be conducted throughout the life of the project; specific closure meetings will be held to obtain views and perspectives of residents in determining the post closure land use.	Achievable	Minor

8 CONCLUSION

8.1 Summary

8.1.1 Components of the ESIA

The principal objective of the ESIA is two-fold: the first objective is to satisfy the requirements of the local environmental regulatory body, EPA-SL for the issuance of the EIA license for the project to commence; the second objective is to adhere to best practice for carrying out an impact assessment for an agricultural/agro-processing project. The study involved predicting the environmental impacts of the project as described, and suggesting mitigation measures where impacts are adverse and enhancement measures where impacts are positive.

The ESIA report comprised:

- i. Compilation of an Environmental and Social Impact Assessment (ESIA) based on the primary and secondary information obtained through field and desk studies;
- ii. Development of Environmental and Social Management Plans based on the proposed impact mitigation measures.
- iii. Formulation of a Community Development Action Plan (CDAP) from socioeconomic analysis and stakeholder discussions;
- iv. Description of the stakeholder, interested and affected parties engagement process in a Public Consultation and Disclosure Plan;

The investigations of impacts on the social environment are a crucial part of the study, since the operation will affect the livelihoods of the communities that reside in and around the project area. The investigation of social impacts has involved the following:

- i. A baseline socio-economic study of communities surrounding in and around the project area;
- ii. Undertaking stakeholders' focus group discussions to sensitise stakeholders and Project Affected Persons (PAPs) on the Project.

Discussions and meetings with stakeholders during the public consultation and disclosure process indicated general acceptability for the project with demonstrated enthusiasm at the local level. Local authorities within the project area expressed their opinions, concerns and general willingness for full co-operation and support during the survey and these were evident during the field investigations.

8.1.2 Key Assessment Findings

8.1.2.1 Physical Environment

There are potential major impacts relating to hydrology and water quality, biodiversity, soil erosion, water & soil pollution due to agro-chemical and water abstraction. Mitigation measures to limit the extent of impacts have been highlighted and will be implemented.

8.1.2.2 Biological Environment

There will be some loss of floral and faunal species during land clearing and preparation, however it is not expected that any species will be wiped out from the region as a result. Mitigation measures have been presented to ensure that minimal clearing is carried out to limit the extent of biodiversity loss.

8.1.2.3 Socio-economic Environment

Perhaps the most critical aspect of the project is the potential reduction of farm land by the communities, potential conflict from issues related to labour, unrealistic expectations held by the communities with regard to benefits created by the project, vehicular traffic and safety risk. The selection of the project site has been judiciously done to avoid the need for community relocation.

The project is will have positive impacts in the area of job creation, improving the quality of life of locals through payment of land lease rent and compensation, outgrower schemes, creation of secondary businesses, etc.

8.2 Conclusion

This Environmental and Social Impact Assessment has been professionally carried out to satisfy the Government of Sierra Leone's legislation and regulations. The study has achieved the following objectives for such an exercise, viz:

- Baseline Survey of the Project Area;
- Execution of an Environmental and Social Impact Assessment (ESIA) and development of Environmental and Social Management Plans (ESMP);

The investigations of impacts on the social environment are a crucial part of the study, since the operation may impact the communities which reside at close proximity to the project site.

The investigation of social impacts has involved the following:

- A baseline socio-economic study of the community envisaged to be impacted by the project activities in both the construction and operational phases;
- A Public Consultation and Disclosure Process (PCDP) undertaken to sensitize stakeholders.

The baseline environmental and socio-economic situations have been presented.

Environmental impacts of the project's activities have been identified for all phases of the project. The most significant of these impacts is the change of morphology and soil and water (surface and ground) pollution due to agro-chemicals utilization. Social impacts include loss of agricultural farm land and unrealistic expectation from the people regarding benefits as the project progresses.

Generally, the investigations reveal that environmental and social problems incurred by the project can be adequately managed and that there are no insurmountable problems that should stop the project from proceeding.

A monitoring system must however be put in place to ensure that management practices for mitigating negative impacts and enhancing those that are positive are affected. It must however be ensured that recommendations made in the Environmental and Social Management Plans are followed through.

9 REFERENCES

- Alatec, S.A., Valenciaport (Alatec-Valenciaport) and CEMMATS Group Ltd, (2008). *Environmental Audit for Freetown Port*. Freetown.
- Altieri, M. A. (1995). *Agroecology: the science of sustainable agriculture*. Boulder:
- Asamoah Larbi (2012). Country Pasture/Forage Resource Profile: Sierra Leone. [online] Available at <http://www.fao.org/ag/agp/agpc/doc/counprof/Sierraleone/Sierraleone.htm> [Accessed 20th September 2016]
- Beier, P., Loe, S. (1992). "In My Experience: A Checklist for Evaluating Impacts to Wildlife Movement Corridors. *Wildlife Society Bulletin*". *Wildlife Society Bulletin*. Allen Press. **20** (4): 434–440.
- Birchall, C.J., Bleeker, P. & Cusani-Visconti, C. (1979). Land in Sierra Leone: A Reconnaissance Survey and Evaluation for Agriculture. Technical Report No. 1 AG: DP/SIL/73/002. UNDP/FAO 1979.
- CEMMATS Group Ltd, (2016). *Freetown Terminal Limited ESIA Scoping Report*. Freetown
- Gleave, M.B. (1996). The Length of the Fallow Period in Tropical Fallow Farming Systems: a discussion with evidence from *Sierra Leone*. *The Geographical Journal* Vol. 162, No. 1 (Mar., 1996), pp. 14-24.
- Ighobor Kingsley, (2014). *Sierra Leone: Nursing Agriculture Back To Health*. [online] Available at <http://www.un.org/africarenewal/magazine/special-edition-agriculture-2014/sierra-leone-nursing-agriculture-back-health>. [Accessed 20th September 2016]
- IUCN. (2016). *IUCN Red List of Threatened Species*. Gland, Switzerland www.iucn.org
- Kingdon, J. (1997). *The Kingdon field guide to African mammals*. Academic Press, San Diego.
- Okoni-Williams, A.D. (2013). Vegetation, carbon and nutrient cycling in the bush fallow system: the implications to the environment and local farming communities in Sierra Leone. PhD Thesis, University of Sierra Leone.
- Shafagoj, J. A. A (2015). *Logistics Capacity Assessment*. [Online] Available at <http://dlca.logcluster.org/display/public/DLCA/2.1+Sierra+Leone+Port+of+Freetown;jsessionid=645327F2D09307EF7D38F4526461E081> [Accessed 7th July 2016]
- Sierra Leone 2015 Population And Housing Census (2016). [online] Available at <https://www.statistics.sl/wp-content/uploads/2016/06/2015-Census-Provisional-Result.pdf> [Accessed 10th October 2016]

UN Office for the Coordination of Humanitarian Affairs (2016). *Sierra Leone: Bo District Profile (04 December 2015)*. [online] Available at <http://reliefweb.int/report/sierra-leone/sierra-leone-bo-district-profile-04-december-2015> [Accessed 28/09/16]

UNDP (2016). *Sierra Leone makes progress in human development, but poverty and inequality persist*. [online] Available at <http://www.sl.undp.org/content/sierraleone/en/home/presscenter/pressreleases/2016/03/10/sierra-leone-makes-steady-progress-in-human-development-despite-ebola-but-deep-poverty-and-inequality-persist/> [Accessed 27/09/16]

10 APPENDICES

ANNEX A: MINUTES OF FOCUS GROUP DISCUSSION MEETINGS

Sierra Tropical Focus Group Minutes in selected villages Problems listed by the group

1. KPATEMA VILLAGE

April 11,

2016

Problems of the village as indicated by the group

1. No school
2. No assembling hall
3. Water problem, bore hole well dries up in the dry season
4. No building for women society

Solution: Need help from the project in collaboration with other relevant NGOs

Livelihood earnings are from Rice farming and small scale diamond mining, but farms are small and alluvial diamond mining is not profitable

End result - Abject poverty

Residents are aware of STL project and look at it for the solution of their problems

Severe water shortage negatively affects palm oil production and the coloration of the oil.

Lack of water is the problem.

Dirty water affects food preparation, bathing and washing of clothes

Preferences for help in order of priority

1. Water
2. School
3. Church
4. Assembly hall
5. Public toilets
6. Housing

2. MOMADU VILLAGE

Similar problems

Lack – Schools

- Water
- Clinics
- Court Barry

The village has 14 houses with a population of 100 people.

Main occupation is cassava and rice farming

Extremely poor village

3. KOMEDE VILLAGE

Problems of village

1. No health facility
2. No school
3. Poor road to village
4. No public toilets
5. No equipment for young people

4. MOSHOGBO VILLAGE

The village has similar needs as above.

Problems

1. No clinic
2. No school
3. No market
4. No public toilets and shops
5. Housing conditions are bad

5. SUMBUYA TOWN

Sumbuya is the head quarter town with limited facilities and also faced with the following problems.

- Reliable Water supply system
- Electricity generation
- Needs a Petrol station
- Large pool of unemployed youths anxiously awaiting to be engaged
- Poor quality of houses

- No Vocational training facilities
- It has a secondary school but poorly equipped
- No Public toilets

In almost every village, the residents are crying for the items listed above and they look for help not only from the project but someone who can associate with NGOs and the government to look into their problems.

ANNEX B: ATTENDANCE LIST FOR FOCUS GROUP DISCUSSIONS

ANNEX C: ECOLOGICAL SPECIES RECORDED

Tree species recorded

Botanical name	Family	IUCN Status	Local status	Uses
<i>Afzelia africana</i>	Caesalpiniaceae	VU	Rare	Valuable timber
<i>Albizia adianthifolia</i>	Mimosaceae	LC	Common	
<i>Albizia zygia</i>	Mimosaceae	LC	Not Common	
<i>Allophylus africanus</i>	Sapindaceae	LC	Common	
<i>Alstonia congensis</i>	Apocynaceae	LC	Common	
<i>Amphimas pterocarpoides</i>	Caesalpiniaceae	LC	Common	
<i>Anisophyllea laurina</i>	Anisophylleaceae	LC	Common	Building poles
<i>Anthocleista nobilis</i>	Gentianaceae	LC	Common	
<i>Anthonota macrophylla</i>	Caesalpiniaceae	LC	Common	
<i>Bombax buonopozense</i>	Bombacaceae	LC	Common	
<i>Canarium schweinfurthii</i>	Burseraceae	LC	Common	
<i>Carapa procera</i>	Meliaceae	LC	Common	
<i>Ceiba pentandra</i>	Bombacaceae	LC	Common	Timber
<i>Chidlowia sanguinea</i>	Caesalpiniaceae	LC	Common	
<i>Cleistopholis patens</i>	Annonaceae	LC	Common	
<i>Cola lateritia</i>	Sterculiaceae	LC	Not Common	
<i>Cola nitida</i>	Sterculiaceae	LC	Common	
<i>Dialum dinklagei</i>	Caesalpiniaceae	LC	Not Common	
<i>Dialum guineense</i>	Caesalpiniaceae	LC	Common	Timber
<i>Ficus exasperate</i>	Moraceae	LC	Common	Timber
<i>Ficus mucoso</i>	Moraceae	LC	Common	
<i>Funtumia africana</i>	Apocynaceae	LC	Not Common	
<i>Garcinia afzelii</i>	Guttiferae	VU	Rare	Valuable timber
<i>Gmelina arborea</i>	Lamiaceae	LC	Common	Timber
<i>Heritiera utilis</i>	Sterculiaceae	VU	Rare	Valuable timber
<i>Hymenocardia lyrata</i>	Euphorbiaceae	LC	Common	
<i>Lophira alata</i>	Ochnaceae	VU	Rare	Valuable timber
<i>Macaranga bati</i>	Euphorbiaceae	LC	Common	
<i>Macaranga heterophylla</i>	Euphorbiaceae	LC	Common	
<i>Milicia regia</i>	Moraceae	VU	Rare	Valuable timber
<i>Morinda geminata</i>	Rubiaceae	LC	Not Common	
<i>Musanga cecropioides</i>	Moraceae	LC	Common	
<i>Myrianthus arboreus</i>	Moraceae	LC	Common	
<i>Newtonia aubrevillei</i>	Mimosaceae	LC	Common	
<i>Parinari excelsa</i>	Chrysobalanaceae	LC	Not Common	Timber
<i>Parkia bicolor</i>	Mimosaceae	LC	Not Common	
<i>Pentadesma butyracea</i>	Guttiferae	LC	Common	
<i>Pentaclethra macrophylla</i>	Mimosaceae	LC	Not Common	
<i>Phyllacasmus africanus</i>	Ixonanthaceae	LC	Common	Timber
<i>Phyllanthus discoideus</i>	Euphorbiaceae	LC	Common	

<i>Piptadeniastrum africanum</i>	Mimosaceae	LC	Common	
<i>Samanea dinklagei</i>	Mimosaceae	LC	Common	
<i>Spondias mombin</i>	Anacardiaceae	LC	Common	
<i>Sterculia tragacantha</i>	Sterculiaceae	LC	Common	
<i>Synsepalum afzelii</i>	Sapotaceae	LC	Common	
<i>Synsepalum brevipes</i>	Sapotaceae	LC	Common	
<i>Synsepalum sp</i>	Sapotaceae	LC	Common	
<i>Tarenna vignei</i>	Rubiaceae	LC	Common	
<i>Terminalia ivorensis</i>	Combretaceae	VU	Rare	Valuable timber
<i>Trichoscypha arborea</i>	Anacardiaceae	LC	Common	
<i>Trichoscypha bijuga</i>	Anacardiaceae	LC	Common	
<i>Uapaca guineensis</i>	Euphorbiaceae	LC	Common	Timber
<i>Vitex micrantha</i>	Verbenaceae	LC	Common	
<i>Xylopia aethiopica</i>	Annoceae	LC	Not Common	

Mammals species that are known to occur within project concession zone

Species	Scientific name	IUCN Status	Local Status	Site 1	Site 2	Site 3	Site 4	Site 5	Site 6	Site 7	Site 8	Site 9	Site 10	Site 11	Site 12	Site 13	Site 14
Chimpanzee	<i>Pan troglodytes verus</i>	EN	VR		x			x	x			x		x		x	
Red colobus Monkey	<i>Procolobus badius</i>	EN	R		x					x	x	x	x		x	x	
Olive Colobus	<i>Poliocolobus verus</i>	LC	R														
Pied Colobus Monkey	<i>Colomus polykomus</i>	VU	NC	x	x	x	x	x	x			x	x	x		x	x
Sooty Mangabay	<i>Cercocebus atys</i>	VU	NC	x	x		x		x				x	x			
Campbell's Monkey	<i>Cercopithecus campbelli</i>	LC	NC							x		x			x	x	x
Maxwel Duiker	<i>Cephalophus maxwelli</i>	LC	C														
Zebra Duiker	<i>Cephalophus zebra</i>	VU	VR				x		x			x	x	x		x	
Red-flanked Duiker	<i>Cephalophus rufilatus</i>	LC	NC	x	x	x	x	x		x	x	x	x				
Bongo	<i>Tragelaphus euryceros</i>	NT	VR	x	x	x	x	x		x	x	x	x				
Water Chevrotain	<i>Hyemoschus aquaticus</i>	DD	NC		x	x	x	x	x		x	x			x	x	
Forest Buffalo	<i>Syncerus caffer</i>	LC	NC	x	x		x	x	x	x	x	x	x	x	x	x	
Bush Pig	<i>Potamochoerus larvatus</i>	LC	C	x	x	x	x	x	x	x	x	x	x	x	x	x	
Marsh Cane Rat	<i>Thryonomys swinderianus</i>	LC	VC	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Fire-footed Rope Squirrels	<i>Funisciurus pyrropus</i>	LC	VC	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Gambian Sun squirrel	<i>Heliosciurus gambianus</i>	LC	C	x		x	x		x	x	x	x	x		x	x	
Brush-tailed Purcupine	<i>Atherurus africanus</i>	LC	NC	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Crested Purcupine	<i>Hystrix cristata</i>	LC	NC	x	x			x	x				x		x		x
Giant Pouch Rat	<i>Cricetomys gambianus</i>	LC	VC	x	x	x	x	x	x	x	x	x	x	x	x	x	x
African Civet	<i>Civettictis civetta</i>	LC	NC	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Tree Pangolins	<i>Phataginus tricuspis</i>	VU	NC	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Scrub Rabbit	<i>Lepus saxatilis</i>	LC	NC													x	x
Common Cusimanse	<i>Crossarchus obscurus</i>	LC	NC		x			x		x	x		x	x	x	x	
Common genet	<i>Genetta genetta</i>	LC	NC	x	x		x		x			x	x			x	x

